

Geology of Hvitland beds (late Pliocene)White Point Lowland, Ellesmere Island, Northwest Territories. by J.G. Fyles [and others]

Natural Resources Canada - Advances in Quaternary Entomology



Description: -

-
Geology - Northwest Territories - Ellesmere Island
Paleoclimatology - Northwest Territories - Ellesmere Island
Paleontology - PlioceneGeology of Hvitland beds (late Pliocene)White Point Lowland, Ellesmere Island, Northwest Territories. by J.G. Fyles [and others]
-Geology of Hvitland beds (late Pliocene)White Point Lowland, Ellesmere Island, Northwest Territories. by J.G. Fyles [and others]
Notes: 11
This edition was published in -



Filesize: 13.102 MB

Tags: #northwest #territories #arctic: #Topics #by #Science.gov

Paleoenvironments of the Canadian high arctic derived from pollen and plant macrofossils: Problems and potentials

Rainfall as a determinant of soil moisture. Oxygen isotope ratio: the ratio of oxygen-18 to oxygen-16 isotopes in oxygen-bearing geologic materials such as carbonate shells of marine organisms, used as a measure of past temperatures. Results are now available of four summers and two winters of standard meteorological observations, and global solar and net radiation measurements as well.

Pliocene terrace gravels of the ancestral Yukon River near Circle, Alaska: Palynology, paleobotany, paleoenvironmental reconstruction and regional correlation

A Frontoclypeus of Agrypnites Phryganeidae; B frontoclypeus of Asynarchus Linnephilidae; C frontoclypeus of Grensia Linnephilidae; D frontoclypeus of Linnephilus; E frontoclypeus of Agrypnia Phryganeidae; F drawing of caddis fly larval frontoclypeus, showing principal diagnostic features photos by the author.

terrestrial paleoclimate record: Topics by Science.gov

General circulation model GCM simulations of future climatic changes, which may be expected with higher levels of carbon dioxide CO2 and other greenhouse gases, show a remarkable consensus that the effects of anthropogenic greenhouse warming will be particularly intense in polar regions e.

A Synthesis of the Long

Salix rotundifolia populations growing adjacent to the Cape Simpson natural oil seep were examined for possible changes which may have occurred as the result of long term exposure to oil. Kettlehole: a steep-sided basin, commonly without surface drainage, in glacial drift deposits especially outwash, formed by the melting of a large, detached block of stagnant ice. Murray Mitchell in Alaska Mitchell, 1956.

Long

Contribution of working group I to the third assessment report of the Intergovernmental Panel on Climate Change. In four populations of polar bears in the eastern Canadian Arctic including Western Hudson Bay , Inuit hunters report more bears near settlements during the open water period in recent years.

History of insects

Zur Frage der Entstehung von Hefeiweiss aus anorganischen Stickstoffverbindungen. This data is primarily from historical hydrographic surveys that were carried out by the Canadian Hydrographic Survey CHS. *Sphaerotheca fuliginea*, which, like most powdery mildews, demands a relatively dry climate, produced ascocarps profusely on several dicotyledons at Hazen Camp, maturing them by late July in the warm season of 1962, about 50 days from the initiation of growth.

Related Books

- [Wuthering Heights \(Emily Brontë\).](#)
- [Yahadut Gruzyah 'al parashat derakhim - \(sikun-benayim shel he-'aśor ha-ḥolef\) 1970-1980](#)
- [Instructors manual for intermediate algebra](#)
- [Tidewrack - neglected facts and intriguing stories from the age of the wooden ships](#)
- [Salop directory 1818](#)