

# Coronary artery diseases - diagnostic and therapeutic imaging approaches

Springer-Verlag - 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes



Description: -

-

Coronary Vessels -- radionuclide imaging.

Coronary Vessels -- radiography.

Coronary Disease -- therapy.

Heart -- Radionuclide imaging.

Coronary heart disease -- Interventional radiology. Coronary artery diseases - diagnostic and therapeutic imaging approaches

- Coronary artery diseases - diagnostic and therapeutic imaging approaches

Notes: Includes bibliographies and index.

This edition was published in 1984



Filesize: 32.74 MB

Tags: #Intravascular #NIRF #Molecular #Imaging #Approaches #in #Coronary #Artery #Disease

## Coronary artery disease in dialysis patients: evidence synthesis, controversies and proposed management strategies

This also explains the persistent use of methods that are entitled to reimbursement despite recognised significant disadvantages and inferior evidence — such as nuclear medicine methods — and are simply underpowered to deliver the insights to a complex pathophysiology, for example, serial echocardiography, cardiac CT or invasive catheterisation.

## Coronary Artery Diseases, Radiology

They used a 20-segment model that divided the myocardium into regions based on the apical, mid and basal cuts on short axis views, and the apical region of the parasternal long axis view. Lung examination may reveal or exclude a pulmonary cause for this presentation wheezes, egophony, absent or asymmetric breath sounds.

## Coronary Artery Disease: Causes, Symptoms, Diagnosis & Treatments

This rather reserved status of cardiac MRI continues to reflect an overall poor availability of CMR in routine practice and there are several aspects responsible for this.

## Effect of Coronary Anatomy and Myocardial Ischemia on Long

However, specificity for the presence of plaque and stenosis for ACS was 54 and 87%, respectively. In the SL group the origin from the LCA was more frequent 80% versus 62%, the tortuosity of the fistulous vessels was more prevalent, and aneurysmal formation was more common than in the PTE group.

## Scientists identify a new diagnostic and therapeutic target for cardiovascular disease

Rest 12-lead ECG was performed in all patients.

## Related Books

- [Nariño, Torres y la revolución francesa](#)
- [Hair sheep of western Africa and the Americas - a genetic resource for the tropics](#)
- [Indian tribes of Alberta](#)
- [Chien jaune](#)
- [Katharina II., die Selbstherrscherin aller Reussen - das Bild der Zarin und ihrer Aussenpolitik in d](#)