

Hémiptères Saldidae et Leptopodidae - dEurope occidentale et du Maghreb

Fédération française des sociétés de sciences naturelles - New shore bug (Hemiptera, Heteroptera, Saldidae) from the Early Cretaceous of China with phylogenetic analyses

Description: -

-
Hockey players -- Czechoslovakia -- Biography.
Hockey -- Czechoslovakia -- History.
United States -- Claims
Bills, Private -- United States
United States. -- Congress -- Private bills
Leptopodidae -- Africa, North.
Shore bugs -- Africa, North.
Leptopodidae -- Europe.
Shore bugs -- Europe. Hémiptères Saldidae et Leptopodidae -
dEurope occidentale et du Maghreb

-
77

Faune de France ;Hémiptères Saldidae et Leptopodidae - dEurope occidentale et du Maghreb

Notes: Includes bibliographical references (p. [211]-226) and indexes.

This edition was published in 1990



Filesize: 14.42 MB

Tags: #A #review #of #species #of #the #genus #Teloleuca #(Heteroptera, #Saldidae) #in #the #fauna #of #Russia #and #adjacent #territories

literatur

Michael Bonk EDV Wolfgang Hanns Laura C. Сложные глаза крупные, выступающие, расположены в заднебоковых углах головы и могут нести как прямые, так и изогнутые щетинки. Data on the distribution of eight species S.

Faune de France, Volume 86: Hyménoptères Sphecidae d'Europe Occidentale Volume 3

В ископаемом состоянии антенны иногда сохраняются достаточно хорошо, однако чаще отдельные членики бывают плохо различимы.

Faune de France (collection d'ouvrages) — Wikipédia

Le contenu est disponible sous licence sauf mention contraire. The body length was measured from the apex of head to the apex of abdomen; body width, at the maximal width of body; pronotum length, along the midline; pronotum width, across the broadest part at its posterior angles; wing length, from the basal to the apex of anterior margin; wing width, at the maximal width of the wing.

literatur

A comprehensive presentation of all 152 species of Pentatomoidea Pentatomidae, Scutelleridae, Cydnidae, Acanthosomatidae, Thyreocoridae, Plataspididae present in France. Pacific Insects 11 3—4 : 571-578. В ископаемом состоянии обычно хорошо различима передняя доля пронотума, иногда сохраняющая рельеф и окраску.

literatur

In Saldinae, base of penis-filum like a watch-spring. A new genus with a new species of Saldidae, Brevimatus pulchaler gen. Kopana

университет Рединга, Англия и И.

Faune n° 77

Некоторые авторы Рыжкова, 2012; Hong, Wang, 1990; Zhang, 1991; Hong, Ren, 1992; Hong, 1995 предлагают также синонимизировать *Enicocoris* Popov, 1980 с *Mesolygaeus* Ping, 1928, описанным в составе семейства Lygaeidae Ping, 1928. Trogositidae, Byturidae, Rhizophagidae, Cucujidae, Erotylidae, Phalacridae, Mycetophagidae, Colydiidae, Endomychidae, Coccinellidae, Sphindidae. Illustrated Catalogue of the American Fauna: Fulgoridae I.

CAB Direct

The semiaquatic and aquatic Hemiptera of California Heteroptera: Hemiptera. Zoological Journal of the Linnean Society 80: 231-238.

Related Books

- [We remember Bataan and Corregidor - the story of the American & Filipino defenders of Bataan and Cor](#)
- [Readings in poetry](#)
- [Historians and the American West](#)
- [Foreign ownership and the auto parts industry. - Prepared as part of a study on Foreign ownership: c](#)
- [Strategien gegen die Arbeitslosigkeit - Analysen zur wirtschaftl. Fehlentwicklung u. wirtschaftspoli](#)