
Three cases of optic nerve tumour

[Adlard and Son] - Eye and Optic Nerve Tumours, overview and information. Patient



Description: -

Islam -- Africa, East -- History -- Sources -- Congresses.

Islam -- Africa, East -- Congresses.

Religion -- Philosophy

Science -- Philosophy

Optic nerve -- Tumors -- Case studies.

Hematemesis. Three cases of optic nerve tumour

-Three cases of optic nerve tumour

Notes: Reprinted from the Opthalmological Societys transactions,

Vol.XXXI, 1911.

This edition was published in 1911



Filesize: 38.110 MB

Tags: #Optic #Nerve #Tumor

Tumors Affecting Vision: Treatment of Optic Nerve and Chiasm Tumors

Such patients should undergo regular clinical examinations every 3—6 months and repeat CT scanning or MR imaging every 6—12 months. Residual or recurrent tumor is often treated with Gamma Knife radiosurgery. Most of these tumours are unilateral, but bilateral cases do occur, most often in adolescent men and women.

Optic Nerve Tumor

Most are choroidal, with the remainder arising from the iris and ciliary body.

Optic Nerve Tumor

Schwannoma Schwannomas are benign tumours that arise from Schwann cells in the peripheral nervous system. Indeed, according to Chutorian et al, 75% of patients with optic nerve gliomas become symptomatic in the first decade of life, and 90% become symptomatic during the first two decades of life.

Eye and Optic Nerve Tumours, overview and information. Patient

Forty-eight cases were collected from the past 21 years at the Department of Ophthalmology, West China Hospital, Sichuan University. Marked, homogenous enhancement with gadolinium is also typical.

Related Books

- Plotinus on number
- Psalms a new translation
- The relation of bacteriology to public health, with special reference to diphtheria diagnosis and to
- Behind closed doors womens oral narratives in Tunis
- Study of the relationships between brown trout and eels in a New Zealand stream