


Immunodiagnosis of schistosomiasis.

University of Birmingham - Non



Description: -

-Immunodiagnosis of schistosomiasis.

-Immunodiagnosis of schistosomiasis.

Notes: Thesis (Ph.D.) - University of Birmingham, Dept of Immunology.

This edition was published in 1983



Filesize: 60.76 MB

Tags: #Present #aspects #of #immunodiagnosis #of #schistosomiasis

Non

The DOR for IHA was better than that for ELISA. Before reagents and procedures can be recommended for routine use of national control programmes, the assays must be standardized with sera from clinically well-characterized patients in geographically defined regions, hence emphasizing the need for a reference serum bank. Eur J Clin Microbiol Infect Dis.

Immunodiagnosis in Schistosomiasis

Stool samples were examined using the Kato method, while circulating anodic antigen CAA and circulating cathodic antigen CCA were determined by highly specific monoclonal antibody-based ELISA's.

CAB Direct

The most efficacious and practical means of diagnosing human schistosomiasis is based on the detection of infection-specific antibodies. Nevertheless, difference of the accuracy between the two methods for diagnosis of schistosomiasis japonica has no statistical significance as shown by the result of Z test. This lack of essential knowledge is largely due to the fact that an unequivocal diagnosis of schistosomiasis is difficult to obtain.

Immunodiagnosis in Schistosomiasis

We found that both IHA and ELISA can be used to rule in or rule out the diagnosis.

Related Books

- [Positioning in media dialogue](#)
- [Beatles trilogy - the last concerts](#)
- [Angels song - a Christmas token.](#)
- [Why Potocki?](#)
- [Coinage of Saorstát Éireann 1928.](#)