

Bacterial wilt of potatoes - pseudomonas solanacearum.

International Potato Center - Biological control of bacterial wilt caused by *Pseudomonas solanacearum* in India with antagonistic bacteria



Description: -

-Bacterial wilt of potatoes - *pseudomonas solanacearum*

-

Psychologie comparee

Technical information bulletin -- 13Bacterial wilt of potatoes - *pseudomonas solanacearum*

Notes: Looseleaf

This edition was published in 1981



Filesize: 24.510 MB

Tags: #Ralstonia #solanacearum #(bacterial #wilt #of #potato)

Bacterial Wilt Disease of Potato Caused by *Ralstonia Solanacearum*

This separation was later confirmed using other molecular markers. Resumenes XXXVI Congreso Anual, Sociedad Agronomica de Chile, Noviembre 1985, Universidad Austral de Chile, Valdivia, Chile.

Bacterial Wilt Disease of Potato Caused by *Ralstonia Solanacearum*

Rotation of groundnut with rice and with maize, wheat, sorghum and sugarcane was effective in reducing incidence and severity ;. Cutting seed potato tubers should be avoided.

Ralstonia solanacearum (bacterial wilt of potato)

In: Plant Pathogenic Bacteria E.

Southern Bacterial Wilt

Annual Review Phytopathology 42: 107-134.

Related Books

- [Incentives and choice in health care](#)
- [Complete art of poetry.](#)
- [Opportunities - coal and cement industries.](#)
- [Attarace themba](#)
- [Newcomers to America, 1400s-1600s](#)