

Early Tertiary flood basalts from Hareoen and western Nugssuaq, west Greenland

Geological Survey of Greenland - Fagaceae pollen from the early Cenozoic of West Greenland: revisiting Engler's and Chaney's Arcto



Description: -

- Basalt - Greenland Early Tertiary flood basalts from Hareoen and western Nugssuaq, west Greenland

- Early Tertiary flood basalts from Hareoen and western Nugssuaq, west Greenland

Notes: 11

This edition was published in -



Filesize: 53.13 MB

Tags: #Fagaceae #pollen #from #the #early #Cenozoic #of #West #Greenland: #revisiting #Engler's #and #Chaney's #Arcto

Browse by Year

Berichte zur Polarforschung 336; 97 pp.

Continental margin off Western India and Deccan Large Igneous Province

REE and multi-element diagrams for representative uncontaminated, enriched and alkaline rocks of the Vaigat Formation.

Ajit Vadakayil: SHIVA LINGAM METEORITE HIT 65 million years ago at Bombay high, discharge of DNA and iridium , dinosaur extinction, birth of humans on planet earth .. Capt Ajit Vadakayil

Cretaceous extinctions: the volcanic hypothesis.

40Ar/39Ar geochronology of the West Greenland Tertiary volcanic province

It doesn't owe its existence to authority figures. K, the first letter of the German word Kreide chalk , is the traditional abbreviation for the Cretaceous Period and Pg is the abbreviation for the Paleogene Period. These data help to constrain the altitude of the lava field to being at, or near, sea level during this time.

Related Books

- [Introduction to individual psychology](#)
- [Komunikologija - psihologija komunikacije](#)
- [New Zealand as it is](#)
- [Workers welfare and law](#)
- [Clinical simulations in pediatrics](#)