


Factors affecting the detection of psychiatric morbidity by general practitioners

University of Birmingham - GP detection of health risk factors among general practice patients at risk of primary and secondary stroke



Description: -

- Factors affecting the detection of psychiatric morbidity by general practitioners

- Factors affecting the detection of psychiatric morbidity by general practitioners

Notes: Thesis (M.Med.Sc.) - University of Birmingham, Department of Public Health and Epidemiology.

This edition was published in 1995



Filesize: 63.910 MB

Tags: #Factors #contributing #to #the #recognition #of #anxiety #and #depression #in #general #practice

Determinants of help

Lifetime history of any psychotic disorder was reported by 3.

The recognition of psychiatric morbidity on a medical oncology ward

Managing depression in primary care: a meta-synthesis of qualitative and quantitative research from the UK to identify barriers and facilitators.

Social

A second qualitative analysis was performed with NVivo using data associated with GPs' main practice settings where GPs' spent most of their work hours--recorded in the questionnaire, namely 1 solo private clinics; 2 private group clinics; 3 HSSCs; and 4 hospitals including general and psychiatric hospitals. Accordingly, primary care is the cornerstone of current efforts to improve the performance and results of healthcare systems.

Detection of child mental health disorders by general practitioners

The finding is in line with the study of Marcus et al. Goldberg and Huxley 15 described the use of a rank order correlation coefficient between GP assessments and screening questionnaire scores as an indicator of accuracy, while Skuse e Williams 33 used the overall agreement rate as the measure.

Physical morbidity and mortality in people with mental illness

This supports the idea of watchful waiting in those with a first episode of depression and a short duration of symptoms. Geneva: World Health Organization; 2001.

The nonspecific symptom screening method: Detection of nonpsychotic morbidity based on nonspecific symptoms

This may, again, be explained by the high comorbidity in our sample. Schomerus G, Matschinger H, Angermeyer MC. The results indicated that the aggregate point prevalence of depression was 12.

Recognition, Management, and Course of Anxiety and Depression in General Practice

Although the main aim of this study was current prevalence of mental disorders, according to the MINI several lifetime prevalence could be also assessed.

Related Books

- [Rasā'il - shi'r](#)
- [Einsame Freundinnen? - lesbisches Leben während der zwanziger Jahre in der Provinz](#)
- [Collected poems, 1923-53.](#)
- [Will socialism save the world?](#)
- [Signal processing and communications - 5th biennial conference proceedings, 1999, July 21-24, 1999](#)