


Mahogany of tropical America - its ecology and management

University of Michigan Press - Академия Google



Description: -

- Antiques / Collectibles

Stamps

Calvinism

Mahogany. Mahogany of tropical America - its ecology and management

- Mahogany of tropical America - its ecology and management

Notes: Bibliography: p. 189-205.

This edition was published in 1966



Filesize: 25.106 MB

Tags: #Swietenia #mahagoni

Swietenia mahagoni

In Floyd RB, Hauxwell C, eds.

EENY336/IN613: Mahogany Shoot Borer, *Hypsipyla grandella* (Zeller) (Insecta: Lepidoptera: Pyralidae: Phycitinae)

Dordrecht: Kluwer Academic Publication, 2001. Their wood is used locally in numerous ways throughout the American Tropics, and they are planted as shade trees in urban areas of this region, but are present only as occasional specimen trees in Florida.

Mahogany

Current status of the taxonomy of *Hypsipyla* Ragonot Pyralidae: Phycitinae. Stable isotopes in ecological research. Effects of partial overstory removal on the natural regeneration of a tropical forest in Quintana Roo, Mexico.

Академия Google

The climate is warm sub-humid, with rains during the summer and partially in the winter, averaging 1290 mm per year with 26 °C average temperature.

Regenerating mahogany (*Swietenia macrophylla*) from seed in Quintana Roo, Mexico: the effects of sowing method and clearing treatment

Today, mahogany harvesters in Brazil are required to retain 20% of commercial trees and perform enrichment plantings in log landings. CAB Direct provides a convenient, single point of access to all of your CABI database subscriptions. Spain itself had no timber suitable for building ships and its unfriendly relations with northern Europe made drawing supplies from that source impossible; consequently it obtained timber from San Domingo, Cuba and Jamaica for building many ships of the Spanish Armada prior to 1588.

‘Marketing’ species conservation

Some of this was re-exported to continental Europe, but most was used by British furniture makers.

EENY336/IN613: Mahogany Shoot Borer, *Hypsipyla grandella* (Zeller) (Insecta: Lepidoptera: Pyralidae: Phycitinae)

Rainfall distribution is seasonal, being below 100 mm from August to November. He suggested, however, that chemical control of these pests might be applicable to nursery situations.

Related Books

- [Wanderer of St. Paul - the first decade, 1867-1877 : a mirror of the German-Catholic immigrant exper](#)
- [Middle management and nurses working in the National Health Service.](#)
- [Duas línguas, dois amigos - uma saudade : Jorge Barbosa em Inglês](#)
- [Lotus guide to 1-2-3 release 2.4](#)
- [Teoria dell'indagine e transazione](#)