


# Pharmacological and functional differentiation of dopamine D1c and D1d receptor subtypes - two novel members of the D1-like receptor family.

National Library of Canada - Protocols and Video Articles Authored by Mona B. Herold  
(Translated to Portuguese)



Description: -

-Pharmacological and functional differentiation of dopamine D1c and D1d receptor subtypes - two novel members of the D1-like receptor family.

-

Veröffentlichungen des Frobenius-Instituts an der Johann Wolfgang Goethe-Universität zu Frankfurt/Main

Veröffentlichungen d. Forschungsinstituts für Kulturmorphologie

Agard lecture series -- 115

Canadian theses = -- Thèses canadiennes Pharmacological and functional differentiation of dopamine D1c and D1d receptor subtypes - two novel members of the D1-like receptor family.

Notes: Thesis (M.Sc.) -- University of Toronto, 1996.

This edition was published in 1996



Filesize: 39.105 MB

Tags: #Analysis #of #two #D1

## Analysis of two D1

The dopaminergic system during singing behavior in zebra finches.

## Protocols and Video Articles Authored by Mona B. Herold (Translated to Portuguese)

Borba JC, Henze IP, Silveira MS et al 2005 Pituitary adenylate cyclase-activating polypeptide PACAP can act as determinant of the tyrosine hydroxylase phenotype of dopaminergic cells during retina development. Several compounds were identified that exhibited D 2 versus D 3 DAR selectivity with the most promising hit compounds being selected for chemical optimization. Rapid regulation of depression-related behaviours by control of midbrain dopamine neurons.

## Comparative pharmacological characterization of D1

Using 10 min sampling intervals they found higher levels of DA in Area X during singing than silence and during directed than undirected singing the directed singing is oriented towards another individual, usually a female, and is accompanied by courtship while undirected singing is not oriented to any particular individual.

## Evolution and Origin of the Diversity of Dopamine Receptors in Vertebrates

Solvation energies, charges on the nucleophile or leaving group, and structural features are compared for the various SN2 reactions in a spectrum of solvents.

## Comparative pharmacological characterization of D1

Negligible TMB signals coming from such negative control samples are necessary to validate the results from the experimental samples. Feedback circuitry within a song-learning pathway.

#### **allele substitution effects: Topics by Science.gov**

Wash in PBS and counterstain culture cells with DAPI. Patients' adherence to and persistence with therapy may be negatively influenced by switching, which will also impact on treatment effectiveness. Furthermore, we identified differences in the magnitude of effects between paternally and maternally inherited DGAT1 promoter VNTR — K232A haplotypes indicating parent-of-origin effects on milk production traits.

#### **Analysis of two D1**

Use-dependent regulation of GABAA receptors. Stabilization was also observed with glycerol and diethylene glycol. Dopamine receptors DARS belong to a large superfamily of neurotransmitter and hormone receptors.

## Related Books

- [Komplexbildung in Lösung - Methoden zur Bestimmung der Zusammensetzung und der Stabilitätskonstant](#)
- [Humphry Davy - life beyond the lamp](#)
- [Rectors grandchildren](#)
- [Effect of praise and competition on the persisting behavior of kindergarten children](#)
- [Cadre de référence pour l'établissement des plans d'intervention pour les élèves handicapés et l](#)