

Russian-German cooperation - the expedition TAYMYR 1995 and the expedition KOLYMA 1995 of the ISSP Pushchino Group

Alfred-Wegener-Institut für Polar- und Meeresforschung - Browse by AWI author ID

Description: -

-

Executives -- Peru.

Communist education -- Bulgaria

Education -- Bulgaria -- Curricula

Science

Environmental Science

Environmental Conservation & Protection - General

Life Sciences - Botany

Nature/Ecology

Plants - General

Life Sciences - Biology - General

Environmental Studies

Nature

Saqqarah (Egypt)

Egypt

Antiquities

Religion / Inspirational

Inspirational - General

Religion

Inspirational

Christianity

Paleoclimatology -- Holocene.

Paleoclimatology -- Russia (Federation) -- Taymyr Peninsula.

Paleoecology -- Holocene.

Paleoecology -- Russia (Federation) -- Taymyr Peninsula.

Russian-German cooperation - the expedition TAYMYR 1995 and the expedition KOLYMA 1995 of the ISSP Pushchino Group

-

Nr. 211.


Berichte zur Polarforschung ;

211

Ber. Polarforsch., Russian-German cooperation - the expedition TAYMYR 1995 and the expedition KOLYMA 1995 of the ISSP Pushchino Group

Notes: Includes bibliographical references (p. 133-137).

This edition was published in 1996



Filesize: 8.28 MB

Missouri Valley, Iowa and Nebraska. Geological Survey of Canada; Forum 1997; abstracts.

Russian

Aproximación a la caracterización estival del campo térmico.

High

The core consists of two major sediment types, clastic varves and sandy layers, which are linked to seasonal sediment supply related to meltwater

Tags: #Browse #by #AWI #author #ID

Browse by AWI author ID

Historia y Geografía, Universidad Rovira i Virgili, Tesis de Licenciatura.

High

Wrocław, Poland: Wydawnictwa Uniwersytetu Wrocławskiego.

Browse by AWI author ID

Annals of glaciology, 21, 213-218.
Solifluction deposits in the late Wisconsinan Peoria Loess of the middle

runoff and episodic events related to turbidity currents, respectively. Waterloo, ON, Canada: Geological Association of Canada. The peculiarities of utilization of the technogenic waste into permafrost.

Laminated Sediments from Levinson

Colloque Formations superficielles et geomorphologie. Climatic warming at the Pleistocene-Holocene transition led to increased biogenic accumulation, originating from both aquatic production and terrestrial supply, but to no significant change in sedimentation rates. Report of geothermal modelling, offshore temperature profiles at Angasak and Amauligak, Beaufort Shelf.

Paleoenvironmental and paleoclimatic records from permafrost deposits in the Arctic region of Northern Siberia

Bosquejo geomorfologico de la region Pampa de Chalia, Alto Rio Mayo, Chubut, Argentina. Morphological features of the active rock glaciers in the Italian Alps and climatic correlations.

Russian

Grand Forks, ND, United States: North Dakota Geological Survey. Ground freezing 88: Proceedings of the fifth international symposium on Ground freezing 88. In: Russian-German Cooperation: The Expedition Taymyr 1995 and the Expedition Kolyma 1995 of the Pushchino Group, Bolshiyarov, D.

Related Books

- [Allegorie, du moyen age au XVIIe siècle. - Chanson et littérature, du XVIIe au XIXe siècle. George](#)
- [Railroad deregulation act of 1979 - hearings before the Subcommittee on Surface Transportation of th](#)
- [New minerals, 1990-1994](#)
- [Seasonal variations in the Canadian economy - employment and unemployment = Variations saisonnières](#)
- [Reoperative aesthetic & reconstructive plastic surgery](#)