

Stratigraphy and sedimentology of the Lower Cretaceous Hulcross and Boulder Creek Formations, northeastern British Columbia

Energy, Mines and Resources - A Morphotype Catalogue, Floristic Analysis and Stratigraphic Description of the Aspen Shale Flora(Cretaceous



Description: -

-

Watercolor painting -- Exhibitions.

Latin America -- History.

America -- Discovery and exploration -- Spanish.

Panama -- History.

Geology - British Columbia
Stratigraphy and sedimentology of the Lower Cretaceous Hulcross and Boulder Creek Formations, northeastern British Columbia

-Stratigraphy and sedimentology of the Lower Cretaceous Hulcross and Boulder Creek Formations, northeastern British Columbia

Notes: 11

This edition was published in -



Filesize: 26.410 MB

Tags: #Browse #subject: #Geology, #Stratigraphic

Dr. Guy Plint

Bustin at the Earth and Ocean Sciences, University of British Columbia. In addition, sandy lithofacies are much less prevalent in the North Telkwa region than in the Goathorn region. During transgression, the uppermost sediments of Unit III were winnowed and reworked by shallow marine wave, current, and tide processes and redeposited as a sheet sandstone.

Cretaceous (Albian to Cenomanian) biostratigraphic, paleogeographic and paleoenvironmental reconstruction of the northern Western Interior Sea: Yukon, Canada

The most important single factor was probably that of local confining pressure in a structurally complex environment.

USGS Geologic Professional Papers 5

Eastern Gravina belt strata accumulated along the western margin of the Stikine, Yukon-Tanana, and Taku terranes. The Hazelton rocks are intruded by the Rocher Deboile stock, which underlies 27 square miles in the north central part of the range. The Otter Lakes Upland in the northeast is a southern extension of the Buffalo Head Hills.

USGS Geologic Professional Papers 5

The matrix is composed of pore-filling carbonate cement, with thin rims of iron oxide around many of the grains.

Stitch in the ditch: Nutzotin Mountains (Alaska) fluvial strata and a dike record ca. 117

Also within the Belle Fourche Member and above the Fish Scales Zone, the *Ostrea beloiti* limestone beds represent a distinctive marker consisting of inoceramid or ostreid fragments McNeil and Caldwell, 1981 which, with an associated bentonite, can be traced as far south as Oklahoma and westward into Montana. The map area is underlain by two magmatic arc successions, the Late Triassic Nicola Group and overlying Early Cretaceous Spences Bridge Group. Mihalynuk Open File 2009-11 is a map of Saltsping Island and an accompanying document of descriptions and comments meant to enhance the 1:25,000 scale map.

Related Books

- [Grillparzers Werke](#)
- [Supervision for success in government - a practical guide for first line managers](#)
- [Indefinitpronomina des Deutschen - Aspekte ihrer Verwendung und ihrer historischen Entwicklung](#)
- [Côtes guerrières - mer, guerre et pouvoirs au Moyen Age : France-façade océanique, XIIIe-XVe siècle](#)
- [Presses of the Pacific Islands 1817-1867 - a history of the first half century of printing in the Pa](#)