

# Habits, food, and economic status of the band-tailed pigeon

U.S. Dept. of the Interior, Fish and Wildlife Service : for sale by Supt. of Docs., U.S. Govt. Print. Off. - Band

| Trap site   | Age | Sex    | Migration     |               |
|-------------|-----|--------|---------------|---------------|
|             |     |        | movement date | Distance (km) |
| Los Alamos  | AHY | Male   | 21 Sep 2015   | 272.17        |
| Los Alamos  | AHY | Male   | 12 Nov 2015   | 284.92        |
| Silver City | AHY | Male   | 19 Nov 2015   | 268.84        |
| Silver City | AHY | Male   | 26 Oct 2015   | 272.07        |
| Weed        | AHY | Female | 26 Dec 2015   | 424.44        |
| Weed        | AHY | Male   | 20 Aug 2015   | 553.72        |
| Weed        | AHY | Female | 25 Sep 2015   | 361.08        |

Description: -

-

Band-tailed pigeon.Habits, food, and economic status of the band-tailed pigeon

-

58

North American fauna ;Habits, food, and economic status of the band-tailed pigeon

Notes: Bibliography: p. 71-74.

This edition was published in 1947



Filesize: 4.16 MB

Tags: #Patagioenas #fasciata

## ADW: Columba fasciata: INFORMATION

Use of mineral sites is most notable in Coastal populations, and less so in Interior populations where mineral needs may instead be satisfied by abrasion of grit that is highly basic Braun 1994.

### Patagioenas fasciata

Concerns for populations due to excessive harvest in the USA were again voiced by Neff 1947 , Morse 1950 and Einarsen 1953 , following a large concentration of Band-tailed Pigeons in California in 1949 McMillan 1949 and subsequent high harvest e. Nesting and reproduction of the band-tailed pigeon in California. During the first few weeks of the nestling's life, the parents' crops produce a thick, milky substance made of protein and fat.

### Band

Flight is usually swift and direct.

### Patagioenas fasciata

Proceedings, Western Association of State Game and Fish Commissioners. Its range in British Columbia expanded northward along the coast and eastward into the Southern Interior in the 1980s, but it has since largely disappeared from the Interior.

### Habits, food, and economic status of the band

Long term BBS trends in the USA are negative but nonsignificant; the annual trend over the last 3 generations in the USA is 2. John is regularly consulted on the status and conservation of birds in British Columbia by the provincial and federal governments, industry, and NGOs. When and

how are populations limited? The reference section is incredibly complete and the thorough indexing makes it rather simple to located any particular feature.

### **Habits, Food, and Economic Status of the Band**

Two broods per season not unusual, and three broods possible Keppie and Braun 2000.

## Related Books

- [Carcere speciale](#)
- [Pionery raketnoi tekhniki - Kibalchich, Tsiolkovskii, Tsander, Kondratyuk : izbrannye trudy](#)
- [Break-up of the Soviet Empire in Eastern Europe.](#)
- [Peril and promise - an inquiry into freedom of the press](#)
- [Roebings Delaware & Hudson Canal aqueducts](#)