


# Rabies

Dept. of Health, Education, and Welfare, Public Health Service, National Institutes of Health - Rabies: Symptoms, Treatment & Prevention



Description: -

-

Food service -- United States

Food industry and trade -- United States

Communicable diseases

RabiesRabies

-

DHEW publication -- no. (NIH) 79-221

DHEW publication ; no. (NIH) 79-221

Infectious diseases researchRabies

Notes: January 1979

This edition was published in 1979



Filesize: 54.93 MB

Tags: #Rabies: #Types, #Symptoms #& #Causes

## Rabies: Types, Symptoms & Causes

Rabies can be prevented by vaccinating pets, staying away from wildlife, and seeking medical care after potential exposures before symptoms start. Hundreds of thousands of animals undergo tests and observation. It from 3 to 12 weeks, but it can take as little as 5 days or more than 2 years.

## Rabies

Rabies is both prevented and treated with a rabies vaccine. Last medically reviewed on October 19, 2015.

## Rabies: Symptoms, causes, treatment, and prevention

If you have been bitten by a wild animal, doctors will typically administer a preventive shot of rabies vaccine to stop the infection before symptoms set in. If you experience any moderate to severe side effects, call your doctor right away. Rabies is a fatal but preventable viral disease.

## Rabies

Determining whether the animal that bit you has rabies In some cases, it's possible to determine whether the animal that bit you has rabies before beginning the series of rabies shots. This group includes people whose activities bring them in frequent contact with animals that could be rabid.

## Related Books

- [Maktub & malfuz literature as a source of socio-political history](#)
- [Shpionazh i biznes](#)
- [Jiaxuan ci bian nian jian zhu](#)
- [Japanese government leadership and management](#)
- [Brüner Beiträge zur Germanistik und Nordistik](#)