


# Polychètes errantes.

**Kraus - Polychæta of the Families Polynoidæ and Acætidæ from the Vicinity of the Panama Canal, collected by Dr. C. Crossland and Dr. Th. Mortensen**



Description: -  
-Polychètes errantes.  
-  
Canada Drb Drea Technical Memorandum -- 84/Y  
Faune de France -- 5Polychètes errantes.  
Notes: First pub. 1923.  
This edition was published in 1969



Filesize: 62.71 MB

Tags: #Pierre #Fauvel

## Pierre Fauvel

Estimation of the *Diopatra neapolitana* annual harvest resulting from digging activity in Canal de Mira, Ria de Aveiro. Most of the species were found in the 0—200 m stratum, so we considered this layer to calculate the standing crop of the entire polychaete community in the whole study area.

## Performance of polychaete assisted sand filters under contrasting nutrient loads in an integrated multi

Les sexes sont séparés chez les Polychètes. Une antenne impaire et deux latérales plus courtes '. Alien species in the Mediterranean Sea by 2010.

## Polychaeta collected by U. D. Gaikwad at Ratnagiri, south of Bombay

All the species found in the present study except *Pedinosoma curtum* that represents only the 0. Dentre as sinapomorfias sugeridas para Phyllococida, pode-se citar os palpos sensoriais posicionados ventralmente; uma faringe eversível axial, também chamada de probóscide; a presença de um cirro anterior alongado; e compostas com um único ligamento.

## Revealing polychaetes invasion patterns: Identification, reproduction and potential risks of the Korean ragworm, *Perinereis lineata* (Treadwell), in the Western Mediterranean

On dit alors qu'elles sont simples, doubles, opposées ou engrenantes. Underwater polychaetes have eversible mouthparts used to capture prey.

## Pierre Fauvel

Les organes nucaux sont des fossettes ciliées, au nombre d'une paire, insérées sur la partie postérieure de l'encéphale et innervées par lui. Ce tube n'est pas fixé et la Pectinaire le traîne avec elle, comme un Dentale sa coquille fig.

## Nereis

Aquatic Conservation: Marine and Freshwater Eco-system, 26, 392-09. Corps de largeur presque uniforme. Polychaetes from the coast of northern Cyprus eastern Mediterranean Sea , with two new records for the Mediterranean Sea.

**Functional morphology and feeding behavior of *Streblospio benedicti* and *S. shrubsolei* (Polychaeta: Spionidae)**

Les palpes sont couverts de fines papilles chitineuses. Partie basilaire couverte de fines papilles coniques, très nombreuses, très serrées, non disposées en rangées longitudinales. This strategy has already been successfully used in a previous study, where PASFs were stocked with juvenile forms of *Perinereis helleri* and *P.*

**[PDF] Polychaetes (Annelida: Polychaeta) described for the Mexican Pacific: an historical review and an updated checklist**

Parapodes subbirèmes, appendices glabres fig. Data were previously checked for normality Anderson—Darling test and homogeneity of variances Bartlett's and Levene's tests.

## Related Books

- [Beginnelsen van beleidsontwikkeling](#)
- [Histoire de la corrida en France - du Second Empire à nos jours](#)
- [Metode Istinbath al-Ahkam K.H.E. Abdurrahman dalam menentukan hukum zakat, infâq, dan shadaqah \(ZIS\)](#)
- [Retail strategy and structure - a management approach](#)
- [South Africa - what of the Church.](#)