

Antarctic ecosystems - environmental contamination, climate change, and human impact

Springer - Climate Change and the Antarctic



Description: -

- Antarctica -- Environmental conditions
- Climatic changes -- Environmental aspects -- Antarctica
- Pollution -- Environmental aspects -- Antarctica
- Biotic communities -- Antarctica
- Nature -- Effect of human beings on -- Antarctica
- Ecology -- Antarctica
- Antarctic ecosystems - environmental contamination, climate change, and human impact

- v. 175

Ecological studies, Antarctic ecosystems - environmental contamination, climate change, and human impact

Notes: Includes bibliographical references (p. [315]-377) and indexes.

This edition was published in 2005



Filesize: 7.410 MB

Tags: #Climate #Change, #Human #Impacts, #and #Coastal #Ecosystems #in #the #Anthropocene

7. Impacts of climate change on biodiversity and ecosystem services

Adélie penguin populations have been declining in recent years due to reductions in krill populations and changing weather conditions in their traditional nesting areas. A summary of sample sizes and biological parameters monitored annually is presented in Table.

Climate Change Impact on Arctic Ecosystems

There are indications that whale populations are beginning to recover.

Arctic climate change and pollution impact little auk foraging and fitness across a decade

They cite evidence that declining seawater pH will particularly affect organisms with calcified shells and skeletal elements, such as molluscs, seastars, sea urchins, coralline algae and cold-water corals. They also highlight evidence suggesting that ocean acidification could profoundly alter the structure and functioning of the planktonic food web, with unknown consequences for animals further up the food chain, including commercially exploited fish. Another key planning area involves prioritizing research as a means of filling in knowledge gaps and facilitating the adoption of effective adaptation responses.

Climate Change and the Antarctic

While little auks foraged in the same areas at the shelf break with or without sea-ice , we found that diving behaviour changed with SIC: Birds performed shallower and shorter dives when SIC was the highest Fig. Aquatic freshwater habitats and wetlands, mangroves, coral reefs, arctic and alpine ecosystems, and cloud forests are particularly vulnerable to the impacts of climate change.

Related Books

- [Ninety years of cinema in Worcester](#)
- [Assessment of commercial industrial projects in developing countries - the Bahraini experience](#)
- [Guangdong jing ji zhi gai ge yan jiu](#)
- [Tree Museum - 2004, the 7th annual exhibition, Exchange - changing the landscape, September 1 to Oct](#)
- [Records of the Department of State relating to political relations between the United States and Mex](#)