

# Multiple medical impairment study

**CMAS Books of MIB, Inc. - Verbal fluency deficits in multiple sclerosis may reflect impaired language ability**

Description: -

-  
Insurance, Life  
Actuarial Analysis  
Morbidity -- Canada  
Morbidity -- United States  
Mortality -- Canada  
Mortality -- United States  
Canada -- Statistics, Medical -- Tables  
Insurance, Health -- Canada -- Tables  
Mortality -- Canada -- Tables  
United States -- Statistics, Medical -- Tables  
Insurance, Health -- United States -- Tables  
Mortality -- United States -- Tables  
Multiple medical impairment study  
-Multiple medical impairment study  
Notes: Includes index.  
This edition was published in 1998



Filesize: 35.35 MB

Tags: #Associations #between #cognitive #impairment #at #onset #and #disability #accrual #in #young #people #with #multiple #sclerosis

## Associations between cognitive impairment at onset and disability accrual in young people with multiple sclerosis

The involvement of audio-motor coupling in the music-supported therapy applied to stroke patients. Due to the heterogeneity of the study populations, study designs and protocols, and variability in the objective testing measures between studies, we were unable to complete a meta-analysis. Limitations Findings from this review were constrained by the limitations of the current literature.

## Rehabilitation of Motor Function after Stroke: A Multiple Systematic Review Focused on Techniques to Stimulate Upper Extremity Recovery

Conclusions: Overall, this retrospective study suggests that multiple myeloma patients are more likely to experience renal impairment with zoledronic acid than with ibandronate. However, further RCTs are needed to ascertain treatment effects of virtual immersion and serious gaming.

## Multiple Sclerosis Clinical Trials

Associations of substance use, psychosis, and mortality among people living in precarious housing or homelessness: a longitudinal, community-based study in Vancouver, Canada.

## Impairment in oral medical comprehension in multiple sclerosis

These exercises can be performed against a manual resistance exerted by the therapist or using weight-bearing apparatus. Gaming literature emphasizes its potential to increase: patient motivation, learning through repetition in an enriched environment, confidence through reinforcement and immediate feedback, and positivity through achievement and social interaction Krichevets et al. Transcranial direct current stimulation of the affected hemisphere does not accelerate recovery of acute stroke patients.

---

## Related Books

- [Sagittarius in Warsaw](#)
- [Tu p'asan](#)
- [Glass ceiling](#)
- [Silvano Barba González - apuntes de su biografía](#)
- [52 offbeat Texas stops - traveling with Bob Phillips, Texas country reporter](#)