

Butterfly farming and trading agency in Irian Jaya

WWF/IUCN Conservation for Development Programme in Indonesia - Insect Zoos



Description: -

-butterfly farming and trading agency in Irian Jaya

-butterfly farming and trading agency in Irian Jaya

Notes: Microfiche. Jakarta : Library of Congress Office ;

Washington, D.C. : Library of Congress Photoduplication Service,

1995. 2 microfiches ; 11 x 15 cm.

This edition was published in 1986



Filesize: 35.16 MB

Tags: #261407387

Status of biodiversity in Papua New Guinea

However, more specimens are needed to draw a final conclusion.

Status of biodiversity in Papua New Guinea

The finger-cutting tradition is also seen as a way to prevent the recurrence of a catastrophe that claimed the life of a grieving family member.

Insect Zoos

Although general statistics on non-indigenous animals in Papua New Guinea are not readily available, there are data on the various species of molluscs, insects, fish, amphibians, birds, and mammals that have been introduced. The smuggler ultimately provided the solution. The freshwater ichthyofauna can be clearly divided into two zoogeographic regions.

The butterfly sting

Many of these organisms are endemic; that is, they are found only in Papua New Guinea or on the island of New Guinea. On a per capita basis, the GDP growth rates for both Papua and Indonesia are lower than those for total GDP.

Nomenclatural updates of *Aristolochia* subgenus *Pararistolochia* (Aristolochiaceae)

Las larvas fueron criadas en el campo y en el laboratorio, se alimentaron de *Ceiba pentandra* y *Chorisia* sp. However, the negative effects of deforestation on the local environment should be a major consideration in the selection of new areas for this and any other plantation crop. Sago is either processed as a pancake or sago congee called , usually eaten with yellow soup made from , or other fishes spiced with , , and other spices.

Butterfly Conservation Management, Annual Review of Entomology

Abstract: All larval instars of *Papilio homerus* have well developed osmeteria. Resume: *Meroleuca Meroleucoides amarillae* n.

Status of biodiversity in Papua New Guinea

Loranthomitoura, a new genus of Eumaeini Lepidoptera: Lycaenidae: Theclinae. Papua, along with West Papua, has a higher degree of autonomy level compared to other. Invasive and Introduced Species Non-indigenous species represent a major economic threat.

Status of biodiversity in Papua New Guinea

The said infrastructure funding for Papua is not insignificant. Information on ecology, hostplants, taxonomy, and zoogeography of the genus is presented.

Related Books

- [Marriage is for grownups](#)
- [Funny friends.](#)
- [Mashāhid al-Jannah fi al-Qur'ān wa-al-Sunnah](#)
- [The Rebuilding of Greater Russia - Putins foreign policy towards the CIS countries \(Routledge Contem](#)
- [Methods in reproductive aquaculture - marine and freshwater species](#)