


Ecodynamics - contributions to theoretical ecology : proceedings of an international workshop, held at the Nuclear Research Centre, Jülich, Fed. Rep. of Germany, 19-20 October 1987

Springer-Verlag - Environment



Description: -

- Ecology -- Congresses. Ecodynamics - contributions to theoretical ecology : proceedings of an international workshop, held at the Nuclear Research Centre, Jülich, Fed. Rep. of Germany, 19-20 October 1987

- Research reports in physics Ecodynamics - contributions to theoretical ecology : proceedings of an international workshop, held at the Nuclear Research Centre, Jülich, Fed. Rep. of Germany, 19-20 October 1987

Notes: Includes bibliographies and index.

This edition was published in 1988



Filesize: 62.108 MB

Tags: #BIOGEOCHEMICAL

Ecodynamics

Table 9: Measurements of variables studied for different seasons. Isolates related to the genus *Arthrobacter* were the most prevalent whereas the genera *Hymenobacter*, *Deinococcus*, *Cryobacterium* and *Sphingomonas* were also recovered in high numbers in both samples. AN: 4098574 170 of 313 TI: Patterns of retention and utilization of aerially deposited nitrogen in boreal peatlands AU: Li,-Yenhung; Vitt,-D.

Environment

Reasonable tracer additions result in only nanomolar increases in the ambient silicic acid and orthophosphate concentrations during production rate experiments.

Environment

Aerosols are essential for cloud formation and therefore also for precipitation, which is the only source for mass gain of the Antarctic ice sheet. Denitrification rates obtained from isotope tracer experiments were compatible with nitrogen consumption rates observed in flux chamber experiments and whole-lake mass balances.

Ecodynamics as a Function of Ecosystem Structure: Forest Dieback as an Example

Areal integrals were calculated for N and P at each station, which ranged in depth between 15 and 150 m.

Full text of Review Vol.35,no.4,

ASPECTS OF FEEDING AND GROWTH IN JUVENILE NORTHERN PIKE, *ESOX LUCIUS*. Because the degradation rate of this material appears to be very high, its presence at the seafloor for several months per year could yield significant phytodetrital contributions to the annual seafloor organic-carbon budget.

Environment

The stream has characteristics typical of many streams throughout the southeastern Coastal Plain, including low gradient, loose sand sediment, and a well-developed flood plain in its lower reach.

BIOGEOCHEMICAL

CSSA Special Publication Number 28.

Ecodynamics : contributions to theoretical ecology : proceedings of an international workshop, held at the Nuclear Research Centre, Juelich, Fed. Rep. of Germany, 19

This may lead to subsequent inappropriate management and inappropriate NHS star ratings.

Related Books

- [În Aponiia na poroge XXI veka - referativnyi sbornik](#)
- [Irish administration, 1801-1914.](#)
- [Quaestio de unico esse in Christo - a doctoribus saeculi XIII disputata.](#)
- [Solubility of nonelectrolytes.](#)
- [Brain and nervous system](#)