


Synopsis of the North American microtine rodents

University of Kansas - Voles (Meadow Mice)



Description: -

-
Judo -- Directories.
Karate -- Directories.
Martial artists -- Biography.
Catalysis.
Metal catalysts.
Mammals -- North America.
Rodents.synopsis of the North American microtine rodents
-
v. 5, no. 27
University of Kansas publications. Museum of Natural History.
[Publications]synopsis of the North American microtine rodents
Notes: Bibliography: p. 485-498.
This edition was published in 1953



Filesize: 57.910 MB

Tags: #Microtus

CiteSeerX — Gradients in density variations of small rodents

Its business office is located at 809 North 1500 West, Salt Lake City, UT 84116, 801 596-1887, email business pglaf.

Morphometric Patterns Among Microtine Rodents. I. Sexual Selection Suggested by Relative Scent Gland Development in Representative Voles (Microtus)

Humboldt Marten *Martes caurina humboldtensis* General Information Official Status: Species of Concern, the Humboldt marten does not have any special status under the Endangered Species Act at this time, but is a candidate for listing as endangered by the State of California.

Microtine rodents lemmings and voles

The time required for the entire process varied from one day to more than two weeks depending on the size of the specimen and on its condition. These morphological characteristics of the stomach are similar across species of *Microtus*. Miscellaneous The mites *Androlaelaps fahrenheitsi*, *Echinonyssus isabellae*, *Euschoengastia zapoteca*, *Eutrombicula alfreddugesi*, *Glycyphagus hypudaei*, *Haemogamasus ambulans*, *Laelaps kochi*, *Laelaps nuttalli*, *Listrophorus pitymys*, *Myocoptes glareoli*, *Neotrombicula claudoi*, *Prolistrophorus sclerobursatus*, and *Pseudoschoengastia pedregalensis* have been found on *Microtus quasiater* ;

Morphometric Patterns Among Microtine Rodents. I. Sexual Selection Suggested by Relative Scent Gland Development in Representative Voles (Microtus)

Another number will appear in volume 11. Within Fennoscandia such indices for *Clethrionomys glareolus* and *Microtus agrestis* were clearly positively related to latitude and snow cover. Here's a look at 10 extinct animals from the last North American ice age, and what scientists know about their lives.

Gradients in density variations of small rodents: the importance of latitude and snow cover

Bristol Foster, Royal Ontario Museum of Zoology, Toronto, Canada, provided specimens of *Phenacomys intermedius*. Its tail is about as long as the hind foot. The baculum of a specimen from Switzerland is weakly developed, of small size shaft 2.

Morphometric Patterns Among Microtine Rodents. I. Sexual Selection Suggested by Relative Scent Gland Development in Representative Voles (*Microtus*)

Mention is made if maturity is known or suspected to differ in specimens being compared. There are a lot of things you can do with Project Gutenberg-tm electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg-tm electronic works. Cyclic density fluctuations may occur every 2-5 years Krebs and Myers 1974.

The Project Gutenberg eBook of The Baculum in Microtine Rodents by Sydney Anderson.

General Habitat Characteristics: American martens are typically associated with closed-canopy, late-successional, mesic coniferous forests with complex physical structure near the ground.

Microtus

Many other protective measures can be taken. The situation in the voles, as we shall see, casts a different light on the matter. S Muskwa ; *Clethrionomys gapperi galei*, 31 mi.

Related Books

- [Normas constitucionais programáticas - normatividade, operatividade e efetividade](#)
- [Kühlkugel und Wärmepfêl](#)
- [Psychoanalysis in colonial India](#)
- [Practical Amstrad machine code programming](#)
- [Tekst, diskurs, kontsept, perevod](#)