

Frogs of Southeastern Brazil.

s.n - Diversity and Abundance of Litter Frogs in a Montane Forest of Southeastern Brazil: Seasonal and Altitudinal Changes on JSTOR



Description: -
-Frogs of Southeastern Brazil.
-
Bulletin (United States National Museum) -- 206Frogs of Southeastern Brazil.
Notes: 1
This edition was published in 1955



Filesize: 12.104 MB

Tags: #Effects #of #Fragmentation #and #Sea

Frogs of Southeastern Brazil

Our international newsletter with news for tropical biologists will begin publication in Q4 2017.

Diversity and abundance of litter frogs in a montane forest of southeastern Brazil: seasonal and altitudinal changes

Our goal is to assess the effects of habitat fragmentation on anuran communities using land-bridge islands as long-term natural experiments.

[PDF] Parameters from the community of leaf

Most specimens were searched for when calling, detected visually, and hand captured. Because of the strong correlation between area, altitude and NBH, the same matrix order is obtained when sorting the islands according to each of these variables. Several studies with different taxonomic groups have demonstrated that insular communities are not randomly distributed; rather, they tend to be structured in a nested pattern —.

Native Amphibians Of Brazil

The only specimen of *Xenohyla truncata* from Marambaia is deposited in the herpetological collection housed in the Museu Nacional MNRJ *Aparasphenodon bruno*i RU 0062, RU 0104, RU 0143, RU 0144, RU 3531, RU 4862, RU 4867; *Aplastodiscus albofrenatus* RU 0078, RU 0159, RU 0253, RU 3502, RU 3758, RU 4297; *Chiasmocleis* cf. These frogs were more frequently found during the rainy season, at the peak of their reproduction in swamps, creeks, and temporary ponds, but individuals were also found during other periods of the year in or near swampy areas or in the forest litter.

SciELO

On six islands there are members of the *Scinax perpusillus* group thought to represent species yet to be described ,.

16 Poisonous Frogs That Are Beautiful but Deadly

Just a few kilometres off the coast there are hundreds of islands and islets. But don't let that small stature fool you.

Diversity and Abundance of Litter Frogs in a Montane Forest of Southeastern Brazil: Seasonal and Altitudinal Changes on JSTOR

About five thousand years ago the relative sea level was 3.

SciELO

Like most other poison frog species, they lose their toxicity in captivity as a result of an altered diet. The region encloses the Sepetiba Bay, and is comprised of an island, known as Ilha da Marambaia, linked to the continent through a sand dune formation approximately 40 km long called Restinga da Marambaia. Box 1897, Lawrence, KS 66044-8897.

Related Books

- [Eighteenth-century literature and culture](#)
- [Massage - a career at your fingertips](#)
- [GENIE LADY](#)
- [Archaeology of Mission Santa Catalina de Gaule.](#)
- [Kyrylivs'kyi monastery](#)