
Conestoga College
MATH73235: Advanced Topics in Mathematics
Spring 2025 Midterm 2
July 15, 2025

Exam Duration: 5pm - 7pm.

Student Name (print): _____

Student ID: _____

This exam contains 10 pages (including this cover page) and 5 questions. The total number of possible points is 50. Enter your answers in the space provided.

- **Organize your work**, in a reasonably neat and coherent way, in the space provided.
- **Mysterious or unsupported answers will not receive full credit.** Correct answers should be supported by calculations, explanations, or algebraic work to receive full credit; incorrect answers supported by substantially correct calculations and explanations may still receive partial credit.
- **Provide exact answers** unless otherwise instructed.
- **Simplify all answers as much as possible.** This means that you need to combine like terms, reduce fractions, etc.
- **Be sure to state units for applied problems.**
- **Clearly identify your answer for each problem.**

Question	Points	Score
1	10	
2	10	
3	10	
4	10	
5	10	
Total:	50	

Do not write in the table to the right.

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

1. (10 points) **Conceptual Questions.** Write an equation, an expression, or a short sentence to describe each of these concepts.

- (a) The expression for the volume integral $\iiint_T f(x, y, z) \, dx dy dz$ when the coordinates are changed from (x, y, z) to some other coordinates (u, v, w) .

Solution.

□

- (b) The main equation for **Stokes' Theorem** for a vector field $\vec{F}$ in a surface S with a boundary curve C traversed such that S remains on the left of C .

Solution.

□

- (c) An expression for a vector $\vec{N}$ that is normal to the surface $z - x^2 - y^2 = 0$.

Solution.

□

- (d) The vector field $\vec{F}$ polar coordinates written as $F_r \hat{e}_r + F_\theta \hat{e}_\theta$ if in Cartesian coordinates we have $\vec{F} = F_1 \hat{i} + F_2 \hat{j} = (x^2 + y^2) \hat{i} - 2y \hat{j}$. *Hint:* $F_r = F_1 \cos \theta + F_2 \sin \theta$, $F_\theta = -F_1 \sin \theta + F_2 \cos \theta$.

Solution.

□

- (e) The expression for the divergence $\nabla \cdot \vec{F}$ in spherical coordinates (ρ, θ, ϕ) if we are given that the divergence of the same vector field is $\nabla \cdot \vec{F} = r^2 + z^2$ in cylindrical coordinates (r, θ, ϕ) .

Solution.

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

2. (10 points) Use the appropriate Lamé coefficients to compute the Laplacian of the scalar function

$$f(x, y, z) = x^2 + y^2 + z^2$$

in *both* Cylindrical and Spherical coordinates. You must compute all the required partial derivatives.

Solution.

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

3. (10 points) Use **Spherical coordinates** to find the surface area of the cone $\phi = \frac{\pi}{6}$ with $0 \leq \rho \leq 2$.

Solution.

□

First name:
Last name:
Student ID:

Name and Student ID must be filled in for this page to be graded.

4. (10 points) Verify Gauss' Divergence Theorem by computing the following surface integral:

$$\oiint_S \vec{F} \cdot \hat{n} \, dA,$$

and the volume integral:

$$\iiint_T (\nabla \cdot \vec{F}) \, dV,$$

where $\vec{F} = x\hat{i} + y\hat{j} + z\hat{k}$, S is the surface of a sphere of radius 2 that forms the boundary of T .

Hint: This problem may be easier to solve in the spherical coordinates. Note that in spherical coordinates, $\vec{F} = \rho\hat{e}_\rho$, while for S , $\hat{n} = \hat{e}_\rho$, and thus

$$\vec{F} \cdot \hat{n} = (\rho\hat{e}_\rho) \cdot (\hat{e}_\rho) = \rho.$$

Solution.

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

Continue problem 4

Solution.

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

5. (10 points) In the first quadrant of the Cartesian plane (i.e., $x \geq 0, y \geq 0$), we may define a set of coordinates known as the hyperbolic coordinates (u, v) , with transformation function

$$x = ve^u, \quad y = ve^{-u}.$$

Find the area of the triangular region R given by

$$0 \leq v \leq u, \quad 0 \leq u \leq 3.$$

Solution.

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

Extra Page: Must submit even if unused.

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

Extra Page: Must submit even if unused.

Formula Sheet

General Coordinates:

$$(x_1, x_2, x_3) \rightarrow (q_1, q_2, q_3):$$

$$x_1 = x_1(q_1, q_2, q_3), \quad x_2 = x_2(q_1, q_2, q_3), \quad x_3 = x_3(q_1, q_2, q_3)$$

$$\text{Derivative Matrix: } D = \begin{bmatrix} \frac{\partial x_1}{\partial q_1} & \frac{\partial x_1}{\partial q_2} & \frac{\partial x_1}{\partial q_3} \\ \frac{\partial x_2}{\partial q_1} & \frac{\partial x_2}{\partial q_2} & \frac{\partial x_2}{\partial q_3} \\ \frac{\partial x_3}{\partial q_1} & \frac{\partial x_3}{\partial q_2} & \frac{\partial x_3}{\partial q_3} \end{bmatrix}$$

$$\text{Lamé coefficients: } h_j^2 = \sum_{k=1}^3 \left(\frac{\partial x_k}{\partial q_j} \right)^2$$

$$\text{Jacobian: } \frac{\partial(x_1, x_2, x_3)}{\partial(q_1, q_2, q_3)} = |J| = |\det D| = |h_1 h_2 h_3|$$

$$\text{Line element: } ds^2 = h_1^2 dq_1^2 + h_2^2 dq_2^2 + h_3^2 dq_3^2$$

$$\text{Volume element: } dV = h_1 h_2 h_3 dq_1 dq_2 dq_3$$

$$\text{Area element for surface with } q_j = \text{const} \ (i, j, k \text{ distinct}): } dA = h_i h_k dq_i dq_k$$

$$\text{Gradient for scalar function } f: \nabla f = \frac{1}{h_1} \frac{\partial f}{\partial q_1} \hat{e}_{q_1} + \frac{1}{h_2} \frac{\partial f}{\partial q_2} \hat{e}_{q_2} + \frac{1}{h_3} \frac{\partial f}{\partial q_3} \hat{e}_{q_3}$$

$$\text{Divergence and curl of vector field } \vec{F} = F_{q_1} \hat{e}_{q_1} + F_{q_2} \hat{e}_{q_2} + F_{q_3} \hat{e}_{q_3}:$$

$$\nabla \cdot \vec{F} = \frac{1}{h_1 h_2 h_3} \left[\frac{\partial}{\partial q_1} (h_2 h_3 F_{q_1}) + \frac{\partial}{\partial q_2} (h_3 h_1 F_{q_2}) + \frac{\partial}{\partial q_3} (h_1 h_2 F_{q_3}) \right]$$

$$\nabla \times \vec{F} = \frac{1}{h_1 h_2 h_3} \begin{vmatrix} h_1 \hat{e}_{q_1} & h_2 \hat{e}_{q_2} & h_3 \hat{e}_{q_3} \\ \frac{\partial}{\partial q_1} & \frac{\partial}{\partial q_2} & \frac{\partial}{\partial q_3} \\ h_1 F_{q_1} & h_2 F_{q_2} & h_3 F_{q_3} \end{vmatrix}$$

Laplacian of scalar field f :

$$\nabla^2 f = \frac{1}{h_1 h_2 h_3} \left[\frac{\partial}{\partial q_1} \left(\frac{h_2 h_3}{h_1} \frac{\partial f}{\partial q_1} \right) + \frac{\partial}{\partial q_2} \left(\frac{h_3 h_1}{h_2} \frac{\partial f}{\partial q_2} \right) + \frac{\partial}{\partial q_3} \left(\frac{h_1 h_2}{h_3} \frac{\partial f}{\partial q_3} \right) \right].$$

Cylindrical Coordinates:

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z.$$

$$h_1 = h_r = 1, \quad h_2 = h_\theta = r, \quad h_3 = h_z = 1.$$

Spherical Coordinates:

$$x = \rho \cos \theta \sin \phi, \quad y = \rho \sin \theta \sin \phi, \quad z = \rho \cos \phi$$

$$h_1 = h_\rho = 1, \quad h_2 = h_\theta = \rho \sin \phi, \quad h_3 = h_\phi = \rho.$$