
Conestoga College
MATH73235: Advanced Topics in Mathematics
Spring 2025 Midterm 2
July 15, 2025

Exam Duration: 5pm - 7pm.

Student Name (print): _____

Student ID: _____

This exam contains 10 pages (including this cover page) and 5 questions. The total number of possible points is 50. Enter your answers in the space provided.

- **Organize your work**, in a reasonably neat and coherent way, in the space provided.
- **Mysterious or unsupported answers will not receive full credit.** Correct answers should be supported by calculations, explanations, or algebraic work to receive full credit; incorrect answers supported by substantially correct calculations and explanations may still receive partial credit.
- **Provide exact answers** unless otherwise instructed.
- **Simplify all answers as much as possible.** This means that you need to combine like terms, reduce fractions, etc.
- **Be sure to state units for applied problems.**
- **Clearly identify your answer for each problem.**

Question	Points	Score
1	10	
2	10	
3	10	
4	10	
5	10	
Total:	50	50

Do not write in the table to the right.

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

1. (10 points) **Conceptual Questions.** Write an equation, an expression, or a short sentence to describe each of these concepts.

- (a) The expression for the volume integral $\iiint_T f(x, y, z) dx dy dz$ when the coordinates are changed from (x, y, z) to some other coordinates (u, v, w) .

Solution.

$$\iiint_T f(x(u, v, w), y(u, v, w), z(u, v, w)) \frac{\partial(x, y, z)}{\partial(u, v, w)} du dv dw$$

□

- (b) The main equation for **Stokes' Theorem** for a vector field $\vec{F}$ in a surface S with a boundary curve C traversed such that S remains on the left of C .

Solution.

$$\iint_S (\nabla \times \vec{F}) \cdot \hat{n} dA = \oint_C \vec{F} \cdot d\vec{r}$$

□

- (c) An expression for a vector $\vec{N}$ that is normal to the surface $z - x^2 - y^2 = 0$.

Solution.

$$\vec{N} = \vec{\nabla}(z - x^2 - y^2) = \begin{bmatrix} -2x \\ -2y \\ 1 \end{bmatrix}$$

□

- (d) The vector field $\vec{F}$ polar coordinates written as $F_r \hat{e}_r + F_\theta \hat{e}_\theta$ if in Cartesian coordinates we have $\vec{F} = F_1 \hat{i} + F_2 \hat{j} = (x^2 + y^2) \hat{i} - 2y \hat{j}$. Hint: $F_r = F_1 \cos \theta + F_2 \sin \theta$, $F_\theta = -F_1 \sin \theta + F_2 \cos \theta$.

Solution.

$$\begin{aligned} \vec{F} &= (F_1 \cos \theta + F_2 \sin \theta) \hat{e}_r + (-F_1 \sin \theta + F_2 \cos \theta) \hat{e}_\theta \\ &= (r^2 \cos \theta - 2r \sin^2 \theta) \hat{e}_r + (-r^2 \sin \theta - 2r \sin \theta \cos \theta) \hat{e}_\theta \end{aligned}$$

□

- (e) The expression for the divergence $\nabla \cdot \vec{F}$ in spherical coordinates (ρ, θ, ϕ) if we are given that the divergence of the same vector field is $\nabla \cdot \vec{F} = r^2 + z^2$ in cylindrical coordinates (r, θ, ϕ) .

Solution.

$$\vec{\nabla} \cdot \vec{F} = r^2$$

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

2. (10 points) Use the appropriate Lamé coefficients to compute the Laplacian of the scalar function

$$f(x, y, z) = x^2 + y^2 + z^2$$

in both Cylindrical and Spherical coordinates. You must compute all the required partial derivatives.

Solution.

In Cylindrical: $f(r, \theta, z) = r^2 + z^2$

$$\begin{aligned}\nabla^2 f &= \frac{1}{r} \left[\frac{\partial}{\partial r} \left(r \frac{\partial f}{\partial r} \right) + \frac{\partial}{\partial \theta} \left(\frac{1}{r} \frac{\partial f}{\partial \theta} \right) + \frac{\partial}{\partial z} \left(r \frac{\partial f}{\partial z} \right) \right] \\ &= \frac{1}{r} \left[\frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} (r^2 + z^2) \right) + \frac{\partial}{\partial z} \left(r \frac{\partial}{\partial z} (r^2 + z^2) \right) \right] \\ &= \frac{1}{r} \left[\frac{\partial}{\partial r} (r(2r)) + \frac{\partial}{\partial z} (r(2z)) \right] = \frac{1}{r} \left[\frac{\partial}{\partial r} (2r^2) + \frac{\partial}{\partial z} (2rz) \right] \\ &= \frac{1}{r} [4r + 2r] = \frac{1}{r} (6r) = 6\end{aligned}$$

In Spherical: $f(\rho, \theta, \phi) = \rho^2$

$$\begin{aligned}\nabla^2 f &= \frac{1}{\rho^2 \sin \phi} \left[\frac{\partial}{\partial \rho} \left(\rho^2 \sin \phi \frac{\partial f}{\partial \rho} \right) + \frac{\partial}{\partial \theta} \left(\frac{\rho}{\sin \phi} \frac{\partial f}{\partial \theta} \right) + \frac{\partial}{\partial \phi} \left(\frac{\rho \sin \theta}{\sin \phi} \frac{\partial f}{\partial \phi} \right) \right] \\ &= \frac{1}{\rho^2 \sin \phi} \left[\frac{\partial}{\partial \rho} \left(\rho^2 \sin \phi \frac{\partial}{\partial \rho} \rho^2 \right) \right] = \frac{1}{\rho^2 \sin \phi} \left[\frac{\partial}{\partial \rho} \left(\rho^2 \sin \phi (2\rho) \right) \right] \\ &= \frac{1}{\rho^2 \sin \phi} \left[\frac{\partial}{\partial \rho} (2\rho^3 \sin \phi) \right] = \frac{1}{\rho^2 \sin \phi} (6\rho^2 \sin \phi) = 6\end{aligned}$$

□

First name:
Last name:
Student ID:

Name and Student ID must be filled in for this page to be graded.

3. (10 points) Use **Spherical coordinates** to find the surface area of the cone $\phi = \frac{\pi}{6}$ with $0 \leq \rho \leq 2$.

Solution.

Since $\phi = \pi/6$ is constant, $dA = \rho \sin \phi \, d\rho \, d\theta = \rho \sin(\pi/6) \, d\rho \, d\theta = \frac{1}{2} \rho \, d\rho \, d\theta$

Then
Surface Area = $\iint_S dA = \int_0^{2\pi} \int_0^2 \frac{1}{2} \rho \, d\rho \, d\theta$
 $= \int_0^{2\pi} d\theta \int_0^2 \frac{1}{2} \rho \, d\rho$
 $= (2\pi) \left[\frac{\rho^2}{4} \Big|_0^2 \right] = 2\pi \left(\frac{4}{4} \right) = 2\pi$

□

First name:
Last name:
Student ID:

Name and Student ID must be filled in for this page to be graded.

4. (10 points) Verify Gauss' Divergence Theorem by computing the following surface integral:

$$\iint_S \vec{F} \cdot \hat{n} \, dA,$$

and the volume integral:

$$\iiint_T (\nabla \cdot \vec{F}) \, dV,$$

where $\vec{F} = x\hat{i} + y\hat{j} + z\hat{k}$, S is the surface of a sphere of radius 2 that forms the boundary of T .

Hint: This problem may be easier to solve in the spherical coordinates. Note that in spherical coordinates, $\vec{F} = \rho\hat{e}_\rho$, while for S , $\hat{n} = \hat{e}_\rho$, and thus

$$\vec{F} \cdot \hat{n} = (\rho\hat{e}_\rho) \cdot (\hat{e}_\rho) = \rho.$$

Solution.

Surface integral: as $\rho = 2$ is constant, $dA = \rho^2 \sin\phi \, d\theta \, d\phi = 4 \sin\phi \, d\theta \, d\phi$

$$\begin{aligned} \text{Then: } \iint_S \vec{F} \cdot \hat{n} \, dA &= \iint_S \rho \, dA \quad (\text{by given hint}) \\ &= \iint_S 2 \, dA = \int_0^{2\pi} \int_0^\pi (2)(4 \sin\phi \, d\phi \, d\theta) \\ &= 8 \int_0^{2\pi} d\theta \int_0^\pi \sin\phi \, d\phi = 8(2\pi) [-\cos\phi]_0^\pi = 8(2\pi)(2) \\ &= 32\pi \end{aligned}$$

Volume integral:

$$\text{Note: } \nabla \cdot \vec{F} = \frac{\partial}{\partial x} x + \frac{\partial}{\partial y} y + \frac{\partial}{\partial z} z = 3$$

So in spherical coordinates

$$\begin{aligned} \iiint_T \nabla \cdot \vec{F} \, dV &= \iiint_T 3 \, dV \\ &= \int_0^{2\pi} \int_0^\pi \int_0^2 3 \rho^2 \sin\phi \, d\phi \, d\theta \, d\rho \\ &= 3 \int_0^{2\pi} d\theta \int_0^\pi \sin\phi \, d\phi \int_0^2 \rho^2 \, d\rho \\ &= 3(2\pi) [-\cos\phi]_0^\pi \left[\frac{\rho^3}{3} \right]_0^2 = 3(2\pi)(2) \left(\frac{8}{3} \right) \\ &= 32\pi \end{aligned}$$

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

Continue problem 4

Solution.

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

5. (10 points) In the first quadrant of the Cartesian plane (i.e., $x \geq 0, y \geq 0$), we may define a set of coordinates known as the hyperbolic coordinates (u, v) , with transformation function

$$x = ve^u, \quad y = ve^{-u}.$$

Find the area of the triangular region R given by

$$0 \leq v \leq u, \quad 0 \leq u \leq 3.$$

Solution.

$$\text{Need } \frac{\partial x}{\partial v} = e^u \quad \frac{\partial x}{\partial u} = ve^u$$

$$\frac{\partial y}{\partial v} = e^{-u} \quad \frac{\partial y}{\partial u} = -ve^{-u}$$

$$\text{Then } \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} ve^u & e^u \\ -ve^{-u} & e^{-u} \end{vmatrix} = (ve^u \cdot e^{-u} - (-ve^{-u} \cdot e^u)) = 2v$$

Then the area of the region is

$$\begin{aligned} \iint_R dA &= \int_0^3 \int_0^u 2v \, dv \, du \\ &= \int_0^3 \left[v^2 \Big|_0^u \right] du = \int_0^3 u^2 \, du = \frac{u^3}{3} \Big|_0^3 \\ &= \frac{27}{3} = 9. \end{aligned}$$

□

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

Extra Page: Must submit even if unused.

First name:

Last name:

Student ID:

Name and Student ID must be filled in for this page to be graded.

Extra Page: Must submit even if unused.

Formula Sheet

General Coordinates:

$$(x_1, x_2, x_3) \rightarrow (q_1, q_2, q_3):$$

$$x_1 = x_1(q_1, q_2, q_3), \quad x_2 = x_2(q_1, q_2, q_3), \quad x_3 = x_3(q_1, q_2, q_3)$$

$$\text{Derivative Matrix: } D = \begin{bmatrix} \frac{\partial x_1}{\partial q_1} & \frac{\partial x_1}{\partial q_2} & \frac{\partial x_1}{\partial q_3} \\ \frac{\partial x_2}{\partial q_1} & \frac{\partial x_2}{\partial q_2} & \frac{\partial x_2}{\partial q_3} \\ \frac{\partial x_3}{\partial q_1} & \frac{\partial x_3}{\partial q_2} & \frac{\partial x_3}{\partial q_3} \end{bmatrix}$$

$$\text{Lamé coefficients: } h_j^2 = \sum_{k=1}^3 \left(\frac{\partial x_k}{\partial q_j} \right)^2$$

$$\text{Jacobian: } \frac{\partial(x_1, x_2, x_3)}{\partial(q_1, q_2, q_3)} = |J| = |\det D| = |h_1 h_2 h_3|$$

$$\text{Line element: } ds^2 = h_1^2 dq_1^2 + h_2^2 dq_2^2 + h_3^2 dq_3^2$$

$$\text{Volume element: } dV = h_1 h_2 h_3 dq_1 dq_2 dq_3$$

$$\text{Area element for surface with } q_j = \text{const} \ (i, j, k \text{ distinct}): dA = h_i h_k dq_i dq_k$$

$$\text{Gradient for scalar function } f: \nabla f = \frac{1}{h_1} \frac{\partial f}{\partial q_1} \hat{e}_{q_1} + \frac{1}{h_2} \frac{\partial f}{\partial q_2} \hat{e}_{q_2} + \frac{1}{h_3} \frac{\partial f}{\partial q_3} \hat{e}_{q_3}$$

$$\text{Divergence and curl of vector field } \vec{F} = F_{q_1} \hat{e}_{q_1} + F_{q_2} \hat{e}_{q_2} + F_{q_3} \hat{e}_{q_3}:$$

$$\nabla \cdot \vec{F} = \frac{1}{h_1 h_2 h_3} \left[\frac{\partial}{\partial q_1} (h_2 h_3 F_{q_1}) + \frac{\partial}{\partial q_2} (h_3 h_1 F_{q_2}) + \frac{\partial}{\partial q_3} (h_1 h_2 F_{q_3}) \right]$$

$$\nabla \times \vec{F} = \frac{1}{h_1 h_2 h_3} \begin{vmatrix} h_1 \hat{e}_{q_1} & h_2 \hat{e}_{q_2} & h_3 \hat{e}_{q_3} \\ \frac{\partial}{\partial q_1} & \frac{\partial}{\partial q_2} & \frac{\partial}{\partial q_3} \\ h_1 F_{q_1} & h_2 F_{q_2} & h_3 F_{q_3} \end{vmatrix}$$

Laplacian of scalar field f :

$$\nabla^2 f = \frac{1}{h_1 h_2 h_3} \left[\frac{\partial}{\partial q_1} \left(\frac{h_2 h_3}{h_1} \frac{\partial f}{\partial q_1} \right) + \frac{\partial}{\partial q_2} \left(\frac{h_3 h_1}{h_2} \frac{\partial f}{\partial q_2} \right) + \frac{\partial}{\partial q_3} \left(\frac{h_1 h_2}{h_3} \frac{\partial f}{\partial q_3} \right) \right].$$

Cylindrical Coordinates:

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z.$$

$$h_1 = h_r = 1, \quad h_2 = h_\theta = r, \quad h_3 = h_z = 1.$$

Spherical Coordinates:

$$x = \rho \cos \theta \sin \phi, \quad y = \rho \sin \theta \sin \phi, \quad z = \rho \cos \phi$$

$$h_1 = h_\rho = 1, \quad h_2 = h_\theta = \rho \sin \phi, \quad h_3 = h_\phi = \rho.$$