

Plan

- Determinant
- Determinants of Elementary Matrices
- Laplace Expansion Theorem
- Cramer's Rule
- Subspaces Associated with Matrices
- Rank Nullity Theorem
- The Fundamental Theorem of Invertible Matrices

Determinant

Determinant

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix.

- If $A = [a]$, **determinant** of A is defined as $\det(A) = a$.

Determinant

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix.

- If $A = [a]$, **determinant** of A is defined as $\det(A) = a$.
- If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then we define $\det(A) = ad - bc$.

Determinant

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix.

- If $A = [a]$, **determinant** of A is defined as $\det(A) = a$.
- If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then we define $\det(A) = ad - bc$.
- Let A_{ij} be the submatrix of A obtained by deleting the i -th row and the j -th column of A .

Determinant

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix.

- If $A = [a]$, **determinant** of A is defined as $\det(A) = a$.
- If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then we define $\det(A) = ad - bc$.
- Let A_{ij} be the submatrix of A obtained by deleting the i -th row and the j -th column of A .
- In general, $\det(A)$ is defined recursively as follows:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}).$$

- Sometimes $\det(A)$ is also denoted by $|A|$.

- Sometimes $\det(A)$ is also denoted by $|A|$.
- We define $\det(A_{ij})$ to be the (i, j) -minor of A .

- Sometimes $\det(A)$ is also denoted by $|A|$.
- We define $\det(A_{ij})$ to be the (i, j) -minor of A .
- The number $C_{ij} = (-1)^{i+j} \det(A_{ij})$ is called the (i, j) -cofactor of A .

- Sometimes $\det(A)$ is also denoted by $|A|$.
- We define $\det(A_{ij})$ to be the (i, j) -minor of A .
- The number $C_{ij} = (-1)^{i+j}\det(A_{ij})$ is called the (i, j) -cofactor of A .
- Thus we can write

$$\det(A) = |A| = \sum_{j=1}^n (-1)^{1+j} a_{1j} \det(A_{1j}) = \sum_{j=1}^n a_{1j} C_{1j}.$$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$.

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$.

● Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

● Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them.

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms?

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$?

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11}a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12}a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13}a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14}a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12}a_{12}$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12} a_{12}$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12} a_{12}$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12} a_{12} + \sigma_{13} a_{13}$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12} a_{12} + \sigma_{13} a_{13}$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11}a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12}a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13}a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14}a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12}a_{12} + \sigma_{13}a_{13}$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11}a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12}a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13}a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14}a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12}a_{12} + \sigma_{13}a_{13}$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11}a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12}a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13}a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14}a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12}a_{12} + \sigma_{13}a_{13} + \sigma_{14}a_{14}$

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?

- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12} a_{12} + \sigma_{13} a_{13} + \sigma_{14} a_{14} = (-1)^{1+2+2+3} \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix}$.

- Take a matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$. Put $\sigma_{ij} = (-1)^{i+j}$. Then $|A| =$

$$\sigma_{11} a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} + \sigma_{12} a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} + \sigma_{13} a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} + \sigma_{14} a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

- Expand each of them. Do you get 12 terms? Do you get a term $(\dots) \begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$? Do you get this matrix twice?
- So, the coefficient of $\begin{vmatrix} a_{31} & a_{34} \\ a_{41} & a_{44} \end{vmatrix}$ is $\sigma_{12} a_{12} + \sigma_{13} a_{13} = (-1)^{1+2+2+3} \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix}$.
- The coefficient of $|A_{1,2|1,3}| = \begin{vmatrix} a_{32} & a_{34} \\ a_{42} & a_{44} \end{vmatrix}$ is $(-1)^{1+2+1+3} \begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix}$.
- In general, the coefficient of $|A_{1,2|i,j}|$ in the double expansion of $|A|$ is $(-1)^{1+2+i+j} \begin{vmatrix} a_{1i} & a_{1j} \\ a_{2i} & a_{2j} \end{vmatrix}$.
- Hence $|A| = \sum_{i < j} (-1)^{1+2+i+j} \begin{vmatrix} a_{1i} & a_{1j} \\ a_{2i} & a_{2j} \end{vmatrix} |A_{1,2|i,j}|$.

Result (**Properties of Determinants**)

- If B is obtained by *interchanging* the *first two rows* of A , then $\det(B) = -\det(A)$.

Result (**Properties of Determinants**)

- If B is obtained by *interchanging* the *first two rows* of A , then $\det(B) = -\det(A)$.
- (By Induction) If B is obtained by interchanging any two *consecutive rows* of A , then $\det(B) = -\det(A)$.

Result (**Properties of Determinants**)

- If B is obtained by *interchanging* the *first two rows* of A , then $\det(B) = -\det(A)$.
- (By Induction) If B is obtained by interchanging any two *consecutive rows* of A , then $\det(B) = -\det(A)$.
- If B is obtained by interchanging *any two rows* of A , then $\det(B) = -\det(A)$.

Result (Properties of Determinants)

- If B is obtained by *interchanging the first two rows* of A , then $\det(B) = -\det(A)$.
- (By Induction) If B is obtained by interchanging any two *consecutive rows* of A , then $\det(B) = -\det(A)$.
- If B is obtained by interchanging *any two rows* of A , then $\det(B) = -\det(A)$.
- If A has a *zero row* then $\det(A) = 0$.

Result (Properties of Determinants)

- If B is obtained by *interchanging the first two rows* of A , then $\det(B) = -\det(A)$.
- (By Induction) If B is obtained by interchanging any two *consecutive rows* of A , then $\det(B) = -\det(A)$.
- If B is obtained by interchanging *any two rows* of A , then $\det(B) = -\det(A)$.
- If A has a *zero row* then $\det(A) = 0$.
- If A has two *identical rows* then $\det(A) = 0$.

Result (Properties of Determinants)

- If B is obtained by *interchanging the first two rows* of A , then $\det(B) = -\det(A)$.
- (By Induction) If B is obtained by *interchanging any two consecutive rows* of A , then $\det(B) = -\det(A)$.
- If B is obtained by *interchanging any two rows* of A , then $\det(B) = -\det(A)$.
- If A has a *zero row* then $\det(A) = 0$.
- If A has two *identical rows* then $\det(A) = 0$.
- If B is obtained by *multiplying a row* of A by k , then $\det(B) = k\det(A)$.

Result (Properties of Determinants)

- If B is obtained by *interchanging the first two rows* of A , then $\det(B) = -\det(A)$.
- (By Induction) If B is obtained by interchanging any two *consecutive rows* of A , then $\det(B) = -\det(A)$.
- If B is obtained by interchanging *any two rows* of A , then $\det(B) = -\det(A)$.
- If A has a *zero row* then $\det(A) = 0$.
- If A has two *identical rows* then $\det(A) = 0$.
- If B is obtained by *multiplying a row* of A by k , then $\det(B) = k\det(A)$.
- If B is obtained by adding a *multiple of one row* of A to another row, then $\det(B) = \det(A)$.

Result (**Determinants of Elementary Matrices**)

*Let E be an $n \times n$ elementary matrix and A be any $n \times n$ matrix.
Then*

① $\det(E) = -1, k \text{ or } 1.$

Result (**Determinants of Elementary Matrices**)

*Let E be an $n \times n$ elementary matrix and A be any $n \times n$ matrix.
Then*

❶ $\det(E) = -1, k \text{ or } 1.$

❷ $\det(EA) = \det(E)\det(A).$

Result (**Determinants of Elementary Matrices**)

Let E be an $n \times n$ elementary matrix and A be any $n \times n$ matrix. Then

1 $\det(E) = -1, k \text{ or } 1.$

2 $\det(EA) = \det(E)\det(A).$

3 E^t is also an elementary matrix and $\det(E) = \det(E^t).$

Result (**Determinants of Elementary Matrices**)

Let E be an $n \times n$ elementary matrix and A be any $n \times n$ matrix. Then

1 $\det(E) = -1, k \text{ or } 1.$

2 $\det(EA) = \det(E)\det(A).$

3 E^t is also an elementary matrix and $\det(E) = \det(E^t).$

Result

- A square matrix A is *invertible* if and only if $\det(A) \neq 0$.

Result

- A square matrix A is *invertible* if and only if $\det(A) \neq 0$.
- Let A be an $n \times n$ matrix. Then $\det(kA) = k^n \det(A)$.

Result

- A square matrix A is *invertible* if and only if $\det(A) \neq 0$.
- Let A be an $n \times n$ matrix. Then $\det(kA) = k^n \det(A)$.
- Let A and B be two $n \times n$ matrices. Then $\det(AB) = \det(A)\det(B)$.

Result

- A square matrix A is *invertible* if and only if $\det(A) \neq 0$.
- Let A be an $n \times n$ matrix. Then $\det(kA) = k^n \det(A)$.
- Let A and B be two $n \times n$ matrices. Then $\det(AB) = \det(A)\det(B)$.
- If the matrix A is invertible then $\det(A^{-1}) = \frac{1}{\det(A)}$.

Result

- A square matrix A is *invertible* if and only if $\det(A) \neq 0$.
- Let A be an $n \times n$ matrix. Then $\det(kA) = k^n \det(A)$.
- Let A and B be two $n \times n$ matrices. Then $\det(AB) = \det(A)\det(B)$.
- If the matrix A is invertible then $\det(A^{-1}) = \frac{1}{\det(A)}$.

★ A matrix A is said to be *singular* or *non-singular* according as $\det(A) = 0$ or $\det(A) \neq 0$.

Result

- The determinant of a *triangular* matrix is the product of the diagonal entries. That is, if $A = [a_{ij}]$ is an $n \times n$ *triangular* matrix then $\det(A) = a_{11} a_{22} \dots a_{nn}$.

Result

- The determinant of a *triangular* matrix is the product of the diagonal entries. That is, if $A = [a_{ij}]$ is an $n \times n$ *triangular* matrix then $\det(A) = a_{11} a_{22} \dots a_{nn}$.
- If R is in row echelon form having a *zero* row, then $\det(R) = 0 = \det(R^t)$.

Result

- The determinant of a *triangular* matrix is the product of the diagonal entries. That is, if $A = [a_{ij}]$ is an $n \times n$ *triangular* matrix then $\det(A) = a_{11} a_{22} \dots a_{nn}$.
- If R is in row echelon form having a *zero* row, then $\det(R) = 0 = \det(R^t)$.
- For any square matrix A , $\det(A^t) = \det(A)$.

Result

- The determinant of a *triangular* matrix is the product of the diagonal entries. That is, if $A = [a_{ij}]$ is an $n \times n$ *triangular* matrix then $\det(A) = a_{11} a_{22} \dots a_{nn}$.
- If R is in row echelon form having a *zero* row, then $\det(R) = 0 = \det(R^t)$.
- For any square matrix A , $\det(A^t) = \det(A)$.

★ Thus a determinant can be expanded *column-wise*.

Result

- The determinant of a *triangular* matrix is the product of the diagonal entries. That is, if $A = [a_{ij}]$ is an $n \times n$ *triangular* matrix then $\det(A) = a_{11} a_{22} \dots a_{nn}$.
- If R is in row echelon form having a *zero* row, then $\det(R) = 0 = \det(R^t)$.
- For any square matrix A , $\det(A^t) = \det(A)$.

★ Thus a determinant can be expanded *column-wise*.

★ All the previous results based on row-wise expansion of determinant are also *valid* for *column-wise* expansion.

Result (Laplace Expansion Theorem)

The determinant of an $n \times n$ matrix $A = [a_{ij}]$, where $n \geq 2$, can be computed as

$$\det(A) = a_{i1}C_{i1} + a_{i2}C_{i2} + \dots + a_{in}C_{in} = \sum_{j=1}^n a_{ij}C_{ij},$$

(this is the *cofactor expansion along the i -th row*),

Result (Laplace Expansion Theorem)

The determinant of an $n \times n$ matrix $A = [a_{ij}]$, where $n \geq 2$, can be computed as

$$\det(A) = a_{i1}C_{i1} + a_{i2}C_{i2} + \dots + a_{in}C_{in} = \sum_{j=1}^n a_{ij}C_{ij},$$

(this is the *cofactor expansion along the i -th row*),

and also as

$$\det(A) = a_{1j}C_{1j} + a_{2j}C_{2j} + \dots + a_{nj}C_{nj} = \sum_{i=1}^n a_{ij}C_{ij},$$

(this is the *cofactor expansion along the j -th column*).

Result

- If B is obtained by interchanging any two *columns* of A , then $\det(B) = -\det(A)$.

Result

- If B is obtained by interchanging any two *columns* of A , then $\det(B) = -\det(A)$.
- If A has a *zero column* then $\det(A) = 0$.

Result

- If B is obtained by interchanging any two *columns* of A , then $\det(B) = -\det(A)$.
- If A has a *zero column* then $\det(A) = 0$.
- If A has two *identical columns* then $\det(A) = 0$.

Result

- If B is obtained by interchanging any two *columns* of A , then $\det(B) = -\det(A)$.
- If A has a *zero column* then $\det(A) = 0$.
- If A has two *identical columns* then $\det(A) = 0$.
- If B is obtained by *multiplying a column* of A by k , then $\det(B) = k\det(A)$.

Result

- If B is obtained by interchanging any two *columns* of A , then $\det(B) = -\det(A)$.
- If A has a *zero column* then $\det(A) = 0$.
- If A has two *identical columns* then $\det(A) = 0$.
- If B is obtained by *multiplying a column* of A by k , then $\det(B) = k\det(A)$.
- If B is obtained by adding a *multiple of one column* of A to another column, then $\det(B) = \det(A)$.

Definition

Let A be an $n \times n$ matrix and let $\mathbf{b} \in \mathbb{R}^n$. Then $A_i(\mathbf{b})$ denotes the matrix obtained by replacing the i -th column of A by \mathbf{b} . That is, if $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$, then $A_i(\mathbf{b}) = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_{i-1} \ \mathbf{b} \ \mathbf{a}_{i+1} \ \dots \ \mathbf{a}_n]$.

Definition

Let A be an $n \times n$ matrix and let $\mathbf{b} \in \mathbb{R}^n$. Then $A_i(\mathbf{b})$ denotes the matrix obtained by replacing the i -th column of A by \mathbf{b} . That is, if $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$, then $A_i(\mathbf{b}) = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_{i-1} \ \mathbf{b} \ \mathbf{a}_{i+1} \ \dots \ \mathbf{a}_n]$.

Result (Cramer's Rule)

Let A be an $n \times n$ invertible matrix and let $\mathbf{b} \in \mathbb{R}^n$. Then the *unique* solution $\mathbf{x} = [x_1, x_2, \dots, x_n]^t$ of the system $A\mathbf{x} = \mathbf{b}$ is given by

$$x_i = \frac{\det(A_i(\mathbf{b}))}{\det(A)} \quad \text{for } i = 1, 2, \dots, n.$$

The Adjoint of a Matrix: Let $A = [a_{ij}]$ be an $n \times n$ matrix and let C_{ij} be the (i, j) -cofactor of A . Then the **adjoint** of A , denoted $\text{adj}(A)$, is defined as

$$\text{adj}(A) = \begin{bmatrix} C_{11} & C_{21} & \dots & C_{n1} \\ C_{12} & C_{22} & \dots & C_{n2} \\ \vdots & \vdots & \vdots & \vdots \\ C_{1n} & C_{2n} & \dots & C_{nn} \end{bmatrix} = [C_{ij}]^t.$$

The Adjoint of a Matrix: Let $A = [a_{ij}]$ be an $n \times n$ matrix and let C_{ij} be the (i, j) -cofactor of A . Then the **adjoint** of A , denoted $\text{adj}(A)$, is defined as

$$\text{adj}(A) = \begin{bmatrix} C_{11} & C_{21} & \dots & C_{n1} \\ C_{12} & C_{22} & \dots & C_{n2} \\ \vdots & \vdots & \vdots & \vdots \\ C_{1n} & C_{2n} & \dots & C_{nn} \end{bmatrix} = [C_{ij}]^t.$$

Result

Let A be an $n \times n$ invertible matrix. Then $A^{-1} = \frac{1}{\det(A)} \cdot \text{adj}(A)$.

The Adjoint of a Matrix: Let $A = [a_{ij}]$ be an $n \times n$ matrix and let C_{ij} be the (i, j) -cofactor of A . Then the **adjoint** of A , denoted $\text{adj}(A)$, is defined as

$$\text{adj}(A) = \begin{bmatrix} C_{11} & C_{21} & \dots & C_{n1} \\ C_{12} & C_{22} & \dots & C_{n2} \\ \vdots & \vdots & \vdots & \vdots \\ C_{1n} & C_{2n} & \dots & C_{nn} \end{bmatrix} = [C_{ij}]^t.$$

Result

Let A be an $n \times n$ invertible matrix. Then $A^{-1} = \frac{1}{\det(A)} \cdot \text{adj}(A)$.

Exercise

Use the **adjoint method** to find the **inverse** of the matrix

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 2 & 4 \\ 1 & 3 & -3 \end{bmatrix}.$$

Subspaces Associated with Matrices

Subspaces Associated with Matrices

Definition

Let A be an $m \times n$ matrix.

- 1 The **null space** of A , denoted $\text{null}(A)$, is the subspace of \mathbb{R}^n consisting of the solutions of the homogeneous linear system $A\mathbf{x} = \mathbf{0}$.

Subspaces Associated with Matrices

Definition

Let A be an $m \times n$ matrix.

- 1 The **null space** of A , denoted $\text{null}(A)$, is the subspace of \mathbb{R}^n consisting of the solutions of the homogeneous linear system $A\mathbf{x} = \mathbf{0}$. In other words,
$$\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\}$$

Subspaces Associated with Matrices

Definition

Let A be an $m \times n$ matrix.

- 1 The **null space** of A , denoted $\text{null}(A)$, is the subspace of \mathbb{R}^n consisting of the solutions of the homogeneous linear system $A\mathbf{x} = \mathbf{0}$. In other words,
$$\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\}$$
- 2 The **column space** of A , denoted $\text{col}(A)$, is the subspace of \mathbb{R}^m spanned by the columns of A .

Subspaces Associated with Matrices

Definition

Let A be an $m \times n$ matrix.

- 1 The **null space** of A , denoted $\text{null}(A)$, is the subspace of \mathbb{R}^n consisting of the solutions of the homogeneous linear system $A\mathbf{x} = \mathbf{0}$. In other words,
$$\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\}$$
- 2 The **column space** of A , denoted $\text{col}(A)$, is the subspace of \mathbb{R}^m spanned by the columns of A .
In other words, $\text{col}(A) = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\}$.

Subspaces Associated with Matrices

Definition

Let A be an $m \times n$ matrix.

- 1 The **null space** of A , denoted $\text{null}(A)$, is the subspace of \mathbb{R}^n consisting of the solutions of the homogeneous linear system $A\mathbf{x} = \mathbf{0}$. In other words,
$$\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\}$$
- 2 The **column space** of A , denoted $\text{col}(A)$, is the subspace of \mathbb{R}^m spanned by the columns of A .
In other words, $\text{col}(A) = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\}$.
- 3 The **row space** of A , denoted $\text{row}(A)$, is the subspace of \mathbb{R}^n spanned by the rows of A .

Subspaces Associated with Matrices

Definition

Let A be an $m \times n$ matrix.

- 1 The **null space** of A , denoted $\text{null}(A)$, is the subspace of \mathbb{R}^n consisting of the solutions of the homogeneous linear system $A\mathbf{x} = \mathbf{0}$. In other words,
$$\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\}$$
- 2 The **column space** of A , denoted $\text{col}(A)$, is the subspace of \mathbb{R}^m **spanned by the columns** of A .
In other words, $\text{col}(A) = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\}$.
- 3 The **row space** of A , denoted $\text{row}(A)$, is the subspace of \mathbb{R}^n **spanned by the rows** of A .
In other words, $\text{row}(A) = \{\mathbf{x}^T A \mid \mathbf{x} \in \mathbb{R}^m\}$
[Here, elements of $\text{row}(A)$ are row vectors. How can they be elements of \mathbb{R}^n ?

Subspaces Associated with Matrices

Definition

Let A be an $m \times n$ matrix.

- 1 The **null space** of A , denoted $\text{null}(A)$, is the subspace of \mathbb{R}^n consisting of the solutions of the homogeneous linear system $A\mathbf{x} = \mathbf{0}$. In other words,
$$\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\}$$
- 2 The **column space** of A , denoted $\text{col}(A)$, is the subspace of \mathbb{R}^m **spanned by the columns** of A .
In other words, $\text{col}(A) = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\}$.
- 3 The **row space** of A , denoted $\text{row}(A)$, is the subspace of \mathbb{R}^n **spanned by the rows** of A .
In other words, $\text{row}(A) = \{\mathbf{x}^T A \mid \mathbf{x} \in \mathbb{R}^m\}$

[Here, elements of $\text{row}(A)$ are row vectors. How can they be elements of \mathbb{R}^n ? In strict sense, $\text{row}(A) := \text{col}(A^T)$.]

Result

- Let B be a matrix that is *row equivalent* to the matrix A .
Then $\text{row}(B) = \text{row}(A)$.

Result

- Let B be a matrix that is *row equivalent* to the matrix A .
Then $\text{row}(B) = \text{row}(A)$.
- For any matrix A , $\text{row}(A) = \text{row}(\text{RREF}(A))$.

Result

- Let B be a matrix that is *row equivalent* to the matrix A .
Then $\text{row}(B) = \text{row}(A)$.
- For any matrix A , $\text{row}(A) = \text{row}(\text{RREF}(A))$.
- For any A , the non-zero rows of $\text{RREF}(A)$ forms a basis of $\text{row}(A)$.

Result

- Let B be a matrix that is *row equivalent* to the matrix A . Then $\text{row}(B) = \text{row}(A)$.
- For any matrix A , $\text{row}(A) = \text{row}(\text{RREF}(A))$.
- For any A , the non-zero rows of $\text{RREF}(A)$ forms a basis of $\text{row}(A)$.

Suppose A and B are row-equivalent. Are $\text{col}(A)$ and $\text{col}(B)$ equal?

Result

- Let B be a matrix that is *row equivalent* to the matrix A . Then $\text{row}(B) = \text{row}(A)$.
- For any matrix A , $\text{row}(A) = \text{row}(\text{RREF}(A))$.
- For any A , the non-zero rows of $\text{RREF}(A)$ forms a basis of $\text{row}(A)$.

Suppose A and B are row-equivalent. Are $\text{col}(A)$ and $\text{col}(B)$ equal? No.

Result

- Let B be a matrix that is *row equivalent* to the matrix A . Then $\text{row}(B) = \text{row}(A)$.
- For any matrix A , $\text{row}(A) = \text{row}(\text{RREF}(A))$.
- For any A , the non-zero rows of $\text{RREF}(A)$ forms a basis of $\text{row}(A)$.

Suppose A and B are row-equivalent. Are $\text{col}(A)$ and $\text{col}(B)$ equal? No. Take $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$.

Result

- Let B be a matrix that is *row equivalent* to the matrix A . Then $\text{row}(B) = \text{row}(A)$.
- For any matrix A , $\text{row}(A) = \text{row}(\text{RREF}(A))$.
- For any A , the non-zero rows of $\text{RREF}(A)$ forms a basis of $\text{row}(A)$.

Suppose A and B are row-equivalent. Are $\text{col}(A)$ and $\text{col}(B)$ equal? No. Take $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$.

Suppose A and B are row-equivalent. Do $\text{col}(A)$ and $\text{col}(B)$ have same dimension?

Result

- Let B be a matrix that is *row equivalent* to the matrix A . Then $\text{row}(B) = \text{row}(A)$.
- For any matrix A , $\text{row}(A) = \text{row}(\text{RREF}(A))$.
- For any A , the non-zero rows of $\text{RREF}(A)$ forms a basis of $\text{row}(A)$.

Suppose A and B are row-equivalent. Are $\text{col}(A)$ and $\text{col}(B)$ equal? No. Take $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$.

Suppose A and B are row-equivalent. Do $\text{col}(A)$ and $\text{col}(B)$ have same dimension? Yes. We will see soon.

Method of finding bases for the Row Space, the Null Space and the Column Space of a matrix:

Method of finding bases for the Row Space, the Null Space and the Column Space of a matrix:

Let A be a given matrix and let R be the reduced row echelon form of A .

Method of finding bases for the Row Space, the Null Space and the Column Space of a matrix:

Let A be a given matrix and let R be the reduced row echelon form of A .

- 1 Use the non-zero rows of R to form a basis for $\text{row}(A)$.

Method of finding bases for the Row Space, the Null Space and the Column Space of a matrix:

Let A be a given matrix and let R be the reduced row echelon form of A .

- 1 Use the non-zero rows of R to form a basis for $\text{row}(A)$.
- 2 Solve the leading variables of $R\mathbf{x} = \mathbf{0}$ in terms of the free variables, set the free variables equal to parameters, substitute back into \mathbf{x} , write the result as a linear combination of k vectors (where k is the number of free variables). These k vectors form a basis for $\text{null}(A)$.

Method of finding bases for the Row Space, the Null Space and the Column Space of a matrix:

Let A be a given matrix and let R be the reduced row echelon form of A .

- 1 Use the non-zero rows of R to form a basis for $\text{row}(A)$.
- 2 Solve the leading variables of $R\mathbf{x} = \mathbf{0}$ in terms of the free variables, set the free variables equal to parameters, substitute back into \mathbf{x} , write the result as a linear combination of k vectors (where k is the number of free variables). These k vectors form a basis for $\text{null}(A)$.
- 3 A basis for $\text{row}(A^t)$ will also be a basis for $\text{col}(A)$.

Method of finding bases for the Row Space, the Null Space and the Column Space of a matrix:

Let A be a given matrix and let R be the reduced row echelon form of A .

- 1 Use the non-zero rows of R to form a basis for $\text{row}(A)$.
- 2 Solve the leading variables of $R\mathbf{x} = \mathbf{0}$ in terms of the free variables, set the free variables equal to parameters, substitute back into \mathbf{x} , write the result as a linear combination of k vectors (where k is the number of free variables). These k vectors form a basis for $\text{null}(A)$.
- 3 A basis for $\text{row}(A^t)$ will also be a basis for $\text{col}(A)$.
- 4 Or, Use the columns of A that correspond to the columns of R containing the leading 1's to form a basis for $\text{col}(A)$.

Subspaces Associated with Matrices

Subspaces Associated with Matrices

Example

Find bases for the *row space*, *column space* and *null space* of the following matrix:

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 2 & 4 \\ 4 & 6 & 2 \end{bmatrix},$$

Subspaces Associated with Matrices

Example

Find bases for the *row space*, *column space* and *null space* of the following matrix:

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 2 & 4 \\ 4 & 6 & 2 \end{bmatrix}, \quad RREF(A) = \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{bmatrix}.$$

Subspaces Associated with Matrices

Example

Find bases for the *row space*, *column space* and *null space* of the following matrix:

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & 2 & 4 \\ 4 & 6 & 2 \end{bmatrix}, \quad RREF(A) = \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & -3 \\ 0 & 0 & 0 \end{bmatrix}.$$

Result

Let $R = [\mathbf{b}_1 \ \mathbf{b}_2 \ \dots \ \mathbf{b}_n]$ be the reduced row echelon form of a matrix $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$ of rank r . Let $\mathbf{b}_{j_1}, \mathbf{b}_{j_2}, \dots, \mathbf{b}_{j_r}$ be the columns of R such that $\mathbf{b}_{j_k} = \mathbf{e}_k$ for $k = 1, \dots, r$. Then $\{\mathbf{a}_{j_1}, \mathbf{a}_{j_2}, \dots, \mathbf{a}_{j_r}\}$ is a basis for $\text{col}(A)$.

Result

The row space and the column space of a matrix A have the same dimension, and $\dim(\text{row}(A)) = \dim(\text{col}(A)) = \text{rank}(A)$.

Result

The row space and the column space of a matrix A have the same dimension, and $\dim(\text{row}(A)) = \dim(\text{col}(A)) = \text{rank}(A)$.

Result

For any matrix A , we have $\text{rank}(A^t) = \text{rank}(A)$.

Result

The row space and the column space of a matrix A have the same dimension, and $\dim(\text{row}(A)) = \dim(\text{col}(A)) = \text{rank}(A)$.

Result

For any matrix A , we have $\text{rank}(A^t) = \text{rank}(A)$.

Nullity: The nullity of a matrix A is the dimension of its null space, and is denoted by $\text{nullity}(A)$.

Result

The row space and the column space of a matrix A have the same dimension, and $\dim(\text{row}(A)) = \dim(\text{col}(A)) = \text{rank}(A)$.

Result

For any matrix A , we have $\text{rank}(A^t) = \text{rank}(A)$.

Nullity: The nullity of a matrix A is the dimension of its null space, and is denoted by $\text{nullity}(A)$.

Result (Rank Nullity Theorem)

Let A be an $m \times n$ matrix. Then

$$\text{rank}(A) + \text{nullity}(A) = n.$$

Result (Fundamental Theorem of Invertible Matrices: II)

Result (Fundamental Theorem of Invertible Matrices: II)

Let A be an $n \times n$ matrix. Then the following statements are equivalent.

Result (Fundamental Theorem of Invertible Matrices: II)

Let A be an $n \times n$ matrix. Then the following statements are equivalent.

- 1 A is *invertible*.
- 2 A^t is *invertible*.
- 3 $A\mathbf{x} = \mathbf{b}$ has a *solution* for every \mathbf{b} in \mathbb{R}^n .
- 4 $A\mathbf{x} = \mathbf{b}$ has a *unique solution* for every \mathbf{b} in \mathbb{R}^n .
- 5 $A\mathbf{x} = \mathbf{0}$ has only the *trivial solution*.
- 6 The reduced row echelon form of A is I_n .
- 7 The rows of A are linearly independent.
- 8 The columns of A are linearly independent.
- 9 $\text{rank}(A) = n$.
- 10 A is a *product of* elementary matrices.

11. $\text{nullity}(A) = 0$.

11. $\text{nullity}(A) = 0$.

12. The column vectors of A span \mathbb{R}^n .

11. $\text{nullity}(A) = 0$.

12. The column vectors of A span \mathbb{R}^n .

13. The column vectors of A form a basis for \mathbb{R}^n .

11. $\text{nullity}(A) = 0$.

12. The column vectors of A span \mathbb{R}^n .

13. The column vectors of A form a basis for \mathbb{R}^n .

14. The row vectors of A span \mathbb{R}^n .

- 11. $\text{nullity}(A) = 0$.
- 12. The column vectors of A span \mathbb{R}^n .
- 13. The column vectors of A form a basis for \mathbb{R}^n .
- 14. The row vectors of A span \mathbb{R}^n .
- 15. The row vectors of A form a basis for \mathbb{R}^n .

Exercise

Show that the vectors $[1, 2, 3]^t$, $[-1, 0, 1]^t$ and $[4, 9, 7]^t$ form a basis for \mathbb{R}^3 .

Exercise

Show that the vectors $[1, 2, 3]^t$, $[-1, 0, 1]^t$ and $[4, 9, 7]^t$ form a basis for \mathbb{R}^3 .

Result

Let A be an $m \times n$ matrix. Then

Exercise

Show that the vectors $[1, 2, 3]^t$, $[-1, 0, 1]^t$ and $[4, 9, 7]^t$ form a basis for \mathbb{R}^3 .

Result

Let A be an $m \times n$ matrix. Then

1 $\text{rank}(A^t A) = \text{rank}(A)$.

Exercise

Show that the vectors $[1, 2, 3]^t$, $[-1, 0, 1]^t$ and $[4, 9, 7]^t$ form a basis for \mathbb{R}^3 .

Result

Let A be an $m \times n$ matrix. Then

- 1 $\text{rank}(A^t A) = \text{rank}(A)$.
- 2 The $n \times n$ matrix $A^t A$ is invertible if and only if $\text{rank}(A) = n$.

Result

Let A, B, T, S be matrices, where T and S are *invertible*.

- 1 If TA and AS are defined, then
$$\text{rank}(TA) = \text{rank}(A) = \text{rank}(AS).$$

Result

Let A, B, T, S be matrices, where T and S are *invertible*.

- 1 If TA and AS are defined, then
$$\text{rank}(TA) = \text{rank}(A) = \text{rank}(AS).$$
- 2 If AB is defined then $\text{rank}(AB) \leq \min\{\text{rank}(A), \text{rank}(B)\}.$

Result

Let A, B, T, S be matrices, where T and S are *invertible*.

- 1 If TA and AS are defined, then
 $\text{rank}(TA) = \text{rank}(A) = \text{rank}(AS)$.
- 2 If AB is defined then $\text{rank}(AB) \leq \min\{\text{rank}(A), \text{rank}(B)\}$.
- 3 If $A + B$ is defined then $\text{rank}(A + B) \leq \text{rank}(A) + \text{rank}(B)$.

Result

Let A, B, T, S be matrices, where T and S are *invertible*.

- 1 If TA and AS are defined, then
 $\text{rank}(TA) = \text{rank}(A) = \text{rank}(AS)$.
- 2 If AB is defined then $\text{rank}(AB) \leq \min\{\text{rank}(A), \text{rank}(B)\}$.
- 3 If $A + B$ is defined then $\text{rank}(A + B) \leq \text{rank}(A) + \text{rank}(B)$.

Result

Let A be an $n \times n$ matrix of *rank* r , where $1 \leq r < n$. Show that there exist elementary matrices E_1, \dots, E_p and F_1, \dots, F_q such

that $E_1 \dots E_p A F_1 \dots F_q = \begin{bmatrix} I_r & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}$.