

Plan

- Eigenvalues and Eigenvectors
- Similar Matrix
- Diagonalizable Matrix

Eigenvalues and Eigenvectors

Eigenvalues and Eigenvectors

- Just like the space \mathbb{R}^n , we also define the space \mathbb{C}^n .

Eigenvalues and Eigenvectors

- Just like the space \mathbb{R}^n , we also define the space \mathbb{C}^n .
- Indeed,

$$\mathbb{C}^n = \{[x_1, x_2, \dots, x_n]^t : x_1, x_2, \dots, x_n \in \mathbb{C}\}.$$

Eigenvalues and Eigenvectors

- Just like the space \mathbb{R}^n , we also define the space \mathbb{C}^n .
- Indeed,

$$\mathbb{C}^n = \{[x_1, x_2, \dots, x_n]^t : x_1, x_2, \dots, x_n \in \mathbb{C}\}.$$

The definitions of vector **addition** and scalar **multiplication** *etc.*, and most of the results that we have studied so far in case of \mathbb{R}^n , can also be accomplished for the space \mathbb{C}^n , **in a similar manner**.

Definition

Let A be an $n \times n$ matrix.

- A complex number λ is called an **eigenvalue** of A if there is $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ such that $A\mathbf{x} = \lambda\mathbf{x}$.

Definition

Let A be an $n \times n$ matrix.

- A complex number λ is called an **eigenvalue** of A if there is $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ such that $A\mathbf{x} = \lambda\mathbf{x}$.
- We call \mathbf{x} an **eigenvector** of A corresponding to λ .

Definition

Let A be an $n \times n$ matrix.

- A complex number λ is called an **eigenvalue** of A if there is $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ such that $A\mathbf{x} = \lambda\mathbf{x}$.
- We call \mathbf{x} an **eigenvector** of A **corresponding to** λ .

Example

For $\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$, eigenvalues: $4, -2$; resp. eigenvectors $\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$.

Definition

Let A be an $n \times n$ matrix.

- A complex number λ is called an **eigenvalue** of A if there is $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ such that $A\mathbf{x} = \lambda\mathbf{x}$.
- We call \mathbf{x} an **eigenvector** of A **corresponding to** λ .

Example

For $\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$, eigenvalues: $4, -2$; resp. eigenvectors $\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$.

Result

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . Then

- λ is an eigenvalue of A **iff** $\det(A - \lambda I) = 0$.

Definition

Let A be an $n \times n$ matrix.

- A complex number λ is called an **eigenvalue** of A if there is $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ such that $A\mathbf{x} = \lambda\mathbf{x}$.
- We call \mathbf{x} an **eigenvector** of A **corresponding to λ** .

Example

For $\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$, eigenvalues: $4, -2$; resp. eigenvectors $\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$.

Result

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . Then

- λ is an eigenvalue of A **iff** $\det(A - \lambda I) = 0$.
- 0 is an **eigenvalue** of A **iff** A is **not invertible**.

Definition

Let A be an $n \times n$ matrix.

- A complex number λ is called an **eigenvalue** of A if there is $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ such that $A\mathbf{x} = \lambda\mathbf{x}$.
- We call \mathbf{x} an **eigenvector** of A **corresponding to** λ .

Example

For $\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$, eigenvalues: $4, -2$; resp. eigenvectors $\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$.

Result

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . Then

- λ is an eigenvalue of A **iff** $\det(A - \lambda I) = 0$.
- 0 is an **eigenvalue** of A **iff** A is **not invertible**.
- ★ The eigenvalues can be directly computed for 2×2 matrices.

Definition

Let A be an $n \times n$ matrix.

- A complex number λ is called an **eigenvalue** of A if there is $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ such that $A\mathbf{x} = \lambda\mathbf{x}$.
- We call \mathbf{x} an **eigenvector** of A **corresponding to** λ .

Example

For $\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$, eigenvalues: $4, -2$; resp. eigenvectors $\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$.

Result

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . Then

- λ is an eigenvalue of A **iff** $\det(A - \lambda I) = 0$.
- 0 is an **eigenvalue** of A **iff** A is **not invertible**.
- ★ The eigenvalues can be directly computed for 2×2 matrices.
- ★ Use **Gauss Jordan Elimination** on $[A - \lambda I | \mathbf{0}]$ to find all possible eigenvectors for λ of A .

Definition

Let λ be an eigenvalue of a matrix A . The collection of all eigenvectors of A corresponding to λ , together with the zero vector, is called the eigenspace of λ , and is denoted by E_λ .

Definition

Let λ be an eigenvalue of a matrix A . The collection of all eigenvectors of A corresponding to λ , together with the zero vector, is called the eigenspace of λ , and is denoted by E_λ .

Result

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . Then

- $E_\lambda = \text{null}(A - \lambda I)$, that is, E_λ is a subspace of \mathbb{C}^n .

Definition

Let λ be an eigenvalue of a matrix A . The collection of all eigenvectors of A corresponding to λ , together with the zero vector, is called the eigenspace of λ , and is denoted by E_λ .

Result

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . Then

- $E_\lambda = \text{null}(A - \lambda I)$, that is, E_λ is a subspace of \mathbb{C}^n .
- Let $\mathbf{v}_1, \dots, \mathbf{v}_k$ be eigenvectors of A corresponding to λ and $\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_k \mathbf{v}_k \neq \mathbf{0}$. Then \mathbf{v} is an eigenvector of A .

Definition

Let λ be an eigenvalue of a matrix A . The collection of all eigenvectors of A corresponding to λ , together with the zero vector, is called the eigenspace of λ , and is denoted by E_λ .

Result

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . Then

- $E_\lambda = \text{null}(A - \lambda I)$, that is, E_λ is a subspace of \mathbb{C}^n .
- Let $\mathbf{v}_1, \dots, \mathbf{v}_k$ be eigenvectors of A corresponding to λ and $\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_k \mathbf{v}_k \neq \mathbf{0}$. Then \mathbf{v} is an eigenvector of A .
- Eigenvalues of a triangular matrix are its diagonal entries.

Definition

Let λ be an eigenvalue of a matrix A . The collection of all eigenvectors of A corresponding to λ , together with the zero vector, is called the eigenspace of λ , and is denoted by E_λ .

Result

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . Then

- $E_\lambda = \text{null}(A - \lambda I)$, that is, E_λ is a subspace of \mathbb{C}^n .
- Let $\mathbf{v}_1, \dots, \mathbf{v}_k$ be eigenvectors of A corresponding to λ and $\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_k \mathbf{v}_k \neq \mathbf{0}$. Then \mathbf{v} is an eigenvector of A .
- Eigenvalues of a triangular matrix are its diagonal entries.
- Eigenvalues of $\left[\begin{array}{c|c} A_p & C \\ \hline O & B_q \end{array} \right]$ are the eigenvalues of A and B .

Definition

Let A be an $n \times n$ matrix. Then

- $P_A(x) = |A - xI|$ is called **characteristic polynomial** of A .

Definition

Let A be an $n \times n$ matrix. Then

- $P_A(x) = |A - xI|$ is called **characteristic polynomial** of A .
- $P_A(x) = 0$ is called **characteristic equation** of A .

Definition

Let A be an $n \times n$ matrix. Then

- $P_A(x) = |A - xI|$ is called **characteristic polynomial** of A .
- $P_A(x) = 0$ is called **characteristic equation** of A .

Example

Take $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -5 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 0 & -3 \\ 1 & 0 & -1 \end{bmatrix}$.

Definition

Let A be an $n \times n$ matrix. Then

- $P_A(x) = |A - xI|$ is called **characteristic polynomial** of A .
- $P_A(x) = 0$ is called **characteristic equation** of A .

Example

Take $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -5 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 0 & -3 \\ 1 & 0 & -1 \end{bmatrix}$.

- $P_A(x) = -(x-1)^2(x-2)$; $E_1 = \text{span}\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}\right)$, $E_2 = \text{span}\left(\begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}\right)$.

Definition

Let A be an $n \times n$ matrix. Then

- $P_A(x) = |A - xI|$ is called **characteristic polynomial** of A .
- $P_A(x) = 0$ is called **characteristic equation** of A .

Example

Take $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -5 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 0 & -3 \\ 1 & 0 & -1 \end{bmatrix}$.

- $P_A(x) = -(x-1)^2(x-2)$; $E_1 = \text{span}\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}\right)$, $E_2 = \text{span}\left(\begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}\right)$.
- $P_B(x) = -x^2(x+2)$; $E_0 = \text{span}\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}\right)$, $E_{-2} = \text{span}\left(\begin{bmatrix} -1 \\ 3 \\ 1 \end{bmatrix}\right)$.

Result (Fundamental Theorem of Invertible Matrices: Version III)

Result (Fundamental Theorem of Invertible Matrices: Version III)

Let A be an $n \times n$ matrix. Then the following statements are equivalent.

Result (Fundamental Theorem of Invertible Matrices: Version III)

Let A be an $n \times n$ matrix. Then the following statements are equivalent.

1. A is *invertible*.
2. A^t is *invertible*.
3. $A\mathbf{x} = \mathbf{b}$ has a *solution* for every \mathbf{b} in \mathbb{R}^n .
4. $A\mathbf{x} = \mathbf{b}$ has a *unique solution* for every \mathbf{b} in \mathbb{R}^n .
5. $A\mathbf{x} = \mathbf{0}$ has only the *trivial solution*.
6. The reduced row echelon form of A is I_n .
7. The rows of A are linearly independent.
8. The columns of A are linearly independent.

- 9. $\text{rank}(A) = n$.
- 10. A is a product of elementary matrices.
- 11. $\text{nullity}(A) = 0$.
- 12. The column vectors of A span \mathbb{R}^n .
- 13. The column vectors of A form a basis for \mathbb{R}^n .
- 14. The row vectors of A span \mathbb{R}^n .
- 15. The row vectors of A form a basis for \mathbb{R}^n .

- 9. $\text{rank}(A) = n$.
- 10. A is a **product of** elementary matrices.
- 11. $\text{nullity}(A) = 0$.
- 12. The column vectors of A span \mathbb{R}^n .
- 13. The column vectors of A form a basis for \mathbb{R}^n .
- 14. The row vectors of A span \mathbb{R}^n .
- 15. The row vectors of A form a basis for \mathbb{R}^n .
- 16. **$\det A \neq 0$.**

- 9. $\text{rank}(A) = n$.
- 10. A is a product of elementary matrices.
- 11. $\text{nullity}(A) = 0$.
- 12. The column vectors of A span \mathbb{R}^n .
- 13. The column vectors of A form a basis for \mathbb{R}^n .
- 14. The row vectors of A span \mathbb{R}^n .
- 15. The row vectors of A form a basis for \mathbb{R}^n .
- 16. $\det A \neq 0$.
- 17. 0 is not an eigenvalue of A .

Result

Let A be a matrix with eigenvalue λ and corresponding eigenvector \mathbf{x} .

Result

Let A be a matrix with eigenvalue λ and corresponding eigenvector \mathbf{x} .

- 1 For any positive integer n , λ^n is an eigenvalue of A^n with corresponding eigenvector \mathbf{x} .

Result

Let A be a matrix with eigenvalue λ and corresponding eigenvector \mathbf{x} .

- 1 For any positive integer n , λ^n is an eigenvalue of A^n with corresponding eigenvector \mathbf{x} .
- 2 If A is invertible, then $\frac{1}{\lambda}$ is an eigenvalue of A^{-1} with corresponding eigenvector \mathbf{x} .

Result

Let A be a matrix with eigenvalue λ and corresponding eigenvector \mathbf{x} .

- 1 For any positive integer n , λ^n is an eigenvalue of A^n with corresponding eigenvector \mathbf{x} .
- 2 If A is invertible, then $\frac{1}{\lambda}$ is an eigenvalue of A^{-1} with corresponding eigenvector \mathbf{x} .
- 3 If A is invertible then for any integer n , λ^n is an eigenvalue of A^n with corresponding eigenvector \mathbf{x} .

Result

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ be eigenvectors of a matrix A with corresponding eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_m$, respectively. Let $\mathbf{x} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_m\mathbf{v}_m$. Then for any positive integer k ,

$$A^k \mathbf{x} = c_1 \lambda_1^k \mathbf{v}_1 + c_2 \lambda_2^k \mathbf{v}_2 + \dots + c_m \lambda_m^k \mathbf{v}_m.$$

Result

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ be eigenvectors of a matrix A with corresponding eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_m$, respectively. Let $\mathbf{x} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_m\mathbf{v}_m$. Then for any positive integer k ,

$$A^k \mathbf{x} = c_1 \lambda_1^k \mathbf{v}_1 + c_2 \lambda_2^k \mathbf{v}_2 + \dots + c_m \lambda_m^k \mathbf{v}_m.$$

Result

Let $\lambda_1, \lambda_2, \dots, \lambda_m$ be *distinct* eigenvalues of a matrix A with corresponding eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$, respectively. Then the set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$ is *linearly independent*.

Similar Matrix

Similar Matrix

Let A and B be two $n \times n$ matrices. Then A is said to be similar to B if there is an $n \times n$ invertible matrix T such that $T^{-1}AT = B$.

Similar Matrix

Let A and B be two $n \times n$ matrices. Then A is said to be similar to B if there is an $n \times n$ invertible matrix T such that $T^{-1}AT = B$.

- If A is similar to B , we write $A \approx B$.

Similar Matrix

Let A and B be two $n \times n$ matrices. Then A is said to be similar to B if there is an $n \times n$ invertible matrix T such that $T^{-1}AT = B$.

- If A is similar to B , we write $A \approx B$.
- If $A \approx B$, we can equivalently write that $A = TBT^{-1}$ or $AT = TB$.

Example

Let $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ and $T = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$.

Then $A \approx B$ since $AT = TB$.

Example

Let $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ and $T = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$.

Then $A \approx B$ since $AT = TB$.

Result

Let A, B and C be $n \times n$ matrices. Then

1 $A \approx A$.

Example

Let $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ and $T = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$.

Then $A \approx B$ since $AT = TB$.

Result

Let A, B and C be $n \times n$ matrices. Then

① $A \approx A$.

② If $A \approx B$ then $B \approx A$.

Example

Let $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ and $T = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$.

Then $A \approx B$ since $AT = TB$.

Result

Let A, B and C be $n \times n$ matrices. Then

- 1 $A \approx A$.
- 2 If $A \approx B$ then $B \approx A$.
- 3 If $A \approx B$ and $B \approx C$ then $A \approx C$.

Result

Let A, B, T be matrices such that T is *invertible* and $B = T^{-1}AT$. Then

Result

Let A, B, T be matrices such that T is *invertible* and $B = T^{-1}AT$. Then

1 $\det A = \det B.$

Result

Let A, B, T be matrices such that T is *invertible* and $B = T^{-1}AT$. Then

- 1 $\det A = \det B$.
- 2 A is invertible *iff* B is invertible.

Result

Let A, B, T be matrices such that T is *invertible* and $B = T^{-1}AT$. Then

- 1 $\det A = \det B$.
- 2 A is invertible *iff* B is invertible.
- 3 A and B have the *same rank*.

Result

Let A, B, T be matrices such that T is *invertible* and $B = T^{-1}AT$. Then

- 1 $\det A = \det B$.
- 2 A is invertible *iff* B is invertible.
- 3 A and B have the *same rank*.
- 4 A and B have the *same characteristic polynomial*.

Result

Let A, B, T be matrices such that T is *invertible* and $B = T^{-1}AT$. Then

- 1 $\det A = \det B$.
- 2 A is invertible *iff* B is invertible.
- 3 A and B have the *same rank*.
- 4 A and B have the *same characteristic polynomial*.
- 5 A and B have the *same* set of eigenvalues.

Result

Let A, B, T be matrices such that T is *invertible* and $B = T^{-1}AT$. Then

- 1 $\det A = \det B$.
 - 2 A is invertible *iff* B is invertible.
 - 3 A and B have the *same rank*.
 - 4 A and B have the *same characteristic polynomial*.
 - 5 A and B have the *same* set of eigenvalues.
- ★ λ is an eigenvalue of B with an eigenvector \mathbf{v} *iff* λ is an eigenvalue of A with an eigenvector $T\mathbf{v}$.

Result

Let A, B, T be matrices such that T is *invertible* and $B = T^{-1}AT$. Then

- 1 $\det A = \det B$.
- 2 A is invertible *iff* B is invertible.
- 3 A and B have the *same rank*.
- 4 A and B have the *same characteristic polynomial*.
- 5 A and B have the *same* set of eigenvalues.
- ★ λ is an eigenvalue of B with an eigenvector \mathbf{v} *iff* λ is an eigenvalue of A with an eigenvector $T\mathbf{v}$.
- 6 The $\dim(E_\lambda)$ for A is same as $\dim(E_\lambda)$ for B .

Diagonalizable Matrix

Diagonalizable Matrix

A matrix A is said to be diagonalizable if there is a diagonal matrix D such that $A \approx D$.

Diagonalizable Matrix

A matrix A is said to be **diagonalizable** if there is a **diagonal** matrix D such that $A \approx D$.

Example

The matrix $A = \begin{bmatrix} 1 & 3 \\ 2 & 2 \end{bmatrix}$ is diagonalizable, since if $D = \begin{bmatrix} 4 & 0 \\ 0 & -1 \end{bmatrix}$ and $T = \begin{bmatrix} 1 & 3 \\ 1 & -2 \end{bmatrix}$ then $AT = TD$.

Diagonalizable Matrix

A matrix A is said to be **diagonalizable** if there is a **diagonal** matrix D such that $A \approx D$.

Example

The matrix $A = \begin{bmatrix} 1 & 3 \\ 2 & 2 \end{bmatrix}$ is diagonalizable, since if $D = \begin{bmatrix} 4 & 0 \\ 0 & -1 \end{bmatrix}$ and $T = \begin{bmatrix} 1 & 3 \\ 1 & -2 \end{bmatrix}$ then $AT = TD$.

Result

Let A be an $n \times n$ matrix. Then A is diagonalizable **iff** A has **n linearly independent eigenvectors**.

- Let A be a diagonalizable matrix.

- Let A be a diagonalizable matrix.
- Suppose $T^{-1}AT = D$, where T is an invertible matrix and D is a diagonal matrix.

- Let A be a diagonalizable matrix.
- Suppose $T^{-1}AT = D$, where T is an invertible matrix and D is a diagonal matrix.
- Then the columns of T are the linearly independent eigenvectors of A .

- Let A be a diagonalizable matrix.
- Suppose $T^{-1}AT = D$, where T is an invertible matrix and D is a diagonal matrix.
- Then the columns of T are the linearly independent eigenvectors of A .
- The diagonal entries of D are the eigenvalues of A corresponding to the columns (eigenvectors of A) of T in the same order.

Example

Check for the *diagonalizability*. If diagonalizable, find a T that diagonalizes it. [Use GJE]

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -5 & 4 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 0 & -3 \\ 1 & 0 & -1 \end{bmatrix}.$$

Example

Check for the *diagonalizability*. If diagonalizable, find a T that diagonalizes it. [Use GJE]

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -5 & 4 \end{bmatrix} \text{ and } B = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 0 & -3 \\ 1 & 0 & -1 \end{bmatrix}.$$

- For A , $E_1 = \text{span}\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}\right)$, $E_2 = \text{span}\left(\begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}\right)$.
- For B , $E_0 = \text{span}\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}\right)$, $E_{-2} = \text{span}\left(\begin{bmatrix} -1 \\ 3 \\ 1 \end{bmatrix}\right)$.

Example

Check for the *diagonalizability*. If diagonalizable, find a T that diagonalizes it. [Use GJE]

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -5 & 4 \end{bmatrix} \text{ and } B = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 0 & -3 \\ 1 & 0 & -1 \end{bmatrix}.$$

- For A , $E_1 = \text{span}\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}\right)$, $E_2 = \text{span}\left(\begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}\right)$.
- For B , $E_0 = \text{span}\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}\right)$, $E_{-2} = \text{span}\left(\begin{bmatrix} -1 \\ 3 \\ 1 \end{bmatrix}\right)$.
- A is **not** diagonalizable.

Example

Check for the *diagonalizability*. If diagonalizable, find a T that diagonalizes it. [Use GJE]

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -5 & 4 \end{bmatrix} \text{ and } B = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 0 & -3 \\ 1 & 0 & -1 \end{bmatrix}.$$

- For A , $E_1 = \text{span}\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}\right)$, $E_2 = \text{span}\left(\begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}\right)$.

- For B , $E_0 = \text{span}\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}\right)$, $E_{-2} = \text{span}\left(\begin{bmatrix} -1 \\ 3 \\ 1 \end{bmatrix}\right)$.

- A is **not** diagonalizable. B is diagonalizable, $T = \begin{bmatrix} 0 & 1 & -1 \\ 1 & 0 & 3 \\ 0 & 1 & 1 \end{bmatrix}$.

Result

If A is an $n \times n$ matrix with n distinct eigenvalues then A is diagonalizable.

Result

If A is an $n \times n$ matrix with n distinct eigenvalues then A is diagonalizable.

Result

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be distinct eigenvalues of a matrix A . If \mathcal{B}_i is a basis for the eigenspace E_{λ_i} , then $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2 \cup \dots \cup \mathcal{B}_k$ is a linearly independent set.

Result

If A is an $n \times n$ matrix with n distinct eigenvalues then A is diagonalizable.

Result

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be distinct eigenvalues of a matrix A . If \mathcal{B}_i is a basis for the eigenspace E_{λ_i} , then $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2 \cup \dots \cup \mathcal{B}_k$ is a linearly independent set.

Definition

Let λ be an eigenvalue of a matrix A .

- The algebraic multiplicity of λ is the multiplicity of λ as a root of the characteristic polynomial of A .

Result

If A is an $n \times n$ matrix with n distinct eigenvalues then A is diagonalizable.

Result

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be distinct eigenvalues of a matrix A . If \mathcal{B}_i is a basis for the eigenspace E_{λ_i} , then $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2 \cup \dots \cup \mathcal{B}_k$ is a linearly independent set.

Definition

Let λ be an eigenvalue of a matrix A .

- The algebraic multiplicity of λ is the multiplicity of λ as a root of the characteristic polynomial of A .
- The geometric multiplicity of λ is the dimension of E_λ .

Diagonalizable Matrix

Result

The *geometric multiplicity* of each eigenvalue of a matrix is *less than or equal* to its *algebraic multiplicity*.

Diagonalizable Matrix

Result

The *geometric multiplicity* of each eigenvalue of a matrix is *less than or equal* to its *algebraic multiplicity*.

Result (The Diagonalization Theorem)

Let A be an $n \times n$ matrix whose *distinct eigenvalues* are $\lambda_1, \lambda_2, \dots, \lambda_k$. Then the following statements are equivalent:

Diagonalizable Matrix

Result

The *geometric multiplicity* of each eigenvalue of a matrix is *less than or equal* to its *algebraic multiplicity*.

Result (The Diagonalization Theorem)

Let A be an $n \times n$ matrix whose *distinct eigenvalues* are $\lambda_1, \lambda_2, \dots, \lambda_k$. Then the following statements are equivalent:

- 1 A is diagonalizable.
- 2 The union \mathcal{B} of the bases of the eigenspaces of A contains n vectors.
- 3 The *algebraic multiplicity* of each eigenvalue *equals* its *geometric multiplicity*.