

Quantum Mechanics
Tutorial-3

1) A particle of mass m is confined to a one dimensional region $0 \leq x \leq a$. The energy eigen states and the corresponding eigen values are given by

$$\begin{aligned}\chi_n &= \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right) \\ E_n &= \frac{n^2 \pi^2 \hbar^2}{2ma^2}.\end{aligned}$$

The normalized wave function of the particle at $t = 0$ is given by

$$\psi(x, 0) = \sqrt{\frac{8}{5\pi}} \left[1 + \cos \frac{\pi x}{a} \right] \sin \frac{\pi x}{a}.$$

- (a) Find the wave function at any time t , $\psi(x, t)$.
- (b) Calculate the average energy at $t = 0$ and $t = t_0$.
- (c) What is the probability that the particle will be found in the left half of the box.

2) The time independent wave function of a free particle in a box is

$$\psi(x) = Ax(a - x) \quad 0 \leq x \leq a.$$

- (a) Calculate A
- (b) Calculate the average energy of the particle.
- (c) Calculate the corresponding momentum wave function.
- (d) What is the probability of finding the particle at energy $E_1 = \frac{\pi^2 \hbar^2}{2ma^2}$.

Normalized energy eigenstates and the corresponding eigen values are as given in problem (1).

3) For a free particle the momentum space wave function at $t = 0$ is given by

$$\phi(p, 0) = Ae^{-\alpha p^2},$$

where α is a constant.

- (a) Find A .
- (b) Find $\psi(x, t)$.

4) Using $[\hat{x}, \hat{p}] = i\hbar$ show that

- (a) $[\hat{x}^2, \hat{p}] = 2i\hbar\hat{x}$.
- (b) Also, prove that $[\hat{x}^n, \hat{p}] = ni\hbar\hat{x}^{n-1}$.