

Plan

- Schur Unitary Triangularization
- Spectral Theorem
- Cayley-Hamilton Theorem

Let $A \in \mathcal{M}_n(\mathbb{C})$, the set of all $n \times n$ complex matrices.

Let $A \in \mathcal{M}_n(\mathbb{C})$, the set of all $n \times n$ complex matrices.

- $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ (a multi-set), where $\lambda_1, \dots, \lambda_n$ are the eigenvalues (not necessarily distinct) of A .

Let $A \in \mathcal{M}_n(\mathbb{C})$, the set of all $n \times n$ complex matrices.

- $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ (a multi-set), where $\lambda_1, \dots, \lambda_n$ are the eigenvalues (not necessarily distinct) of A . This $\sigma(A)$ is called the spectrum of A .

Let $A \in \mathcal{M}_n(\mathbb{C})$, the set of all $n \times n$ complex matrices.

- $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ (a multi-set), where $\lambda_1, \dots, \lambda_n$ are the eigenvalues (not necessarily distinct) of A . This $\sigma(A)$ is called the spectrum of A .
- If λ is an eigenvalue of A with corresponding eigenvector \mathbf{v} , then (λ, \mathbf{v}) is called an eigenpair of A .

Let $A \in \mathcal{M}_n(\mathbb{C})$, the set of all $n \times n$ complex matrices.

- $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ (a multi-set), where $\lambda_1, \dots, \lambda_n$ are the eigenvalues (not necessarily distinct) of A . This $\sigma(A)$ is called the spectrum of A .
- If λ is an eigenvalue of A with corresponding eigenvector \mathbf{v} , then (λ, \mathbf{v}) is called an eigenpair of A .
- If $A = (a_{ij})$, then $\text{Tr}(A) = a_{11} + \dots + a_{nn}$ is called the trace of A .

Let $A \in \mathcal{M}_n(\mathbb{C})$, the set of all $n \times n$ complex matrices.

- $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ (a multi-set), where $\lambda_1, \dots, \lambda_n$ are the eigenvalues (not necessarily distinct) of A . This $\sigma(A)$ is called the spectrum of A .
- If λ is an eigenvalue of A with corresponding eigenvector \mathbf{v} , then (λ, \mathbf{v}) is called an eigenpair of A .
- If $A = (a_{ij})$, then $\text{Tr}(A) = a_{11} + \dots + a_{nn}$ is called the trace of A .
- A is called a normal matrix if $A^*A = AA^*$. For example,
 $\begin{bmatrix} i & 0 \\ 0 & 2 \end{bmatrix}, \begin{bmatrix} i & 2 \\ 2 & i \end{bmatrix}.$

Let $A \in \mathcal{M}_n(\mathbb{C})$, the set of all $n \times n$ complex matrices.

- $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ (a multi-set), where $\lambda_1, \dots, \lambda_n$ are the eigenvalues (not necessarily distinct) of A . This $\sigma(A)$ is called the spectrum of A .
- If λ is an eigenvalue of A with corresponding eigenvector \mathbf{v} , then (λ, \mathbf{v}) is called an eigenpair of A .
- If $A = (a_{ij})$, then $\text{Tr}(A) = a_{11} + \dots + a_{nn}$ is called the trace of A .
- A is called a normal matrix if $A^*A = AA^*$. For example,
$$\begin{bmatrix} i & 0 \\ 0 & 2 \end{bmatrix}, \quad \begin{bmatrix} i & 2 \\ 2 & i \end{bmatrix}.$$
- A is called a unitary matrix if $A^*A = I = AA^*$. For example,
$$\begin{bmatrix} i & 0 \\ 0 & 1 \end{bmatrix}, \quad \frac{1}{2} \begin{bmatrix} i & 1 \\ 1 & i \end{bmatrix}.$$

Let $A \in \mathcal{M}_n(\mathbb{C})$, the set of all $n \times n$ complex matrices.

- $\sigma(A) := \{\lambda_1, \dots, \lambda_n\}$ (a multi-set), where $\lambda_1, \dots, \lambda_n$ are the eigenvalues (not necessarily distinct) of A . This $\sigma(A)$ is called the spectrum of A .
- If λ is an eigenvalue of A with corresponding eigenvector \mathbf{v} , then (λ, \mathbf{v}) is called an eigenpair of A .
- If $A = (a_{ij})$, then $\text{Tr}(A) = a_{11} + \dots + a_{nn}$ is called the trace of A .
- A is called a normal matrix if $A^*A = AA^*$. For example, $\begin{bmatrix} i & 0 \\ 0 & 2 \end{bmatrix}$, $\begin{bmatrix} i & 2 \\ 2 & i \end{bmatrix}$.
- A is called a unitary matrix if $A^*A = I = AA^*$. For example, $\begin{bmatrix} i & 0 \\ 0 & 1 \end{bmatrix}$, $\frac{1}{2} \begin{bmatrix} i & 1 \\ 1 & i \end{bmatrix}$.
- A is called an orthogonal matrix if A is real and $A^t A = I = A A^t$. For example, $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$.

Schur unitary triangularization

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real,

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n .

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = \begin{bmatrix} w_1 & \cdots & w_n \end{bmatrix}$.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis,

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Put $U = W \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Put $U = W \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$. Note: U is unitary.

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Put $U = W \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$. Note: U is unitary. Now

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Put $U = W \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$. Note: U is unitary. Now

$$U^*AU = \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} W^*AW \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$$

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Put $U = W \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$. Note: U is unitary. Now

$$U^*AU = \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} W^*AW \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$$

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Put $U = W \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$. Note: U is unitary. Now

$$\begin{aligned} U^*AU &= \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} W^*AW \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix} \\ &= \begin{bmatrix} \lambda_1 & * \\ 0 & U'^*A'U' \end{bmatrix} \end{aligned}$$

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Put $U = W \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$. Note: U is unitary. Now

$$\begin{aligned} U^*AU &= \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} W^*AW \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix} \\ &= \begin{bmatrix} \lambda_1 & * \\ 0 & U'^*A'U' \end{bmatrix} = \begin{bmatrix} \lambda_1 & * \\ 0 & T' \end{bmatrix}. \end{aligned}$$

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

Pr. Use induction on n . Case $n = 1$ is trivial. Let $n > 1$. Let (λ_1, w_1) be an eigenpair of A , $\|w_1\| = 1$. Take an orthonormal basis $\{w_1, \dots, w_n\}$ and form $W = [w_1 \ \dots \ w_n]$. Then, $W^*AW = \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix}$, $A' \in \mathcal{M}_{n-1}$.

By induction hypothesis, $\exists U'$ unitary, s.t. $T' = U'^*A'U'$ is upper triangular.

Put $U = W \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix}$. Note: U is unitary. Now

$$\begin{aligned} U^*AU &= \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} W^*AW \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & U'^* \end{bmatrix} \begin{bmatrix} \lambda_1 & * \\ 0 & A' \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & U' \end{bmatrix} \\ &= \begin{bmatrix} \lambda_1 & * \\ 0 & U'^*A'U' \end{bmatrix} = \begin{bmatrix} \lambda_1 & * \\ 0 & T' \end{bmatrix}. \end{aligned}$$

The rest follows from: 'if A, λ are real then $\exists x \neq 0$ real, s.t. $Ax = \lambda x$ '. ■

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.
Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

A

A

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.
Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

$$W^*AW$$

$$\left[\begin{array}{c|c} \lambda_1 & * \\ \hline 0 & A_1 \end{array} \right]$$

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.
Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

$$\begin{bmatrix} 1 & 0 \\ 0 & U_1^* \end{bmatrix} W^* A W \begin{bmatrix} 1 & 0 \\ 0 & U_1 \end{bmatrix}$$

$$\left[\begin{array}{cc|c} \lambda_1 & 0 & * \\ 0 & \lambda_2 & * \\ \hline 0 & 0 & A_2 \end{array} \right]$$

Th. Let $A \in \mathcal{M}_n(\mathbb{C})$. Then $\exists U$ unitary, s.t. $T = U^*AU$ is upper triangular.

Further, if A and $\sigma(A)$ are real, then U can be real orthogonal.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & U_2^* \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & U_1^* \end{bmatrix} W^* A W \begin{bmatrix} 1 & 0 \\ 0 & U_1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & U_2 \end{bmatrix}$$

$$\left[\begin{array}{ccc|c} \lambda_1 & 0 & 0 & * \\ 0 & \lambda_2 & 0 & * \\ 0 & 0 & \lambda_3 & * \\ \hline 0 & 0 & 0 & A_3 \end{array} \right]$$

Applications: Schur unitary triangularization (SUT)

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

So $\sum \lambda_i =$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} =$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) =$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) =$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) =$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i =$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} =$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T =$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU =$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** **A** we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A .

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Th[spectral Theorem]. Let A be normal ($AA^* = A^*A$).

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Th[spectral Theorem]. Let A be normal ($AA^* = A^*A$). Then $\exists U$ unitary, s.t. $U^*AU = D$ is diagonal.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Th[spectral Theorem]. Let A be normal ($AA^* = A^*A$). Then $\exists U$ unitary, s.t. $U^*AU = D$ is diagonal. Further, $\text{DIAG } D = \sigma(A)$.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Th[spectral Theorem]. Let A be normal ($AA^* = A^*A$). Then $\exists U$ unitary, s.t. $U^*AU = D$ is diagonal. Further, **DIAG** $D = \sigma(A)$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Th[spectral Theorem]. Let A be normal ($AA^* = A^*A$). Then $\exists U$ unitary, s.t. $U^*AU = D$ is diagonal. Further, **DIAG** $D = \sigma(A)$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^*A = AA^*$, we get $T^*T = TT^*$.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Th[spectral Theorem]. Let A be normal ($AA^* = A^*A$). Then $\exists U$ unitary, s.t. $U^*AU = D$ is diagonal. Further, **DIAG** $D = \sigma(A)$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^*A = AA^*$, we get $T^*T = TT^*$. Note: $\sum |t_{1i}|^2 = (TT^*)_{11} = (T^*T)_{11} = |t_{11}|^2$.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. Υ

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Th[spectral Theorem]. Let A be normal ($AA^* = A^*A$). Then $\exists U$ unitary, s.t. $U^*AU = D$ is diagonal. Further, **DIAG** $D = \sigma(A)$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^*A = AA^*$, we get $T^*T = TT^*$. Note: $\sum |t_{1i}|^2 = (TT^*)_{11} = (T^*T)_{11} = |t_{11}|^2$. So $t_{12} = \dots = t_{1n} = 0$.

Remark. In SUT, as $U^*AU = T$, we have

$$\{\lambda_1, \dots, \lambda_n\} = \sigma(A) = \sigma(T) = \{t_{11}, \dots, t_{nn}\}.$$

Further, we can get the λ_i s in the diagonal of T in any prescribed order. **Y**

Cor. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. Then $\text{DET } A = \prod \lambda_i$ and $\text{TR } A = \sum \lambda_i$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. So $\sigma(T) = \sigma(A)$.

$$\text{So } \sum \lambda_i = \sum t_{ii} = \text{TR}(T) = \text{TR}(U^*AU) = \text{TR}(AUU^*) = \text{TR}(A).$$

$$\text{And } \prod \lambda_i = \prod t_{ii} = \text{DET } T = \text{DET } U^*AU = \text{DET } A. \quad \blacksquare$$

- By **DIAG** A we denote the diagonal $\{a_{11}, \dots, a_{nn}\}$ of A . By **DIAG** v we denote the diagonal matrix A with $a_{11} = v_1, \dots, a_{nn} = v_n$.

Th[spectral Theorem]. Let A be normal ($AA^* = A^*A$). Then $\exists U$ unitary, s.t. $U^*AU = D$ is diagonal. Further, **DIAG** $D = \sigma(A)$.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^*A = AA^*$, we get $T^*T = TT^*$. Note: $\sum |t_{1i}|^2 = (TT^*)_{11} = (T^*T)_{11} = |t_{11}|^2$. So $t_{12} = \dots = t_{1n} = 0$. Repeating the process, T is diagonal. ■

Applications: Schur unitary triangularization

Cor. Let A be Hermitian.

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular.

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$.

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & \textcolor{red}{0} & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & \color{red}{0} & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & \textcolor{red}{0} & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & \color{red}{0} & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} = \begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & \textcolor{red}{0} & * \\ 0 & 0 & 0 & * \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & \textcolor{red}{0} & * \\ 0 & 0 & 0 & * \end{bmatrix} = \begin{bmatrix} 0 & & & \\ 0 & & & \\ 0 & & & \\ 0 & & & \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & \textcolor{red}{0} & * \\ 0 & 0 & 0 & * \end{bmatrix} = \begin{bmatrix} 0 & 0 & & \\ 0 & 0 & & \\ 0 & 0 & & \\ 0 & 0 & & \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & \textcolor{red}{0} & * \\ 0 & 0 & 0 & * \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & \textcolor{red}{0} & * \\ 0 & 0 & 0 & * \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• Multiply

$$\begin{bmatrix} 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & \textcolor{red}{0} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$.

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular; $t_{ii} = \lambda_i$ for each i .

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular; $t_{ii} = \lambda_i$ for each i . Let $p(x)$ be the characteristic polynomial of A .

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular; $t_{ii} = \lambda_i$ for each i . Let $p(x)$ be the characteristic polynomial of A . So $p(x) = \prod(x - \lambda_i)$.

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular; $t_{ii} = \lambda_i$ for each i . Let $p(x)$ be the characteristic polynomial of A . So $p(x) = \prod (x - \lambda_i)$. So

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular; $t_{ii} = \lambda_i$ for each i . Let $p(x)$ be the characteristic polynomial of A .

So $p(x) = \prod (x - \lambda_i)$. So

$$p(A) = \prod (A - \lambda_i I)$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular; $t_{ii} = \lambda_i$ for each i . Let $p(x)$ be the characteristic polynomial of A .

So $p(x) = \prod (x - \lambda_i)$. So

$$p(A) = \prod (A - \lambda_i I) = \prod (UTU^* - \lambda_i UIU^*)$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular; $t_{ii} = \lambda_i$ for each i . Let $p(x)$ be the characteristic polynomial of A .

So $p(x) = \prod (x - \lambda_i)$. So

$$\begin{aligned} p(A) &= \prod (A - \lambda_i I) = \prod (UTU^* - \lambda_i UIU^*) \\ &= U \left[(T - \lambda_1 I)(T - \lambda_2 I) \cdots (T - \lambda_n I) \right] U^* \end{aligned}$$

Cor. Let A be Hermitian. Then $\exists U$ unitary, s.t. $U^*AU = D$ is real diagonal.

Further, if A is real, then U can be chosen real orthogonal.

Pr. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular. As $A^* = A$, we get $T^* = T$. So T is real diagonal. ■

• So

$$\begin{bmatrix} 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \\ 0 & 0 & 0 & * \end{bmatrix} \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0.$$

Cor[Cayley-Hamilton]. Every matrix satisfies its characteristic equation.

Pr. Let $\sigma(A) = \{\lambda_1, \dots, \lambda_n\}$. By SUT, $\exists U$ unitary, s.t. $U^*AU = T$ is upper triangular; $t_{ii} = \lambda_i$ for each i . Let $p(x)$ be the characteristic polynomial of A . So $p(x) = \prod(x - \lambda_i)$. So

$$\begin{aligned} p(A) &= \prod(A - \lambda_i I) = \prod(UTU^* - \lambda_i UIU^*) \\ &= U \left[(T - \lambda_1 I)(T - \lambda_2 I) \cdots (T - \lambda_n I) \right] U^* = U 0 U^* = 0. \end{aligned}$$