2019117

[3]

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 1 of 18

 $\operatorname{Roll\ No.:}\ 170108024$

Name: Mukul Ranjan

Mukul Ranjan

Important Instructions: Read carefully!

- Write your answers in the space provided against each question. If required, you may continue your solutions only on pages 16, 17, 18. Answers written elsewhere will not be evaluated.
- · No extra sheets will be provided for writing solutions.
- You can use the supplement(s) for rough work. These sheets will not be evaluated.
- No marks will be awarded if you write only the final answer.
- 1. Check whether the following systems of linear equations are equivalent or not:

Space only for Q. 1

AWD :

Augumented Matrix of System-I

Do not write below this line

| PageID: 170108024-01

 ${\bf MarkingScheme:~1+1+1}$

PageMark: **1+1+1**

 \mathbf{c}

PageID: 170108024-01

MarkingScheme: 1+1+1

PageMark: **1+1+1**

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 2 of 18

 $\operatorname{Roll\ No.:}\ 170108024$

Name: Mukul Ranjan

Space only for Q. 1

Now writing the argumented modpin of System-I

$$\begin{bmatrix} 47 & 1 & 40 \\ 13 & -1 & 20 \\ 123 & 3 \\ 041 & 17 \end{bmatrix} \xrightarrow{R_3 \leftarrow R_3 - R_2} \begin{bmatrix} 47 & 146 \\ 13 & -120 \\ 0 & -14 & 17 \end{bmatrix}$$

embine on Page

PageID: 170108024-02

MarkingScheme:

PageMark:

PageID: 170108024-02

MarkingScheme:

PageMark:

Missing a negative sign here.

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 16 of 18

Roll No.: 170108024

Name: Mukul Ranjan

Extra space for answers, NOT rough work. Doing rough work here or not mentioning the Q. No. clearly will invite NEGATIVE marking.

Q.1 continue

Since the both system has some solution sit hence they are aquivalent.

Continue Quertion-2(a)

$$T \begin{bmatrix} n_1 \\ n_2 \end{bmatrix} = \begin{bmatrix} 2 m_1 - m_3 \\ -m_1 + m_2 + m_3 \\ -m_1 + m_2 + m_3 \end{bmatrix}$$

$$2 m_2 + 2 m_3$$

DO NOT WRITE BELOW THIS LINE

PageID: 170108024-16

MarkingScheme:

PageMark:

PageID: 170108024-16

MarkingScheme:

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 3 of 18

 $\mathrm{Roll\ No.:}\ 170108024$

Name: Mukul Ranjan

2. Let $T: \mathbb{R}^3 \to \mathbb{R}^4$ be a linear transformation such that

$$T\left(\begin{bmatrix}1\\1\\1\end{bmatrix}\right) = \begin{bmatrix}1\\0\\1\\0\end{bmatrix}, \qquad T\left(\begin{bmatrix}1\\1\\0\end{bmatrix}\right) = \begin{bmatrix}1\\2\\1\\2\end{bmatrix}, \qquad T\left(\begin{bmatrix}0\\1\\1\end{bmatrix}\right) = \begin{bmatrix}0\\1\\0\\2\end{bmatrix}$$

(a) Determine a matrix A such that T(x) = Ax for all $x \in \mathbb{R}^3$.

[3

Space only for Q. 2 (a)

Let
$$n = \begin{bmatrix} n_1 \\ n_2 \\ n_3 \end{bmatrix} \in \mathbb{R}^3$$

$$\begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \\ m_3 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 2 & 2 \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \\ m_3 \end{bmatrix}$$

solving this equ equising argumented neutrix

$$\begin{bmatrix}
1 & 1 & 0 & 1 & 1 & 0 \\
1 & 1 & 1 & 0 & 21 \\
1 & 0 & 1 & 1 & 0 \\
0 & 0 & 0 & 0 & 22
\end{bmatrix}
\xrightarrow{R_2 \leftarrow R_2 - R_1}
\begin{bmatrix}
1 & 1 & 0 & | & 1 & 0 \\
0 & 0 & 1 & | & -1 & 1 \\
1 & 0 & 1 & | & 1 & 0 \\
0 & 0 & 0 & 0 & 22
\end{bmatrix}$$

Thus, we have
$$\begin{bmatrix}
n_1 \\
n_3
\end{bmatrix} = \begin{bmatrix}
-n_1 + n_2 + n_3 \\
2n_1 - n_3 \\
-n_1 + n_2 + n_3 \\
2n_2 + 2n_3
\end{bmatrix}$$

DO NOT WRITE BELOW THIS LINE

continu of Page 16

| | PageID: 170108024-03

MarkingScheme: 1+1+1

PageMark: 0+0+0

PageID: 170108024-03

 ${\bf Marking Scheme: \ 1+1+1}$

PageMark: 0+0+0

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 4 of 18

 $\operatorname{Roll\ No.:}\ 170108024$

Name: Mukul Ranjan

(b) Justify whether such a matrix A as above is unique.

[2_]

Space only for Q. 2 (b)

Since, the Sept part of argumented matrix of O is in RREF which is unique hence matrix A must be unique.

PageID: 170108024-04

MarkingScheme: 2

PageMark: $\mathbf{0}$

PageID: 170108024-04

MarkingScheme: 2

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 16 of 18

Roll No.: 170108024

Name: Mukul Ranjan

Extra space for answers, NOT rough work. Doing rough work here or not mentioning the Q. No. clearly will invite NEGATIVE marking.

Q.1 continue

Since the both system has some solution sit hence they are aquivalent.

Continue Quertion-2(a)

$$T \begin{bmatrix} n_1 \\ n_2 \end{bmatrix} = \begin{bmatrix} 2 m_1 - m_3 \\ -m_1 + m_2 + m_3 \\ -m_1 + m_2 + m_3 \end{bmatrix}$$

$$2 m_2 + 2 m_3$$

DO NOT WRITE BELOW THIS LINE

PageID: 170108024-16

MarkingScheme:

PageMark:

PageID: 170108024-16

MarkingScheme:

 $\#08 \ of \ 38$ /MA 101/Midsem/2017-18 **2203**/20-Sep-2017/p. 5 of 18

Roll No.: 170108024

Name: Mukul Ranjan

3. State with proper justification, whether each of the following statements is **True** or **False**.

(a) If A be a nonzero matrix such that $A^{31} = 0$ then all eigenvalues of A are equal to 0 and A is not diagonalizable.

Space only for Q. 3 (a)

A + 0

Photo

AM = AX

we know that Rigen value of An is 2n.

hence, ASINC = 231 oc

AM = AM

A·AN= AAM= A(AM)

=> A2 N = A(AN) = A2N :. A31 N = A31 M

A31 = 0

Agimeo = ygimeo

N+0

DO NOT WRITE BELOW THIS LINE

231=0 : A= 0

thus all the eight value of A = 0 and hence A is not diagonisable.

stadement is time. Hence,

PageID: 170108024-05

MarkingScheme: 1

PageMark: 0

PageID: 170108024-05

MarkingScheme: 1

PageMark: $\mathbf{0}$

Wrong argument. A matrix with all eigenvalue zero may be diagonaliz#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 6 of 18

Roll No.: 170108024

Name: Mukul Ranjan

(b) Any square matrix is similar to its Reduced Row Echelon Form.

[1]

Space only for Q. 3 (b)

any square matrix A. be the R.R.E.F of

R = RR.BFLA)

we know know you swo matrix, so solve similar

det (A) = det (B)

home same steeped io) skee should

we know that det (A) & det R But

thus, The given any square metris is not similar to lits R.R.E.T.

set of elgenvalues same xinton house similar Row transformation does Holl me Jenous eigen values hence eigen values of R.R.1 are different thus, they need not p reserve are not similar.

DO NOT WRITE BELOW THIS LINE

PageID: $170108024-06$	PageID: $170108024-06$
MarkingScheme: 1	MarkingScheme: 1
PageMark: 0	PageMark: 0

To show a statement is false you are Why $det(A) \neq det(R)$? Give an examrequired to give a counter example ple. with proper justification.

 $\#08 \ of \ 38$ /MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 7 of 18

Roll No.: 170108024

Name: Mukul Ranjan

det A = ± 1

4. (a) Let A be a real $n \times n$ matrix such that A is similar to A^{2017} . Determine all the possible values for the determinant of A.

we know the determinant of similar matrices an equal; as AP=PB dut (AP) = det(PB) det A. alet P = det P. det A det A = det B

Now, det A = det (A 2017) we know that det (An) = (det A) =) det A = (det A) 2017 =) (det A)2017_ detA = 0 det A [det A) 2016 [] = 0 det A = 0 on blut A) 2016 = 1

thus, det A = 0, 00 1, -1

DO NOT WRITE BELOW THIS LINE

PageID: 170108024-07

MarkingScheme: 1

PageMark: 1

PageID: 170108024-07

MarkingScheme: 1

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 8 of 18

Roll No.: 170108024

Name: Mukul Ranjan

(b) Let v_1 and v_2 be two nonzero eigenvectors corresponding to distinct eigenvalues λ_1 and λ_2 of a 5 × 5 matrix A. Show that $v_1 - v_2$ can not be an eignvector of A.

[2]

Space only for Q. 4

eigen values of the eigenvector v, and vr of a vector

we know that, $A V_1 = A_1 V_1 - \Theta$ and $A V_2 = A_2 V_2 - \Theta$

Subtracting @ and @

AU1-AU2 = A1U1-A2U2

From agn (11) we to can see that

Vi-Uz will be the eight walke of A vector of

A iff Ai=Az lent Ai # Az

hence (Ui-Vz) can not be an eigenvector of

PageID: 170108024-08

 ${\bf MarkingScheme:}\ \, {\bf 1+1}$

PageMark: 0+0

PageID: 170108024-08

 ${\it MarkingScheme:}\ {\bf 1+1}$

PageMark: 0+0

need proper justification

- Incorrect justification.
- Need to justify why $v_1 v_2$ will be an eigen vector iff $\lambda_1 = \lambda_2$.

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 9 of 18

Roll No.: 170108024 Name: Mukul Ranjan

5. (a) Let A be a $p \times n$ matrix and B be a $q \times n$ matrix such that $\operatorname{rank}(A) + \operatorname{rank}(B) < n$. Show that there exists a nonzero vector $x \in \mathbb{R}^n$ such that Ax = 0 and Bx = 0.

rank(A) < n-rank(B)

rank(A) + nullity of A)=1

rank(A) < nullity (B)

rank(B) + nullity(B)

rank(B) + nullity(B) Since, rank A + rank B & M Row (A) C mill (B)

> thus dimenul (B) = 0 Some vector on EIRM such that

> > BN = 0

Similarly

rank B < n- rank (A) namber a mulliby A Row (B) C null (A) : dim/null (A) + 0 here there must esist some Such short AN=0

PageID: 170108024-09

MarkingScheme: 1+1

PageMark: 0+0

PageID: 170108024-09

MarkingScheme: 1+1

PageMark: 0+0

Why common solution exixts

your aim is to show that vector x is comman, such that Ax=0, Bx=0. in your case vector may be different.

DO NOT WRITE BELOW THIS LINE

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 10 of 18

 $\operatorname{Roll\ No.:}\ 170108024$

Name: Mukul Ranjan

(b) Let v_1, \ldots, v_k be vectors in \mathbb{R}^n and A be an invertible $n \times n$ matrix. Then show that $\dim (\operatorname{Span}(\{v_1, \ldots, v_k\})) = \dim (\operatorname{Span}(\{Av_1, \ldots, Av_k\}))$

Space only for Q. 5 (b)

Colla) must loe span en as a is non maluis.

thus, dim (sol A) = n

Span (AVI -- AVN) = XAVI + d2 AV2+ - - 1 dk AVR

since A is invertible matur's number of elements wester elements in the spen & y, -- , vn y nust be equal to member of elements in the span Sav, Avr -- Avr ? as any of the span Sav, Avr -- Avr ? as any of the mow of A is not zero, hence it does not appert on nullipy any element of span (v1, v2, -v appert on nullipy any element of span (v1, v2, -v

:. dim (span (su, - - un 4)) = dim (span (spa

| PageID: 170108024-10

 ${\bf MarkingScheme:}\ {\bf 1+1}$

PageMark: 0+0

PageID: **170108024-10**

MarkingScheme: 1+1

PageMark: 0+0

wrong INCORRECT

Do not write below this line

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 11 of 18

Roll No.: 170108024 Name: Mukul Ranjan

6. Let A be an $n \times n$ matrix, where $n \geq 2$.

(a) Show that $\det(\operatorname{adj}(A)) = \det(A)^{n-1}$.

[2]

Space only for Q. 6 (a)

Case I

when det A #0

we know that

Jerking dolerminant both side -

det A. alet (Ady A) = (det A) M

det A det B det A det B det

det (Ady'A) = (det(A))^n-1

det (Ady'A) = det(A)^n-1

Muoved Tax; (det A) = detIAM

Twe have, det(AB)=

Can I det A = 0

set assume that Ady

then we have

to Priore that, det (Ady'A) = 0

It's assumblet (Ardy'A) ‡ 0 thus, Ady'A is inventible.

(Ady'A). A = (dy/A) In = 0

continue on Page 17

S) Adja

DO NOT WRITE BELOW THIS LINE

PageID: **170108024-11**

 ${\bf MarkingScheme:}\ {\bf 1+1}$

PageMark: 1+1

PageID: **170108024-11**

 ${\it MarkingScheme:}\ {\bf 1+1}$

PageMark: 1+1

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 12 of 18

 $\operatorname{Roll\ No.:}\ 170108024$

Name: Mukul Ranjan

(b) If A has rank n-1 then show that adj(A) has rank 1. (You may use the rank-nullity theorem.)

[3]

Space only for Q. 6 (b)

we have,

Rank (A) & milliby (A) = n

Rank'A = 80 m- millity (A)

Rank (A)= n-1 : mullity (A)=1

matrix A contain 1 zero mon. thus

 $A = \begin{bmatrix} a_{11} & a_{12} & - & - & a_{1n} \\ a_{21} & a_{22} & - & - & a_{2n} \\ a_{n-1} & a_{n-2n} \\ - & a_{2n} \end{bmatrix}$

Since · Sulorde copulor hence. det Rouk (Adj'A) = n

DO NOT WRITE BELOW THIS LINE

PageID: 170108024-12

MarkingScheme: 1+1+1

PageMark: 0+0+0

PageID: 170108024-12

MarkingScheme: 1+1+1

PageMark: 0+0+0

Incorrect reasoning. rank(A) = n - 11 implies that REF of A has exactly one zero row, but A need not have a zero row. See grading scheme for an example.

You had to prove that rank(adj(A) =

Incorrect solution.

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 17 of 18

Roll No.: 170108024 Name: Mukul Ranjan

Extra space for answers, NOT rough work. Doing rough work here or not mentioning the Q. No. clearly will invite NEGATIVE marking.

Quest. 6. (a)

$$(Adj'A)\cdot A = 0$$

$$(Adj'A)^{-1}(Adj'A)\cdot A = 0$$

$$A = 0$$

heme.

Question-8:

$$W = \begin{cases} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} & \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} \end{cases}$$

$$\omega = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Calculating
$$W^{\perp}$$
:

 $v_1 = m_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
 $v_2 = m_2 - pmoj_2 x_2$

DO NOT WRITE BELOW THIS LINE

PageID: 170108024-17

MarkingScheme:

PageMark:

 ${\it PageID:}~ 170108024\text{--}17$

MarkingScheme:

#08 of 38/MA 101/Midsem/2017-182203/20-Sep-2017/p. 13 of 18

Roll No.: 170108024 Name: Mukul Ranjan

7. Let A be an $n \times n$ $(n \ge 3)$ matrix with entries a_{ij} as follows:

$$a_{ij} = \begin{cases} 3 & \text{if } i = j \\ 2 & \text{if } i \neq j \end{cases}$$

- (a) Without computing the characteristic polynomial, determine all the eigenvalues of A, and a basis for each eigenspace of A. (You need to justify why there is no eigenvalue other than those you have found.)
- (b) Find an invertible matrix P and a diagonal matrix D such that $P^{-1}AP = D$.

(Answering it only for special cases like n=3 or 4 will be awarded zero marks.)

Space only for Q. 7

DO NOT WRITE BELOW THIS LINE

A =
$$\begin{bmatrix} 3 & 2 & 2 \\ 2 & 3 & 2 & 1 \\ 2 & 2 & 3 & 1 \\ 2 & 2 & -1 & -1 & 3 \end{bmatrix}$$
 Since A is symmetric matrix

AT = A

[4]

[1]

PageID: 170108024-13

MarkingScheme: 1+1+1+1+1

PageMark: **0+0+0+0+0**

PageID: 170108024-13

MarkingScheme: 1+1+1+1+1

PageMark: 0+0+0+0+0

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 14 of 18

Roll No.: 170108024 Name: Mukul Ranjan

Space only for Q. 7

PageID: 170108024-14	PageID: 170108024-14	
MarkingScheme:	MarkingScheme:	
PageMark:	PageMark:	

BlankPage

BlankPage

DO NOT WRITE BELOW THIS LINE

#08 of 38/MA 101/Midsem/2017-18**2203**/20-Sep-2017/p. 15 of 18

Roll No.: 170108024 Name: Mukul Ranjan

8. Let

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 & 2 \\ -1 & 1 & 3 & 7 & 0 \\ 0 & 2 & 4 & 8 & 2 \\ 1 & -3 & -7 & -15 & -2 \end{bmatrix}, \quad W = \operatorname{col}(A) \quad \text{and} \quad w = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

Determine the orthogonal projection of w onto W^{\perp} .

[3]

Space only for Q. 8

$$-C_1 + 2C_2 = C_3$$
 $-3C_1 + 44C_2 = C_4$
 $C_1 + C_2 = C_4$

linearly independent.

: Col(A) = Span
$$\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} \right\}$$

Since relation between colours does not in Row fronsponnation thus, col 2 and in A is also independent. Cal 2 DO NOT WRITE BELOW THIS LINE Continue of Page - 17

PageID: 170108024-15

MarkingScheme: 1+1+1

PageMark: **1+1+0**

PageID: 170108024-15

MarkingScheme: 1+1+1

PageMark: **1+1+0**

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 17 of 18

Roll No.: 170108024 Name: Mukul Ranjan

Extra space for answers, NOT rough work. Doing rough work here or not mentioning the Q. No. clearly will invite NEGATIVE marking.

Quest. 6. (a)

$$(Adj'A)\cdot A = 0$$

$$(Adj'A)^{-1}(Adj'A)\cdot A = 0$$

$$A = 0$$

heme.

Question-8:

$$W = \begin{cases} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} & \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} \end{cases}$$

$$\omega = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Calculating
$$W^{\perp}$$
:

 $v_1 = m_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
 $v_2 = m_2 - pmoj_2 x_2$

DO NOT WRITE BELOW THIS LINE

PageID: 170108024-17

MarkingScheme:

PageMark:

 ${\it PageID:}~ 170108024\text{--}17$

MarkingScheme:

#08 of 38/MA 101/Midsem/2017-18 2203/20-Sep-2017/p. 18 of 18

 $\operatorname{Roll\ No.:}\ 170108024$

Name: Mukul Ranjan

Extra space for answers, NOT rough work. Doing rough work here or not mentioning the Q. No. clearly will invite $\overrightarrow{NEGATIVE}$ marking.

DO NOT WRITE BELOW THIS LINE

PageID: 170108024-18	PageID: 170108024-18		
MarkingScheme:	MarkingScheme:		
PageMark:	PageMark:		

The basis obtained by you is the basis of tained by you is not for for W not for W^{\perp} . You did mistake. W^{\perp} . This is for W.

MA101:2017-18:Odd:Midsem

 $\underline{\mathbf{Mathematics}\text{-}\mathbf{I}}$

Graded Answer Script of 170108024:Mukul Ranjan;T7;2203;8;38

Question	YourScore	ClassAverage	ClassMax	ClassMin
1	3	1.02	3	0
2	0	2.49	5	0
3	0	.32	2	0
4	1	1.32	3	0
5	0	.27	4	0
6	2	1.02	5	0
7	0	.29	5	0
8	2	.50	3	0
TOTAL	8	7.27	26	0