

Basic Definitions, Existence and Uniqueness Results for First-Order IVP

Department of Mathematics
IIT Guwahati

Texts/References:

S. L. Ross, Differential Equations, John Wiley & Son Inc, 2004.

W. E. Boyce and R. C. DiPrima, Elementary Differential Equations and Boundary Value Problems, John Wiley & Son, 2001.

E. A. Coddington, An Introduction to Ordinary Differential Equations, Prentice Hall India, 1995.

E. L. Ince, Ordinary Differential Equations, Dover Publications, 1958.

Dennis G. Zill and Michael R. Cullen, Differential equations with boundary value problems

Definition: An equation involving derivatives of one or more dependent variables with respect to one or more independent variables is said to be a **differential equation**(DE).

Definition: A DE involving ordinary derivatives of one or more dependent variables w.r.t a single independent variable is called an **ordinary differential equation**(ODE).

A general form of the n th order ODE:

$$F(x, y(x), y'(x), y''(x), \dots, y^{(n)}(x)) = 0, \quad (1)$$

where $y'(x) = \frac{dy}{dx}$, $y''(x) = \frac{d^2y}{dx^2}$, \dots , $y^{(n)}(x) = \frac{d^ny}{dx^n}$.

- The **order** of a DE is the order of the highest derivative that occurs in the equation.
- The **degree** of a DE is the power of the highest order derivative occurring in the differential equation.
- Eq. (1) is **linear** if F is linear in $y, y', y'', \dots, y^{(n)}$, with coefficients depending on the independent variable x . Eq. (1) is called **nonlinear** if it is not linear.

Examples:

- $y''(x) + 3y'(x) + xy(x) = 0$
(second-order, first-degree, linear)
- $y''(x) + 3y(x)y'(x) + xy(x) = 0$
(second-order, first-degree, nonlinear)
- $(y''(x))^2 + 3y'(x) + xy^2(x) = 0$
(second-order, second-degree, nonlinear)

Definition: A DE involving partial derivatives of one or more dependent variables w.r.t more than one independent variable is called a **partial differential equation**(PDE).

A PDE for a function $u(x_1, x_2, \dots, x_n)$ ($n \geq 2$) is a relation of the form

$$F(x_1, x_2, \dots, x_n, u, u_{x_1}, u_{x_2}, \dots, u_{x_1 x_1}, u_{x_1 x_2}, \dots) = 0, \quad (2)$$

where F is a given function of the **independent variables** x_1, x_2, \dots, x_n , and of the unknown function u and of a finite number of its partial derivatives.

Examples:

- $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 0$ (first-order equation)
- $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ (second-order equation)

We shall consider only ODE.

Definition: A function $\phi(x) \in C^n((a, b))$ that satisfies

$$F(x, \phi(x), \phi'(x), \phi''(x), \dots, \phi^n(x)) = 0, \quad x \in (a, b)$$

is called an **explicit solution** to the equation on (a, b) .

Example: $\phi(x) = x^2 - x^{-1}$ is an **explicit solution** to

$$y''(x) - 2\frac{y}{x^2} = 0.$$

Note that $\phi(x)$ is an explicit solution on $(-\infty, 0)$ and also on $(0, \infty)$.

Definition: (Initial Value Problem)

Find a solution $y(x) \in C^n((a, b))$ that satisfies

$$F(x, y, y'(x), \dots, y^{(n)}(x)) = 0, \quad x \in (a, b)$$

and the n initial conditions(IC)

$$y(x_0) = y_0, \quad y'(x_0) = y_1, \dots, y^{(n-1)}(x_0) = y_{n-1},$$

where $x_0 \in (a, b)$ and y_0, y_1, \dots, y_{n-1} are given constants.

First-order IVP: $F(x, y, y'(x)) = 0, \quad y(x_0) = y_0.$

Second-order IVP: $F(x, y, y'(x), y''(x)) = 0,$
 $y(x_0) = y_0, \quad y'(x_0) = y_1.$

Example: The function $\phi(x) = \sin x - \cos x$ is a solution to IVP: $y''(x) + y(x) = 0, \quad y(0) = -1, \quad y'(0) = 1.$ on $\mathbb{R}.$

Consider the following IVPs:

$$|y'| + 2|y| = 0, \quad y(0) = 1 \text{ (no solution)}.$$

$$y'(x) = x, \quad y(0) = 1 \text{ (a unique solution } y = \frac{1}{2}x^2 + 1 \text{)}.$$

$$xy'(x) = y - 1, \quad y(0) = 1 \text{ (many solutions } y = 1 + cx \text{)}.$$

Observation:

Thus, an IVP

$$F(x, y, y') = 0, \quad y(x_0) = y_0$$

may have none, precisely one, or more than one solution.

Well-posed IVP

An IVP is said to be **well-posed** if

- it has a solution,
- the solution is unique and,
- the solution is continuously depends on the initial data y_0 and f .

Theorem(Peano's Theorem):

Let $\mathcal{R} : |x - x_0| \leq a, |y - y_0| \leq b$ be a rectangle. If $f \in C(\mathcal{R})$ then the IVP

$$y'(x) = f(x, y), \quad y(x_0) = y_0$$

has **at least one solution** $y(x)$. This solution is defined for all x in the interval $|x - x_0| \leq h$, where

$$h = \min\left\{a, \frac{b}{K}\right\}, \quad K = \max_{(x,y) \in \mathcal{R}} |f(x, y)|.$$

Example: Let $\mathcal{R} : |x - 0| \leq 3, |y - 0| \leq 3$ be a rectangle. Let $f(x, y) = xy$. Then $f \in C(\mathcal{R})$. Then the IVP

$$y'(x) = f(x, y), \quad y(0) = 0$$

has at least one solution $y(x)$. This solution is defined for all x in the interval $|x - 0| \leq h$, where

$$h = \min\left\{3, \frac{3}{K}\right\}, \quad K = \max_{(x,y) \in \mathcal{R}} |xy|.$$

Theorem(Picard's Theorem):

Let $f \in C(\mathcal{R})$ and satisfy the **Lipschitz condition** with respect to y in \mathcal{R} , i.e., there exists a number L such that

$$|f(x, y_2) - f(x, y_1)| \leq L|y_2 - y_1| \quad \forall (x, y_1), (x, y_2) \in \mathcal{R}.$$

Then, the IVP

$$y'(x) = f(x, y), \quad y(x_0) = y_0$$

has a unique solution $y(x)$. This solution is defined for all x in the interval $|x - x_0| \leq h$, where

$$h = \min\left\{a, \frac{b}{K}\right\}, \quad K = \max_{(x,y) \in \mathcal{R}} |f(x, y)|$$

Example: Consider the IVP:

$$y'(x) = |y|, \quad y(1) = 1.$$

$f(x, y) = |y|$ is continuous and satisfies Lipschitz condition w.r.t y in every domain \mathcal{R} of the xy -plane. The point $(1, 1)$ certainly lies in some such domain \mathcal{R} . The IVP has a unique solution ϕ defined on some $|x - 1| \leq h$ about $x_0 = 1$.

Corollary to Picard's Theorem:

Let $f, \frac{\partial f}{\partial y} \in C(\mathcal{R})$. Then the IVP

$$y'(x) = f(x, y), \quad y(x_0) = y_0$$

has a **unique solution** $y(x)$. This solution is defined for all x in the interval $|x - x_0| \leq h$, where

$$h = \min\left\{a, \frac{b}{K}\right\}, \quad K = \max_{(x,y) \in \mathcal{R}} |f(x, y)|.$$