

# Basic Definitions, Existence and Uniqueness Results for First-Order IVP

Department of Mathematics  
IIT Guwahati

## Texts/References:

S. L. Ross, Differential Equations, John Wiley & Son Inc, 2004.

W. E. Boyce and R. C. DiPrima, Elementary Differential Equations and Boundary Value Problems, John Wiley & Son, 2001.

E. A. Coddington, An Introduction to Ordinary Differential Equations, Prentice Hall India, 1995.

E. L. Ince, Ordinary Differential Equations, Dover Publications, 1958.

Dennis G. Zill and Michael R. Cullen, Differential equations with boundary value problems

**Example:** Let  $\mathcal{R} : |x| \leq 5, |y| \leq 3$  be the rectangle.  
Consider the IVP

$$y' = 1 + y^2, \quad y(0) = 0$$

over  $\mathcal{R}$ .

Here,  $a = 5$ ,  $b = 3$ . Then

$$\max_{(x,y) \in \mathcal{R}} |f(x,y)| = \max_{(x,y) \in \mathcal{R}} |1 + y^2| = 10 (= K),$$

$$\max_{(x,y) \in \mathcal{R}} \left| \frac{\partial f}{\partial y} \right| = \max_{(x,y) \in \mathcal{R}} 2|y| = 6 (= L).$$

$$h = \min\left\{a, \frac{b}{K}\right\} = \min\left\{5, \frac{3}{10}\right\} = 0.3 < 5.$$

Note that the solution of the IVP is  $y = \tan x$ . This solution is valid in the interval  $|x| \leq 0.3$  instead of the entire interval  $|x| \leq 5$ .

**Example(Non-uniqueness):** Consider the IVP:

$$y' = 3 y^{\frac{2}{3}} \text{ for } x \in \mathbb{R}, \quad y(0) = 0.$$

For each real number  $c \geq 0$ , let

$$y_c(x) = \begin{cases} 0 & \text{if } 0 \leq x < c, \\ (x - c)^3 & \text{if } c \leq x < \infty, \end{cases}.$$

It is easy to verify that  $y_c(x)$  is a solution to the IVP. Therefore, this IVP has **infinitely many solutions**.

# The Method of Successive Approximations

Consider the IVP

$$y'(x) = f(x, y), \quad y(x_0) = y_0. \quad (1)$$

**Key Idea:** Replacing the IVP (1) by an the equivalent integral equation

$$y(x) = y_0 + \int_{x_0}^x f(t, y(t)) dt. \quad (2)$$

Note that (1) and (2) are equivalent.

A first (rough) approximation to a solution is given by  $y_0(x) = y_0$ . A second approximation  $y_1(x)$  is obtained as follows:

$$y_1(x) = y_0 + \int_{x_0}^x f(t, y_0(t)) dt.$$

The next step is to use  $y_1(x)$  to generate another approximation  $y_2(x)$  in the same way:

$$y_2(x) = y_0 + \int_{x_0}^x f(t, y_1(t)) dt.$$

At the  $n$ th step, we have

$$y_n(x) = y_0 + \int_{x_0}^x f(t, y_{n-1}(t)) dt.$$

This procedure is called **Picard's method of successive approximations**.

**Example:** Consider IVP:  $y' = y$ ,  $y(0) = 1$ .

$$y_n(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!}.$$

Note that  $y_n(x) \rightarrow e^x$  as  $n \rightarrow \infty$ .

**Theorem:** Let  $\mathcal{R} : |x - x_0| \leq a, |y - y_0| \leq b$  be a rectangle where  $a, b > 0$ . Let  $f \in C(\mathcal{R})$  and let  $|f(x, y)| \leq M$  for all  $(x, y) \in \mathcal{R}$ . Further suppose that  $f$  satisfies Lipschitz condition w.r.t  $y$  with constant  $K$  in  $\mathcal{R}$ . Then the successive approximations

$$y_0(x) \equiv y_0$$

$$y_{k+1}(x) = y_0 + \int_{x_0}^x f(t, y_k(t)) dt, \quad k = 0, 1, 2, 3, \dots$$

converge uniformly on the interval  $I : |x - x_0| \leq h$  where  $h = \min\{a, \frac{b}{M}\}$  to a solution  $y(x)$  of the IVP  $y' = f(x, y), y(x_0) = y_0$ .

**Theorem(Continuous dependence on initial data):**

Let  $f, \frac{\partial f}{\partial y} \in C(\mathcal{R})$  and  $(x_0, y_0), (x_0, y_{0m}) \in \mathcal{R}$ . Let  $\phi(x)$  be the solution of

$$y' = f(x, y), y(x_0) = y_0,$$

and let  $\phi_m(x)$  be the solution of

$$y' = f(x, y), y(x_0) = y_{0m},$$

in  $\mathcal{R}$  for  $|x - x_0| \leq h$ . Then, for  $|x - x_0| \leq h$ , we have

$$|\phi(x) - \phi_m(x)| \leq |y_0 - y_{0m}|e^{Lh},$$

where  $|\frac{\partial f}{\partial y}(x, y)| \leq L$  for all  $(x, y) \in \mathcal{R}$ .

Further, as  $y_{0m} \rightarrow y_0$ ,  $\phi_m \rightarrow \phi$  uniformly on  $[x_0 - h, x_0 + h]$ .



## Theorem(Continuous dependence on $f$ ):

Let  $f, f_m, \frac{\partial f}{\partial y}, \frac{\partial f_m}{\partial y} \in C(\mathcal{R})$ , and  $(x_0, y_0) \in \mathcal{R}$ . Let  $\phi(x)$  be the solution of

$$y' = f(x, y), \quad y(x_0) = y_0,$$

and  $\phi_m(x)$  be the solution of

$$y' = f_m(x, y), \quad y(x_0) = y_0.$$

Assume that both  $\phi(x), \phi_m(x)$  exist on  $[x_0 - h, x_0 + h]$ .

Then, for  $|x - x_0| \leq h$ , we have

$$|\phi(x) - \phi_m(x)| \leq h e^{\hat{L}h} \max_{(x,y) \in \mathcal{R}} \left\{ |f(x, y) - f_m(x, y)| \right\},$$

$\hat{L} = \min\{L, L_m\}$ ,  $|\frac{\partial f}{\partial y}(x, y)| \leq L, |\frac{\partial f_m}{\partial y}(x, y)| \leq L_m \forall (x, y) \in \mathcal{R}$ . Further, as  $f_m \rightarrow f, \phi_m \rightarrow \phi$  uniformly on  $[x_0 - h, x_0 + h]$ .

## Separable Equations

**Definition:** A first-order equation  $y'(x) = f(x, y)$  is separable if it can be written in the form

$$\frac{dy}{dx} = g(x)p(y)$$

**Method for solving separable equations:** To solve the equation

$$\frac{dy}{dx} = g(x)p(y),$$

we write it as  $h(y)dy = g(x)dx$ , where  $h(y) := \frac{1}{p(y)}$ .

Integrating both sides

$$\int h(y)dy = \int g(x)dx \implies H(y) = G(x) + C,$$

which gives an implicit solution to the differential equation.

**Formal justification of method:** Writing the equation in the form

$$h(y) \frac{dy}{dx} = g(x), \quad h(y) := \frac{1}{p(y)}.$$

Let  $H(y)$  and  $G(x)$  be such that

$$H'(y) = h(y), \quad G'(x) = g(x).$$

Then

$$H'(y) \frac{dy}{dx} = G'(x).$$

Since  $\frac{d}{dx}H(y(x)) = H'(y(x)) \frac{dy}{dx}$  (by chain rule), we obtain

$$\frac{d}{dx}H(y(x)) = \frac{d}{dx}G(x) \Rightarrow H(y(x)) = G(x) + C.$$

**Remark:** In finding a one-parameter family of solutions in the separation process, we assume that  $p(y) \neq 0$ . Then we must find the solutions  $y = y_0$  of the equation  $p(y) = 0$  and determine whether any of these are solutions of the original equation which were lost in the formal separation process.

**Example:** Consider  $(x - 4)y^4 dx - x^3(y^2 - 3)dy = 0$ . Separating the variable by dividing  $x^3 y^4$ , we obtain

$$\frac{(x - 4)dx}{x^3} - \frac{(y^2 - 3)dy}{y^4} = 0$$

The general solution is  $-\frac{1}{x} + \frac{2}{x^2} + \frac{1}{y} - \frac{1}{y^3} = C, \quad y \neq 0$

**Note:**  $y = 0$  is a solution of the original equation which was lost in the separation process.

# First-Order Linear Equations

A linear first-order equation can be expressed in the form

$$a_1(x) \frac{dy}{dx} + a_0(x)y = b(x), \quad (3)$$

where  $a_1(x)$ ,  $a_0(x)$  and  $b(x)$  depend only on the independent variable  $x$ , not on  $y$ .

**Examples:**

$$(1 + 2x) \frac{dy}{dx} + 6y = e^x \text{ (linear)}$$

$$\sin x \frac{dy}{dx} + (\cos x)y = x^2 \text{ (linear)}$$

$$\frac{dy}{dx} + xy^3 = x^2 \text{ (not linear)}$$

**Theorem (Existence and Uniqueness):**

Suppose  $a_1(x)$ ,  $a_0(x)$ ,  $b(x) \in C((a, b))$ ,  $a_1(x) \neq 0$  and  $x_0 \in (a, b)$ . Then for any  $y_0 \in \mathbb{R}$ , there exists a unique solution  $y(x) \in C^1((a, b))$  to the IVP

$$a_1(x) \frac{dy}{dx} + a_0(x)y = b(x), \quad y(x_0) = y_0.$$