

Constrained extrema and Lagrange multipliers

Inverse and implicit function theorems

Department of Mathematics
IIT Guwahati

Constrained extrema of $f : \mathbb{R}^n \rightarrow \mathbb{R}$

Let $U \subset \mathbb{R}^n$ be open and $f, g : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be continuous.
Then

Maximize or Minimize $f(\mathbf{x})$
Subject to the constraint $g(\mathbf{x}) = \alpha$.

Constrained extrema of $f : \mathbb{R}^n \rightarrow \mathbb{R}$

Let $U \subset \mathbb{R}^n$ be open and $f, g : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be continuous.
Then

Maximize or Minimize $f(\mathbf{x})$
Subject to the constraint $g(\mathbf{x}) = \alpha$.

Example: Find the extreme values of $f(x, y) = x^2 - y^2$ on the circle $x^2 + y^2 = 1$.

It turns out that f attains minimum at $(0, \pm 1)$ and maximum at $(\pm 1, 0)$ although $\nabla f(0, \pm 1) \neq 0$ and $\nabla f(\pm 1, 0) \neq 0$.

Test for constrained extrema of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

Theorem: Let $f, g : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 . Suppose that f has an extremum at $(a, b) \in U$ such that $g(a, b) = \alpha$ and that $\nabla g(a, b) \neq (0, 0)$. Then there is a $\lambda \in \mathbb{R}$, called **Lagrange multiplier**, such that $\nabla f(a, b) = \lambda \nabla g(a, b)$.

Test for constrained extrema of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

Theorem: Let $f, g : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 . Suppose that f has an extremum at $(a, b) \in U$ such that $g(a, b) = \alpha$ and that $\nabla g(a, b) \neq (0, 0)$. Then there is a $\lambda \in \mathbb{R}$, called **Lagrange multiplier**, such that $\nabla f(a, b) = \lambda \nabla g(a, b)$.

Proof: Let $\mathbf{r}(t)$ be a local parametrization of the curve $g(x, y) = \alpha$ such that $\mathbf{r}(0) = (a, b)$. Then $f(\mathbf{r}(t))$ has an extremum at $t = 0$. Therefore

$$\left. \frac{df(\mathbf{r}(t))}{dt} \right|_{t=0} = \nabla f(a, b) \bullet \mathbf{r}'(0) = 0.$$

Test for constrained extrema of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

Theorem: Let $f, g : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 . Suppose that f has an extremum at $(a, b) \in U$ such that $g(a, b) = \alpha$ and that $\nabla g(a, b) \neq (0, 0)$. Then there is a $\lambda \in \mathbb{R}$, called **Lagrange multiplier**, such that $\nabla f(a, b) = \lambda \nabla g(a, b)$.

Proof: Let $\mathbf{r}(t)$ be a local parametrization of the curve $g(x, y) = \alpha$ such that $\mathbf{r}(0) = (a, b)$. Then $f(\mathbf{r}(t))$ has an extremum at $t = 0$. Therefore

$$\left. \frac{df(\mathbf{r}(t))}{dt} \right|_{t=0} = \nabla f(a, b) \bullet \mathbf{r}'(0) = 0.$$

Now $g(\mathbf{r}(t)) = \alpha \Rightarrow \nabla g(a, b) \bullet \mathbf{r}'(0) = 0$. This shows that $\mathbf{r}'(0) \perp \nabla g(a, b)$ and $\mathbf{r}'(0) \perp \nabla f(a, b)$. Hence $\nabla f(a, b) = \lambda \nabla g(a, b)$ for some $\lambda \in \mathbb{R}$. ■

Method of Lagrange multipliers for $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

To find extremum of f subject to the constraint $g(x, y) = \alpha$, define $L(x, y, \lambda) := f(x, y) - \lambda(g(x, y) - \alpha)$ and solve the equations

Method of Lagrange multipliers for $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

To find extremum of f subject to the constraint $g(x, y) = \alpha$, define $L(x, y, \lambda) := f(x, y) - \lambda(g(x, y) - \alpha)$ and solve the equations

$$\begin{aligned}\frac{\partial L}{\partial x} &= 0 \Rightarrow \frac{\partial f}{\partial x} = \lambda \frac{\partial g}{\partial x}, \\ \frac{\partial L}{\partial y} &= 0 \Rightarrow \frac{\partial f}{\partial y} = \lambda \frac{\partial g}{\partial y}, \\ \frac{\partial L}{\partial \lambda} &= 0 \Rightarrow g(x, y) = \alpha.\end{aligned}$$

Method of Lagrange multipliers for $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

To find extremum of f subject to the constraint $g(x, y) = \alpha$, define $L(x, y, \lambda) := f(x, y) - \lambda(g(x, y) - \alpha)$ and solve the equations

$$\begin{aligned}\frac{\partial L}{\partial x} &= 0 \Rightarrow \frac{\partial f}{\partial x} = \lambda \frac{\partial g}{\partial x}, \\ \frac{\partial L}{\partial y} &= 0 \Rightarrow \frac{\partial f}{\partial y} = \lambda \frac{\partial g}{\partial y}, \\ \frac{\partial L}{\partial \lambda} &= 0 \Rightarrow g(x, y) = \alpha.\end{aligned}$$

- The auxiliary function $L(x, y, \lambda)$ is called **Lagrangian** which converts constrained extrema to unconstrained extrema.

Method of Lagrange multipliers for $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

To find extremum of f subject to the constraint $g(x, y) = \alpha$, define $L(x, y, \lambda) := f(x, y) - \lambda(g(x, y) - \alpha)$ and solve the equations

$$\begin{aligned}\frac{\partial L}{\partial x} &= 0 \Rightarrow \frac{\partial f}{\partial x} = \lambda \frac{\partial g}{\partial x}, \\ \frac{\partial L}{\partial y} &= 0 \Rightarrow \frac{\partial f}{\partial y} = \lambda \frac{\partial g}{\partial y}, \\ \frac{\partial L}{\partial \lambda} &= 0 \Rightarrow g(x, y) = \alpha.\end{aligned}$$

- The auxiliary function $L(x, y, \lambda)$ is called **Lagrangian** which converts constrained extrema to unconstrained extrema.
- Critical points of L are eligible solutions for constrained extrema.

Example

Find the extreme values of $f(x, y) = x^2 - y^2$ on the circle $x^2 + y^2 = 1$.

Example

Find the extreme values of $f(x, y) = x^2 - y^2$ on the circle $x^2 + y^2 = 1$.

The equations $f_x = \lambda g_x$, $f_y = \lambda g_y$ and $g(x, y) = 1$ give $2x = \lambda 2x$, $-2y = \lambda 2y$ and $x^2 + y^2 = 1$. The first equation shows either $x = 0$ or $\lambda = 1$.

Example

Find the extreme values of $f(x, y) = x^2 - y^2$ on the circle $x^2 + y^2 = 1$.

The equations $f_x = \lambda g_x$, $f_y = \lambda g_y$ and $g(x, y) = 1$ give $2x = \lambda 2x$, $-2y = \lambda 2y$ and $x^2 + y^2 = 1$. The first equation shows either $x = 0$ or $\lambda = 1$.

If $x = 0$ then $y = \pm 1 \Rightarrow \lambda = -1$. Thus $(x, y, \lambda) := (0, \pm 1, -1)$ are possible solutions.

Example

Find the extreme values of $f(x, y) = x^2 - y^2$ on the circle $x^2 + y^2 = 1$.

The equations $f_x = \lambda g_x$, $f_y = \lambda g_y$ and $g(x, y) = 1$ give $2x = \lambda 2x$, $-2y = \lambda 2y$ and $x^2 + y^2 = 1$. The first equation shows either $x = 0$ or $\lambda = 1$.

If $x = 0$ then $y = \pm 1 \Rightarrow \lambda = -1$. Thus $(x, y, \lambda) := (0, \pm 1, -1)$ are possible solutions.

If $\lambda = 1$ then $y = 0 \Rightarrow x = \pm 1$. Thus $(x, y, \lambda) := (\pm 1, 0, 1)$ are also possible solutions.

Example

Find the extreme values of $f(x, y) = x^2 - y^2$ on the circle $x^2 + y^2 = 1$.

The equations $f_x = \lambda g_x$, $f_y = \lambda g_y$ and $g(x, y) = 1$ give $2x = \lambda 2x$, $-2y = \lambda 2y$ and $x^2 + y^2 = 1$. The first equation shows either $x = 0$ or $\lambda = 1$.

If $x = 0$ then $y = \pm 1 \Rightarrow \lambda = -1$. Thus $(x, y, \lambda) := (0, \pm 1, -1)$ are possible solutions.

If $\lambda = 1$ then $y = 0 \Rightarrow x = \pm 1$. Thus $(x, y, \lambda) := (\pm 1, 0, 1)$ are also possible solutions.

Now $f(0, 1) = f(0, -1) = -1$ and $f(1, 0) = f(-1, 0) = 1$ so that minimum and maximum values are -1 and 1 .

Finding global extrema of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $U \subset \mathbb{R}^2$ be a region with smooth closed boundary curve C . To find global extremum of f in U :

Finding global extrema of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $U \subset \mathbb{R}^2$ be a region with smooth closed boundary curve C . To find global extremum of f in U :

- Locate all critical points of f in U .

Finding global extrema of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $U \subset \mathbb{R}^2$ be a region with smooth closed boundary curve C . To find global extremum of f in U :

- Locate all critical points of f in U .
- Find eligible global extremum of f on the curve C by using Lagrange multipliers or parametrization.

Finding global extrema of $f : \mathbb{R}^2 \rightarrow \mathbb{R}$

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $U \subset \mathbb{R}^2$ be a region with smooth closed boundary curve C . To find global extremum of f in U :

- Locate all critical points of f in U .
- Find eligible global extremum of f on the curve C by using Lagrange multipliers or parametrization.
- Choose points among eligible solutions in C and the critical points at which f attains extreme values. These extreme values are global extremum.

Example: global extrema

Find global maximum and global minimum of the function $f(x, y) := (x^2 + y^2)/2$ such that $x^2/2 + y^2 \leq 1$.

Example: global extrema

Find global maximum and global minimum of the function $f(x, y) := (x^2 + y^2)/2$ such that $x^2/2 + y^2 \leq 1$.

Since $f_x = x$ and $f_y = y$, $(0, 0)$ is the only critical point.

Example: global extrema

Find global maximum and global minimum of the function $f(x, y) := (x^2 + y^2)/2$ such that $x^2/2 + y^2 \leq 1$.

Since $f_x = x$ and $f_y = y$, $(0, 0)$ is the only critical point.

Next, consider $L(x, y, \lambda) := (x^2 + y^2)/2 - \lambda(x^2/2 + y^2 - 1)$. Then Lagrange multiplier equations are

$$x = \lambda x, \quad y = 2\lambda y, \quad x^2/2 + y^2 = 1.$$

If $x = 0$ then $y = \pm 1$ and $\lambda = 1/2$. If $y = 0$ then $x = \pm\sqrt{2}$ and $\lambda = 1$. If $xy \neq 0$ then $\lambda = 1$ and $\lambda = 1/2$ -which is not possible.

Example: global extrema

Find global maximum and global minimum of the function $f(x, y) := (x^2 + y^2)/2$ such that $x^2/2 + y^2 \leq 1$.

Since $f_x = x$ and $f_y = y$, $(0, 0)$ is the only critical point.

Next, consider $L(x, y, \lambda) := (x^2 + y^2)/2 - \lambda(x^2/2 + y^2 - 1)$. Then Lagrange multiplier equations are

$$x = \lambda x, \quad y = 2\lambda y, \quad x^2/2 + y^2 = 1.$$

If $x = 0$ then $y = \pm 1$ and $\lambda = 1/2$. If $y = 0$ then $x = \pm\sqrt{2}$ and $\lambda = 1$. If $xy \neq 0$ then $\lambda = 1$ and $\lambda = 1/2$ -which is not possible.

Thus $(0, \pm 1)$ and $(\pm\sqrt{2}, 0)$ are eligible solutions for the boundary curve. We have $f(0, \pm 1) = 1/2$, $f(\pm\sqrt{2}, 0) = 1$ and $f(0, 0) = 0$.

Test for constrained extrema of $f : \mathbb{R}^n \rightarrow \mathbb{R}$

Theorem: Let $f, g : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be C^1 . Suppose that f has an extremum at $\mathbf{p} \in U$ such that $g(\mathbf{p}) = \alpha$ and $\nabla g(\mathbf{p}) \neq \mathbf{0}$. Then there is a $\lambda \in \mathbb{R}$ such that $\nabla f(\mathbf{p}) = \lambda \nabla g(\mathbf{p})$.

Test for constrained extrema of $f : \mathbb{R}^n \rightarrow \mathbb{R}$

Theorem: Let $f, g : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be C^1 . Suppose that f has an extremum at $\mathbf{p} \in U$ such that $g(\mathbf{p}) = \alpha$ and $\nabla g(\mathbf{p}) \neq \mathbf{0}$. Then there is a $\lambda \in \mathbb{R}$ such that $\nabla f(\mathbf{p}) = \lambda \nabla g(\mathbf{p})$.

Proof:

Test for constrained extrema of $f : \mathbb{R}^n \rightarrow \mathbb{R}$

Theorem: Let $f, g : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be C^1 . Suppose that f has an extremum at $\mathbf{p} \in U$ such that $g(\mathbf{p}) = \alpha$ and $\nabla g(\mathbf{p}) \neq \mathbf{0}$. Then there is a $\lambda \in \mathbb{R}$ such that $\nabla f(\mathbf{p}) = \lambda \nabla g(\mathbf{p})$.

Proof:

If $g(\mathbf{x}) = \alpha$ is a closed surface then global extremum is obtained by finding all points where $\nabla f(\mathbf{x}) = \lambda \nabla g(\mathbf{x})$ and choosing those where f is largest or smallest.

Test for constrained extrema of $f : \mathbb{R}^n \rightarrow \mathbb{R}$

Theorem: Let $f, g : U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be C^1 . Suppose that f has an extremum at $\mathbf{p} \in U$ such that $g(\mathbf{p}) = \alpha$ and $\nabla g(\mathbf{p}) \neq \mathbf{0}$. Then there is a $\lambda \in \mathbb{R}$ such that $\nabla f(\mathbf{p}) = \lambda \nabla g(\mathbf{p})$.

Proof:

If $g(\mathbf{x}) = \alpha$ is a closed surface then global extremum is obtained by finding all points where $\nabla f(\mathbf{x}) = \lambda \nabla g(\mathbf{x})$ and choosing those where f is largest or smallest.

The Lagrangian is given by $L(\mathbf{x}, \lambda) := f(\mathbf{x}) - \lambda(g(\mathbf{x}) - \alpha)$. So, the multiplier equations are $\nabla L(\mathbf{x}, \lambda) = \mathbf{0}$.

Example

Maximize the function $f(x, y, z) := x + z$ subject to the constraint $x^2 + y^2 + z^2 = 1$.

Example

Maximize the function $f(x, y, z) := x + z$ subject to the constraint $x^2 + y^2 + z^2 = 1$.

The multiplier equations are

$$1 = 2x\lambda, \quad 0 = 2y\lambda, \quad 1 = 2z\lambda, \quad x^2 + y^2 + z^2 = 1.$$

From first and 3rd equation, $\lambda \neq 0$. Thus, by second equation, $y = 0$. By first and 3rd equations $x = z \Rightarrow x = z = \pm 1/\sqrt{2}$.

Example

Maximize the function $f(x, y, z) := x + z$ subject to the constraint $x^2 + y^2 + z^2 = 1$.

The multiplier equations are

$$1 = 2x\lambda, \quad 0 = 2y\lambda, \quad 1 = 2z\lambda, \quad x^2 + y^2 + z^2 = 1.$$

From first and 3rd equation, $\lambda \neq 0$. Thus, by second equation, $y = 0$. By first and 3rd equations $x = z \Rightarrow x = z = \pm 1/\sqrt{2}$.

Hence $\mathbf{p} := (1/\sqrt{2}, 0, 1/\sqrt{2})$ and $\mathbf{q} := (-1/\sqrt{2}, 0, -1/\sqrt{2})$ are eligible solutions. This shows that $f(\mathbf{p}) = \sqrt{2}$ and $f(\mathbf{q}) = -\sqrt{2}$.

What does the Implicit function theorem say?

Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 . Consider the curve

$$V(F) := \{(x, y) \in \mathbb{R}^2 : F(x, y) = 0\}.$$

Does there exist $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $V(F) = \text{Graph}(f)$?

Equivalently, can $F(x, y) = 0$ be solved either for x or for y ?

What does the Implicit function theorem say?

Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 . Consider the curve

$$V(F) := \{(x, y) \in \mathbb{R}^2 : F(x, y) = 0\}.$$

Does there exist $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $V(F) = \text{Graph}(f)$?

Equivalently, can $F(x, y) = 0$ be solved either for x or for y ?

Considering $F(x, y) := x^2 + y^2 - 1$, it follows that $F(x, y) = 0$ cannot be solved for y or x .

What does the Implicit function theorem say?

Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 . Consider the curve

$$V(F) := \{(x, y) \in \mathbb{R}^2 : F(x, y) = 0\}.$$

Does there exist $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $V(F) = \text{Graph}(f)$?

Equivalently, can $F(x, y) = 0$ be solved either for x or for y ?

Considering $F(x, y) := x^2 + y^2 - 1$, it follows that $F(x, y) = 0$ cannot be solved for y or x .

The **implicit function theorem** says that if $F(a, b) = 0$ and $\nabla F(a, b) \neq (0, 0)$ then in a **neighbourhood of (a, b)** , we have

$$V(F) = \text{Graph}(f)$$

for some function f .

Implicit function theorem

Theorem: Let $F : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 , where U is open. Consider the curve $V(F) := \{(x, y) \in U : F(x, y) = 0\}$. Let $(a, b) \in V(F)$. Suppose that $\partial_y F(a, b) \neq 0$.

Implicit function theorem

Theorem: Let $F : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 , where U is open. Consider the curve $V(F) := \{(x, y) \in U : F(x, y) = 0\}$. Let $(a, b) \in V(F)$. Suppose that $\partial_y F(a, b) \neq 0$.

- Then there exists $r > 0$ and a C^1 function $g : (a - r, a + r) \rightarrow \mathbb{R}$ such that $F(x, g(x)) = 0$ for $x \in (a - r, a + r)$.

Implicit function theorem

Theorem: Let $F : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 , where U is open. Consider the curve $V(F) := \{(x, y) \in U : F(x, y) = 0\}$. Let $(a, b) \in V(F)$. Suppose that $\partial_y F(a, b) \neq 0$.

- Then there exists $r > 0$ and a C^1 function $g : (a - r, a + r) \rightarrow \mathbb{R}$ such that $F(x, g(x)) = 0$ for $x \in (a - r, a + r)$.
- For $W := (a - r, a + r) \times (b - r, b + r)$, we have

$$W \cap V(F) = \text{Graph}(g).$$

Implicit function theorem

Theorem: Let $F : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be C^1 , where U is open. Consider the curve $V(F) := \{(x, y) \in U : F(x, y) = 0\}$. Let $(a, b) \in V(F)$. Suppose that $\partial_y F(a, b) \neq 0$.

- Then there exists $r > 0$ and a C^1 function $g : (a - r, a + r) \rightarrow \mathbb{R}$ such that $F(x, g(x)) = 0$ for $x \in (a - r, a + r)$.
- For $W := (a - r, a + r) \times (b - r, b + r)$, we have

$$W \cap V(F) = \text{Graph}(g).$$

- Further, $\partial_x F(a, b) + \partial_y F(a, b)g'(a) = 0$.

Implicit derivative

Thus if $\partial_y f(a, b) \neq 0$ then in some disk about (a, b) the set of points (x, y) satisfying $F(x, y) = 0$ is the graph of a function $y = g(x)$ with

$$\frac{dy}{dx}\bigg|_{x=a} = g'(a) = -F_x(a, b)/F_y(a, b).$$

Implicit derivative

Thus if $\partial_y f(a, b) \neq 0$ then in some disk about (a, b) the set of points (x, y) satisfying $F(x, y) = 0$ is the graph of a function $y = g(x)$ with

$$\frac{dy}{dx}\bigg|_{x=a} = g'(a) = -F_x(a, b)/F_y(a, b).$$

Example: Consider $F(x, y) := e^{x-2+(y-1)^2} - 1$ and the equation $F(x, y) = 0$. Then $F(2, 1) = 0$, $\partial_x F(2, 1) = 1$, and $\partial_y F(2, 1) = 0$.

Hence $x = g(y)$ for some C^1 function $g : (1 - r, 1 + r) \rightarrow \mathbb{R}$. Moreover, $g'(1) = 0$.

Implicit function theorem for $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$

Let $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ be C^1 , where U is open. Consider the level set $V(F) := \{(\mathbf{x} \in U : F(\mathbf{x}, y) = 0\}$. Let $(\mathbf{a}, b) \in V(F)$. Suppose that $\partial_y F(\mathbf{a}, b) \neq 0$.

Implicit function theorem for $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$

Let $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ be C^1 , where U is open. Consider the level set $V(F) := \{(\mathbf{x} \in U : F(\mathbf{x}, y) = 0\}$. Let $(\mathbf{a}, b) \in V(F)$. Suppose that $\partial_y F(\mathbf{a}, b) \neq 0$.

- Then there exists $r > 0$ and a C^1 function $g : B(\mathbf{a}, r) \rightarrow \mathbb{R}$ such that $F(\mathbf{x}, g(\mathbf{x})) = 0$ for $\mathbf{x} \in B(\mathbf{a}, r)$.

Implicit function theorem for $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$

Let $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ be C^1 , where U is open. Consider the level set $V(F) := \{(\mathbf{x} \in U : F(\mathbf{x}, y) = 0\}$. Let $(\mathbf{a}, b) \in V(F)$. Suppose that $\partial_y F(\mathbf{a}, b) \neq 0$.

- Then there exists $r > 0$ and a C^1 function $g : B(\mathbf{a}, r) \rightarrow \mathbb{R}$ such that $F(\mathbf{x}, g(\mathbf{x})) = 0$ for $\mathbf{x} \in B(\mathbf{a}, r)$.
- For $W := B(\mathbf{a}, r) \times (b - r, b + r)$, we have

$$W \cap V(F) = \text{Graph}(g).$$

Implicit function theorem for $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$

Let $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ be C^1 , where U is open. Consider the level set $V(F) := \{(\mathbf{x} \in U : F(\mathbf{x}, y) = 0\}$. Let $(\mathbf{a}, b) \in V(F)$. Suppose that $\partial_y F(\mathbf{a}, b) \neq 0$.

- Then there exists $r > 0$ and a C^1 function $g : B(\mathbf{a}, r) \rightarrow \mathbb{R}$ such that $F(\mathbf{x}, g(\mathbf{x})) = 0$ for $\mathbf{x} \in B(\mathbf{a}, r)$.
- For $W := B(\mathbf{a}, r) \times (b - r, b + r)$, we have

$$W \cap V(F) = \text{Graph}(g).$$

- Further, $\partial_i F(\mathbf{a}, b) + \partial_y F(\mathbf{a}, b) \partial_i g(\mathbf{a}) = 0, i = 1, 2, \dots, n$

Inverse function theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be linear and represented (standard basis) by an $n \times n$ matrix A . Then f is invertible on $\mathbb{R}^n \Leftrightarrow \det(A) \neq 0$.

Inverse function theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be linear and represented (standard basis) by an $n \times n$ matrix A . Then f is invertible on $\mathbb{R}^n \Leftrightarrow \det(A) \neq 0$.

Note that $J_f(\mathbf{x}) = A$. Thus $\det(J_f(\mathbf{x})) \neq 0 \implies f$ is invertible on \mathbb{R}^n .

Inverse function theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be linear and represented (standard basis) by an $n \times n$ matrix A . Then f is **invertible** on $\mathbb{R}^n \Leftrightarrow \det(A) \neq 0$.

Note that $J_f(\mathbf{x}) = A$. Thus $\det(J_f(\mathbf{x})) \neq 0 \implies f$ is invertible on \mathbb{R}^n .

Example: Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, (x, y) \mapsto (e^x \cos y, e^x \sin y)$.
Then

$$J_f(x, y) = \begin{bmatrix} e^x \cos y & -e^x \sin y \\ e^x \sin y & e^x \cos y \end{bmatrix} \Rightarrow \det(J_f(x, y)) = e^{2x} \neq 0.$$

Inverse function theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be linear and represented (standard basis) by an $n \times n$ matrix A . Then f is invertible on $\mathbb{R}^n \Leftrightarrow \det(A) \neq 0$.

Note that $J_f(\mathbf{x}) = A$. Thus $\det(J_f(\mathbf{x})) \neq 0 \Rightarrow f$ is invertible on \mathbb{R}^n .

Example: Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, (x, y) \mapsto (e^x \cos y, e^x \sin y)$.
Then

$$J_f(x, y) = \begin{bmatrix} e^x \cos y & -e^x \sin y \\ e^x \sin y & e^x \cos y \end{bmatrix} \Rightarrow \det(J_f(x, y)) = e^{2x} \neq 0.$$

But $f(x, y) = f(x, y + 2\pi) \Rightarrow f$ is not invertible on \mathbb{R}^2 .

Inverse function theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be linear and represented (standard basis) by an $n \times n$ matrix A . Then f is **invertible** on $\mathbb{R}^n \Leftrightarrow \det(A) \neq 0$.

Note that $J_f(\mathbf{x}) = A$. Thus $\det(J_f(\mathbf{x})) \neq 0 \implies f$ is invertible on \mathbb{R}^n .

Example: Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, (x, y) \mapsto (e^x \cos y, e^x \sin y)$.
Then

$$J_f(x, y) = \begin{bmatrix} e^x \cos y & -e^x \sin y \\ e^x \sin y & e^x \cos y \end{bmatrix} \Rightarrow \det(J_f(x, y)) = e^{2x} \neq 0.$$

But $f(x, y) = f(x, y + 2\pi) \Rightarrow f$ is not invertible on \mathbb{R}^2 .

Moral: Nonsingularity of $J_f(\mathbf{x})$ does not guarantee invertibility of $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ on \mathbb{R}^n .

Fact: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be C^1 and $f'(x_0) \neq 0$. Then

- f is invertible in a neighborhood of x_0 ,
- the inverse is continuously differentiable, and
- $(f^{-1})'(f(x_0)) = \frac{1}{f'(x_0)}$.

Fact: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be C^1 and $f'(x_0) \neq 0$. Then

- f is invertible in a neighborhood of x_0 ,
- the inverse is continuously differentiable, and
- $(f^{-1})'(f(x_0)) = \frac{1}{f'(x_0)}$.

Theorem: Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be C^1 and $\det(J_f(\mathbf{a})) \neq 0$. Then there are open subsets $U \subset \mathbb{R}^n$ and $V \subset \mathbb{R}^n$ such that

- $\mathbf{a} \in U$ and $f(\mathbf{a}) \in V$,
- $f : U \rightarrow V$ is bijective,
- $f^{-1} : V \rightarrow U$ is C^1 and $J_{f^{-1}}(f(\mathbf{a})) = (J_f(\mathbf{a}))^{-1}$.

Fact: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be C^1 and $f'(x_0) \neq 0$. Then

- f is invertible in a neighborhood of x_0 ,
- the inverse is continuously differentiable, and
- $(f^{-1})'(f(x_0)) = \frac{1}{f'(x_0)}$.

Theorem: Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be C^1 and $\det(J_f(\mathbf{a})) \neq 0$. Then there are open subsets $U \subset \mathbb{R}^n$ and $V \subset \mathbb{R}^n$ such that

- $\mathbf{a} \in U$ and $f(\mathbf{a}) \in V$,
- $f : U \rightarrow V$ is bijective,
- $f^{-1} : V \rightarrow U$ is C^1 and $J_{f^{-1}}(f(\mathbf{a})) = (J_f(\mathbf{a}))^{-1}$.
- **Stronger version:** $J_{f^{-1}}(\mathbf{y}) = (J_f(f^{-1}(\mathbf{y})))^{-1}$ for $\mathbf{y} \in V$.

Example

Consider the system $u = x \cos y$, $v = x \sin y$. Then x and y can be expressed as C^1 functions of (u, v) in a neighbourhood of (a, b) when $a \neq 0$.

Example

Consider the system $u = x \cos y$, $v = x \sin y$. Then x and y can be expressed as C^1 functions of (u, v) in a neighbourhood of (a, b) when $a \neq 0$.

Let $f(x, y) := (x \cos y, x \sin y)$. Then

$$J_f(a, b) = \begin{bmatrix} \cos b & -a \sin b \\ \sin b & a \cos b \end{bmatrix} \Rightarrow \det(J_f(a, b)) = a.$$

Example

Consider the system $u = x \cos y$, $v = x \sin y$. Then x and y can be expressed as C^1 functions of (u, v) in a neighbourhood of (a, b) when $a \neq 0$.

Let $f(x, y) := (x \cos y, x \sin y)$. Then

$$J_f(a, b) = \begin{bmatrix} \cos b & -a \sin b \\ \sin b & a \cos b \end{bmatrix} \Rightarrow \det(J_f(a, b)) = a.$$

Set $(x, y) := f^{-1}(u, v)$. Then

$$\begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix} = J_{f^{-1}}(f(a, b)) = \begin{bmatrix} \cos b & -a \sin b \\ \sin b & a \cos b \end{bmatrix}^{-1}.$$

*** End ***