

Adverse Events of CTCAE Grade 3 or Higher by SOC and PT (Safety Analysis Set)

		Number (%) of patients	
System Organ Class	Preferred Term	TREATMENT A	TREATMENT B
BLOOD AND LYMPHATIC SYSTEM DISORDERS	Anaemia	9 (11.8%)	3 (3.9%)
	Neutropenia	2 (2.6%)	0 (0.0%)
CARDIAC DISORDERS	Acute coronary syndrome	1 (1.3%)	0 (0.0%)
	Atrioventricular block	0 (0.0%)	1 (1.3%)
EAR AND LABYRINTH DISORDERS	Vertigo positional	1 (1.3%)	0 (0.0%)
ENDOCRINE DISORDERS	Adrenal insufficiency	0 (0.0%)	1 (1.3%)
GASTROINTESTINAL DISORDERS	Abdominal pain	1 (1.3%)	0 (0.0%)
	Colitis	0 (0.0%)	1 (1.3%)
	Gastric haemorrhage	1 (1.3%)	0 (0.0%)
	Gastric ulcer perforation	0 (0.0%)	1 (1.3%)
	Intestinal obstruction	0 (0.0%)	1 (1.3%)
	Nausea	1 (1.3%)	0 (0.0%)
	Small intestinal obstruction	1 (1.3%)	0 (0.0%)
	Upper gastrointestinal haemorrhage	0 (0.0%)	1 (1.3%)
GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS	Asthenia	1 (1.3%)	0 (0.0%)
	Fatigue	2 (2.6%)	1 (1.3%)
	Pyrexia	1 (1.3%)	1 (1.3%)
HEPATOBIILIARY DISORDERS	Drug-induced liver injury	1 (1.3%)	0 (0.0%)
INFECTIONS AND INFESTATIONS	Escherichia bacteraemia	0 (0.0%)	1 (1.3%)
	Kidney infection	1 (1.3%)	0 (0.0%)
	Peritonitis	1 (1.3%)	0 (0.0%)
	Pneumonia	2 (2.6%)	2 (2.6%)
	Septic shock	1 (1.3%)	0 (0.0%)
	Urinary tract infection	6 (7.9%)	4 (5.3%)
INJURY, POISONING AND PROCEDURAL COMPLICATIONS	Femur fracture	0 (0.0%)	1 (1.3%)
	Hip fracture	1 (1.3%)	0 (0.0%)

AE onset >= first dose and <= 28 days after TRTEDT; Denominator = SAFFL='Y' from ADSL

Adverse Events of CTCAE Grade 3 or Higher by SOC and PT (Safety Analysis Set)

		Number (%) of patients	
System Organ Class	Preferred Term	TREATMENT A	TREATMENT B
INVESTIGATIONS	Urostomy complication	1 (1.3%)	0 (0.0%)
	Amylase increased	0 (0.0%)	1 (1.3%)
	Blood creatinine increased	1 (1.3%)	0 (0.0%)
	Gamma-glutamyltransferase increased	0 (0.0%)	2 (2.6%)
	Lipase increased	0 (0.0%)	1 (1.3%)
	Neutrophil count decreased	1 (1.3%)	0 (0.0%)
METABOLISM AND NUTRITION DISORDERS	Dehydration	1 (1.3%)	0 (0.0%)
	Hyperglycaemia	1 (1.3%)	1 (1.3%)
	Hyperkalaemia	1 (1.3%)	1 (1.3%)
	Hypermagnesaemia	1 (1.3%)	0 (0.0%)
	Hypoglycaemia	0 (0.0%)	1 (1.3%)
	Hypokalaemia	0 (0.0%)	1 (1.3%)
	Hyponatraemia	1 (1.3%)	1 (1.3%)
	Type 2 diabetes mellitus	0 (0.0%)	1 (1.3%)
MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS	Bone pain	1 (1.3%)	0 (0.0%)
	Pain in extremity	0 (0.0%)	1 (1.3%)
PSYCHIATRIC DISORDERS	Anxiety	0 (0.0%)	1 (1.3%)
RENAL AND URINARY DISORDERS	Acute kidney injury	4 (5.3%)	4 (5.3%)
	Chronic kidney disease	1 (1.3%)	1 (1.3%)
	Haematuria	1 (1.3%)	1 (1.3%)
	Hydronephrosis	3 (3.9%)	0 (0.0%)
	Renal failure	4 (5.3%)	0 (0.0%)
	Renal impairment	1 (1.3%)	0 (0.0%)
	Ureteric obstruction	1 (1.3%)	0 (0.0%)
	Urinary bladder haemorrhage	0 (0.0%)	1 (1.3%)
	Urinary retention	0 (0.0%)	1 (1.3%)
REPRODUCTIVE SYSTEM AND BREAST DISORDERS	Vaginal haemorrhage	0 (0.0%)	1 (1.3%)
RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS	Immune-mediated pneumonitis	1 (1.3%)	0 (0.0%)

AE onset >= first dose and <= 28 days after TRTEDT; Denominator = SAFFL='Y' from ADSL

Adverse Events of CTCAE Grade 3 or Higher by SOC and PT (Safety Analysis Set)

		Number (%) of patients	
System Organ Class	Preferred Term	TREATMENT A	TREATMENT B
	Pneumonia aspiration	1 (1.3%)	0 (0.0%)
	Pulmonary thrombosis	1 (1.3%)	0 (0.0%)
	Respiratory failure	1 (1.3%)	0 (0.0%)
SKIN AND SUBCUTANEOUS TISSUE DISORDERS	Psoriasis	0 (0.0%)	1 (1.3%)
	Rash	0 (0.0%)	1 (1.3%)
VASCULAR DISORDERS	Hypertension	1 (1.3%)	1 (1.3%)
	Hypotension	1 (1.3%)	0 (0.0%)
ZZZ	Any Grade >=3 AE	34 (44.7%)	27 (35.5%)

AE onset >= first dose and <= 28 days after TRTEDT; Denominator = SAFFL='Y' from ADSL