

# FYS3150/FYS4150

## Project 1

Magnus Ulimoen  
Krister Stræte Karlsen

September 2, 2015

### 1 Motivation

Solutions of the Poissons equation are a vital component to many aspects physics.

### 2 a)

#### Formulation

Given the following ODE with boundary conditions,

$$-u''(x) = f(x), \quad x \in (0, 1), \quad u(0) = u(1) = 0. \quad (1)$$

and the discretized form following a symmetric Taylor expansion

$$-\frac{v_{i+1} + v_{i-1} - 2v_i}{h^2} = f_i \quad \text{for } i = 1, \dots, n, \quad v_0 = v_n = 0 \quad (2)$$

we are going to show it can be written as system of linear equations of the form:

$$\mathbf{A}\mathbf{v} = \tilde{\mathbf{b}}, \quad (3)$$

## Solution

Multiplying the discretized equation (2) by  $h^2$  we get:

$$-v_{i-1} + 2v_i - v_{i+1} = h^2 f_i \quad \text{for } i = 1, \dots, n$$

Filling in for  $i$  and choosing  $\tilde{b}_i = h^2 f_i$  we obtain the following set of equations:

$$\begin{aligned} 2v_1 - v_2 &= \tilde{b}_1 \\ -v_1 + 2v_2 - v_3 &= \tilde{b}_2 \\ &\vdots \\ -v_{i-1} + 2v_i - v_{i+1} &= \tilde{b}_i \\ &\vdots \\ -v_{n-1} + 2v_n &= \tilde{b}_n \end{aligned}$$

Now one can easily see that this system of linear equations can be written in the form of (3), where

$$\mathbf{A} = \begin{pmatrix} 2 & -1 & 0 & \dots & \dots & 0 \\ -1 & 2 & -1 & 0 & \dots & \dots \\ 0 & -1 & 2 & -1 & 0 & \dots \\ & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & & -1 & 2 & -1 \\ 0 & \dots & & 0 & -1 & 2 \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ \dots \\ \dots \\ \dots \\ v_n \end{pmatrix}, \quad \tilde{\mathbf{b}} = \begin{pmatrix} \tilde{b}_1 \\ \tilde{b}_2 \\ \dots \\ \dots \\ \dots \\ \tilde{b}_n \end{pmatrix}.$$

## Algorithm for a tridiagonal system of linear equations

We start by looking at the system of equations:

$$\begin{aligned} b_1 v_1 + c_1 v_2 &= \tilde{b}_1 & (1*) \\ a_2 v_1 + b_2 v_2 + c_3 v_3 &= \tilde{b}_2 & (2*) \\ a_3 v_2 + b_3 v_3 + c_3 v_4 &= \tilde{b}_3 & (3*) \\ &\vdots \\ a_n v_{n-1} + b_n v_n &= \tilde{b}_n & (n*) \end{aligned}$$

If we solve (1\*) for  $v_1$  and insert it into (2\*) we obtain the following "modified second equation":

$$(b_1 b_2 - a_2 c_1) v_2 + b_1 c_2 v_3 = b_1 \tilde{b}_2 - a_2 \tilde{b}_1$$

Now having successfully removed  $v_1$  from the second equation we can go on and solve it for  $v_2$  and insert it into the third equation obtaining:

$$(b_3(b_1 b_2 - a_2 c_1) - a_3 b_1 c_2) v_3 + c_3(b_1 b_2 - a_2 c_1) v_4 = (b_1 b_2 - a_2 c_1) \tilde{b}_3 - a_3 b_1 \tilde{b}_2 + a_2 a_3 \tilde{b}_1$$

The two modified equations may be written as

$$\begin{aligned} v_2 &= \frac{b_1 \tilde{b}_2 - a_2 \tilde{b}_1}{b_1 b_2 - a_2 c_1} - \frac{b_1 c_2}{b_1 b_2 - a_2 c_1} v_3 = \beta_3 + \gamma_3 v_3 \\ v_3 &= \frac{(b_1 b_2 - a_2 c_1) \tilde{b}_3 - a_3(b_1 \tilde{b}_2 - a_2 \tilde{b}_1)}{b_3(b_1 b_2 - a_2 c_1) - a_3 b_1 c_2} - \frac{c_3(b_1 b_2 - a_2 c_1)}{b_3(b_1 b_2 - a_2 c_1) - a_3 b_1 c_2} v_4 \\ &= \beta_4 + \gamma_4 v_4 = \frac{\tilde{b}_3 - a_3 \beta_3}{a_3 \gamma_3 + b_3} + \frac{-c_3}{a_3 \gamma_3 + b_3} v_4 \end{aligned}$$

This process can be repeated up until the last equation. This is the forward substitution step. From the last equation we compute  $v_n$  and get all we need to compute  $v_{n-1}$ , then  $v_{n-2}$ , and so on. This is the backward substitution part of the algorithm. A shrewd reader might see that the coefficients,  $\beta$  and  $\gamma$ , take a recursive form

$$\beta_{i+1} = \frac{\tilde{b}_i - a_i \beta_i}{a_i \gamma_i + b_i}, \quad \gamma_{i+1} = \frac{-c_i}{a_i \gamma_i + b_i},$$

and the equation for  $v_{i-1}$  reads:

$$v_{i-1} = \beta_i + \gamma_i v_i \tag{4}$$

It follows from (1\*) that  $\beta_1 = \gamma_1 = 0$ . From combining (n\*) and (4) we get

$$v_n = \frac{\tilde{b}_n - a_n \beta_n}{a_n \gamma_n + b_n} = \beta_{n+1} + \gamma_{n+1} v_{n+1}, \quad v_{n+1} = 0.$$

The goal of the IMIDLERTID stupid formulation above is to underline the importance of setting  $v_{n+1} = 0$  in order to obtain the right formula for  $v_n$ .

Having all the nessecary ingredients the algorithm reads as follows.

#### Algorithm I

```

 $a_i = c_i = -1, \quad i = 1, 2, 3, \dots, n$ 
 $b_i = 2, \quad i = 1, 2, 3, \dots, n$ 
 $\beta_1 = \gamma_1 = 0$ 
for  $i = 1, 2, \dots, n - 1$ 
     $\beta_{i+1} = \frac{\tilde{b}_i - a_i \beta_i}{a_i \gamma_i + b_i}, \quad \gamma_{i+1} = \frac{-c_i}{a_i \gamma_i + b_i}$ 

 $v_{n+1} = 0$ 
for  $j = n + 1, n, \dots, 1$ 
     $v_{i-1} = \beta_i + \gamma_i v_i$ 

```

This is often referred to as *Thomas Algorithm*, an algorithm for solving tridiagonal systems of linear equations.