

Update_Writeup

Aidan Mullan

2/26/2019

```
## Warning in read_fun(path = path, sheet_i = sheet, limits = limits, shim =  
## shim, : Expecting numeric in I1195 / R1195C9: got '100'  
## Warning: NAs introduced by coercion
```

Effective Dose

Summary Statistics

Using the fixed data for effective dosage of radiation administered to patients, we retrieve the plots shown in Figure 1. Here we can see the right-skew in the histogram suggesting several patients recieved much higher amounts of radiation than average. The mean effective dose given to a patient was 5.631 millisieverts, with a range of 0.1 to 48.8 millisieverts. Out of all 2382 patients, only 54 recieved a radiation dose above 20 millisieverts.

Table 1: Summary Statistics of Radiation and Contrast

	Min	1st Q	Median	Mean	3rd Q	Max
Effective Dose	0.10	3.06	4.30	5.631	6.36	48.83
Contrast	5.0	75.0	95.0	88.5	98.0	150.0

Figure 1a: Histogram of Effective Dose





Figure 1b: Boxplot of Effective Dose




##	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
##	0.098	3.055	4.298	5.631	6.356	48.832

We are also interested in the relationship between the effective dose administered and both the age and BMI of the patient. For the case of age, patients were grouped into 10-year age groups. The number of patients that belong to group is given below. Figure 2 shows the average radiation administered for each of these age groups.

##	<20	20-30	30-40	40-50	50-60	60-70	70-80	80-90	>90
##	55	133	222	294	514	562	397	179	26

Figure 2: Mean Effective Dose by Age



Contrast Given