

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A \text{ where } X_A = \frac{\text{moles } A}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = {}^\circ\text{C} + 273$$

$$D = \frac{m}{V}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

Molarity, M = moles of solute per liter of solution

$$A = abc$$

$$\text{Gas constant, } R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$= 0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1}$$

$$= 62.36 \text{ L torr mol}^{-1} \text{ K}^{-1}$$

$$1 \text{ atm} = 760 \text{ mm Hg}$$

$$= 760 \text{ torr}$$

$$\text{STP} = 0.00^\circ\text{C and } 1.000 \text{ atm}$$

THERMOCHEMISTRY/ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

Faraday's constant, $F = 96,485$ coulombs per mole of electrons

$$1 \text{ volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$

Appendix B: AP Chemistry Equations and Constants

Throughout the test the following symbols have the definitions specified unless otherwise noted.

$$\begin{aligned} \text{mm Hg} &= \text{millimeters of mercury} \\ \text{J, kJ} &= \text{joule(s), kilojoule(s)} \\ V &= \text{volt(s)} \\ \text{mol} &= \text{mole(s)} \end{aligned}$$

$$\begin{aligned} \text{L, mL} &= \text{liter(s), milliliter(s)} \\ g &= \text{gram(s)} \\ \text{nm} &= \text{nanometer(s)} \\ \text{atm} &= \text{atmosphere(s)} \end{aligned}$$

ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

$$\begin{aligned} E &= \text{energy} \\ \nu &= \text{frequency} \\ \lambda &= \text{wavelength} \end{aligned}$$

$$\text{Planck's constant, } h = 6.626 \times 10^{-34} \text{ J s}$$

$$\text{Speed of light, } c = 2.998 \times 10^8 \text{ m s}^{-1}$$

$$\text{Avogadro's number} = 6.022 \times 10^{23} \text{ mol}^{-1}$$

$$\text{Electron charge, } e = -1.602 \times 10^{-19} \text{ coulomb}$$

EQUILIBRIUM

$$K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}, \text{ where } aA + bB \rightleftharpoons cC + dD$$

Equilibrium Constants

K_c (molar concentrations)

K_p (gas pressures)

K_a (weak acid)

K_b (weak base)

K_w (water)

$$K_p = \frac{(P_C)^c(P_D)^d}{(P_A)^a(P_B)^b}$$

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[H^+], \text{ pOH} = -\log[OH^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log\left(\frac{[A^-]}{[HA]}\right)$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

KINETICS

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

$$k = \text{rate constant}$$

$$t = \text{time}$$

$$t_{1/2} = \text{half-life}$$