

Vitamin C



- Reduce Stress in Dairy cows & Poultry.
- Protects the cells from Oxidative stress and damage.
- Assist liver cells to detox mycotoxins and heavy metals.

Vitamin C

Composition

L-Ascorbic Acid > 97%, a white crystalline powder

DAIRY Indications

- Vitamin-C reduce stress in dairy cows particularly heat stress and transition stress.
- Vitamin-C protects the cells from oxidative stress and damage.
- Vitamin-C assist liver cells to detox mycotoxins and heavy metals.
- Vitamin-C is involved in collagen synthesis which can support hoof health.

Dosage

Dairy Cow: 10 gm / Cow / day
Calf : 01 gm / day

POULTRY Indications

- Support in case of stress such as heat stress, advance climate condition, transportation, pullet replacement, overcrowding, culling, vaccination and anthelmintic use etc.
- During debeaking to reduce hemorrhage. Intestinal ulcers, scissoring ulcer or any kind of wound for proper healing by collagen synthesis.
- To prevent thin-shelled egg formation in case of excessive environment temperature, supportive treatment in case of coccidiosis for healing of intestinal ulcer.

Dosage

Preventive Dose: 1 gram / 10 litre of drinking water
Treatment Dose: 2 gram / 10 litre of drinking water

**Packing
25 Kg**

HEAD OFFICE

GHAZNAVI BLOCK PLOT #. 29,
COMMERCIAL AREA BAHRIA TOWN, LAHORE
CELL: 0345-8432667, Tel: 042-36448267

SITE OFFICE

55/5-L - 2KM SAHIWAL TO OKARA TOLL PLAZA
MAIN GT ROAD, SAHIWAL. CELL: 0300-9692667

HOT LINE

0303-8847336

info@greenlandzone.com
www.greenlandzone.com