ACADGILD

SESSION 9: Statistical Inference

Assignment 2

Submitted by: Munmun Ghosal
Login Id: munmun55@gmail.com
(M):+91-8007178659

Data Analytics

Table of Contents

1. Problem Statement

a. Calculate the p-value for the test in Problem no 2.
b. How do you test the proportions and compare against hypothetical props? Test hypothesis: proportion of automatic cars is 40%

2. Solution

a. Calculate the p-value for the test in Problem no 2.

The R-script for the given problem is as follows:

```
library(readr)
library(psych)
mtcars <- read_csv("E:/munmun_acadgild/acadgild data analytics/supporting
files/mtcars.csv")
View(mtcars)
mtcars
str(mtcars)
describe(mtcars$am)
table(mtcars$am)

# Calculate the P Value for the test in Problem 2.

t.test(mtcars$am,mu=10,conf.level = 0.95)
t.test(mpg~am,data = mtcars)

# OR

phat <- 13/(13 + 19)
(phat - 0.4)/sqrt(0.4 * 0.6/(13 + 19))

prop.test(13, 13 + 19, p = 0.4, alternative = "less",
        conf.level = 0.95, correct = FALSE)
```

The output of the R-Script (from Console window) is given as follows:

```
> library(readr)
> library(psych)
> mtcars <- read_csv("E:/munmun_acadgild/acadgild data analytics/supporting
files/mtcars.csv")
Parsed with column specification:
cols(
  model = col_character(),
  mpg = col_double(),
```

```
  cyl = col_double(),
  disp = col_double(),
  hp = col_double(),
  drat = col_double(),
  wt = col_double(),
  qsec = col_double(),
  vs = col_double(),
  am = col_double(),
  gear = col_double(),
  carb = col_double()
)
> View(mtcars)
```



Untitled1* | mtcars

Filter

	model	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
1	Mazda RX4	21.0	6	160.0	110	3.90	2.620	16.46	0	1	4	4
2	Mazda RX4 Wag	21.0	6	160.0	110	3.90	2.875	17.02	0	1	4	4
3	Datsun 710	22.8	4	108.0	93	3.85	2.320	18.61	1	1	4	1
4	Hornet 4 Drive	21.4	6	258.0	110	3.08	3.215	19.44	1	0	3	1
5	Hornet Sportabout	18.7	8	360.0	175	3.15	3.440	17.02	0	0	3	2
6	Valiant	18.1	6	225.0	105	2.76	3.460	20.22	1	0	3	1
7	Duster 360	14.3	8	360.0	245	3.21	3.570	15.84	0	0	3	4
8	Merc 240D	24.4	4	146.7	62	3.69	3.190	20.00	1	0	4	2
9	Merc 230	22.8	4	140.8	95	3.92	3.150	22.90	1	0	4	2
10	Merc 280	19.2	6	167.6	123	3.92	3.440	18.30	1	0	4	4
11	Merc 280C	17.8	6	167.6	123	3.92	3.440	18.90	1	0	4	4
12	Merc 450SE	16.4	8	275.8	180	3.07	4.070	17.40	0	0	3	3
13	Merc 450SL	17.3	8	275.8	180	3.07	3.730	17.60	0	0	3	3
14	Merc 450SLC	15.2	8	275.8	180	3.07	3.780	18.00	0	0	3	3
15	Cadillac Fleetwood	10.4	8	472.0	205	2.93	5.250	17.98	0	0	3	4
16	Lincoln Continental	10.4	8	460.0	215	3.00	5.424	17.82	0	0	3	4
17	Chrysler Imperial	14.7	8	440.0	230	3.23	5.345	17.42	0	0	3	4
18	Fiat 128	32.4	4	78.7	66	4.08	2.200	19.47	1	1	4	1
19	Honda Civic	30.4	4	75.7	52	4.93	1.615	18.52	1	1	4	2
20	Toyota Corolla	33.9	4	71.1	65	4.22	1.835	19.90	1	1	4	1
21	Toyota Corona	21.5	4	120.1	97	3.70	2.465	20.01	1	0	3	1
22	Dodge Challenger	15.5	8	318.0	150	2.76	3.520	16.87	0	0	3	2

Showing 1 to 23 of 32 entries

```
> mtcars
# A tibble: 32 x 12
   model              mpg   cyl  disp    hp  drat    wt  qsec    vs    am
gear  carb
   <chr>            <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <dbl>
<dbl> <dbl>
 1 Mazda RX4         21       6  160    110  3.9   2.62  16.5      0     1
4     4
 2 Mazda RX4 Wag     21       6  160    110  3.9   2.88  17.0      0     1
4     4
 3 Datsun 710        22.8     4  108     93  3.85  2.32  18.6      1     1
4     1
 4 Hornet 4 Drive    21.4     6  258    110  3.08  3.22  19.4      1     0
3     1
```

```
 5 Hornet Sportabout  18.7     8  360    175  3.15  3.44  17.0     0     0
3     2
 6 Valiant            18.1     6  225    105  2.76  3.46  20.2     1     0
3     1
 7 Duster 360         14.3     8  360    245  3.21  3.57  15.8     0     0
3     4
 8 Merc 240D          24.4     4  147.    62  3.69  3.19  20       1     0
4     2
 9 Merc 230           22.8     4  141.    95  3.92  3.15  22.9     1     0
4     2
10 Merc 280           19.2     6  168.   123  3.92  3.44  18.3     1     0
4     4
# ... with 22 more rows
> str(mtcars)
Classes ‘spec_tbl_df’, ‘tbl_df’, ‘tbl’ and 'data.frame':     32 obs. of  12
variables:
 $ model: chr  "Mazda RX4" "Mazda RX4 Wag" "Datsun 710" "Hornet 4 Drive" ...
 $ mpg  : num  21 21 22.8 21.4 18.7 18.1 14.3 24.4 22.8 19.2 ...
 $ cyl  : num  6 6 4 6 8 6 8 4 4 6 ...
 $ disp : num  160 160 108 258 360 ...
 $ hp   : num  110 110 93 110 175 105 245 62 95 123 ...
 $ drat : num  3.9 3.9 3.85 3.08 3.15 2.76 3.21 3.69 3.92 3.92 ...
 $ wt   : num  2.62 2.88 2.32 3.21 3.44 ...
 $ qsec : num  16.5 17 18.6 19.4 17 ...
 $ vs   : num  0 0 1 1 0 1 0 1 1 1 ...
 $ am   : num  1 1 1 0 0 0 0 0 0 0 ...
 $ gear : num  4 4 4 3 3 3 3 4 4 4 ...
 $ carb : num  4 4 1 1 2 1 4 2 2 4 ...
 - attr(*, "spec")=
  .. cols(
  ..   model = col_character(),
  ..   mpg = col_double(),
  ..   cyl = col_double(),
  ..   disp = col_double(),
  ..   hp = col_double(),
  ..   drat = col_double(),
  ..   wt = col_double(),
  ..   qsec = col_double(),
  ..   vs = col_double(),
  ..   am = col_double(),
  ..   gear = col_double(),
  ..   carb = col_double()
  .. )
> #summary(mtcars$am)
> describe(mtcars$am)
   vars  n mean  sd median trimmed mad min max range skew kurtosis   se
X1    1 32 0.41 0.5      0    0.38   0   0   1     1 0.36    -1.92 0.09
> table(mtcars$am)

 0  1
19 13
> t.test(mtcars$am,mu=10,conf.level = 0.95)

        One Sample t-test

data:  mtcars$am
```

```
t = -108.76, df = 31, p-value < 2.2e-16
alternative hypothesis: true mean is not equal to 10
95 percent confidence interval:
 0.2263446 0.5861554
sample estimates:
mean of x
  0.40625

> t.test(mpg~am,data = mtcars)

        Welch Two Sample t-test

data:  mpg by am
t = -3.7671, df = 18.332, p-value = 0.001374
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 -11.280194  -3.209684
sample estimates:
mean in group 0 mean in group 1
       17.14737        24.39231

>
> # OR
>
> phat <- 13/(13 + 19)
> (phat - 0.4)/sqrt(0.4 * 0.6/(13 + 19))
[1] 0.07216878
>
>
> prop.test(13, 13 + 19, p = 0.4, alternative = "less",
+           conf.level = 0.95, correct = FALSE)

        1-sample proportions test without continuity correction

data:  13 out of 13 + 19, null probability 0.4
X-squared = 0.0052083, df = 1, p-value = 0.5288
alternative hypothesis: true p is less than 0.4
95 percent confidence interval:
 0.0000000 0.5508812
sample estimates:
      p
0.40625
```

b. How do you test the proportions and compare against hypothetical props?

Test hypothesis: proportion of automatic cars is 40%

The R-script for the given problem is as follows:

```
prop.test(13, 32, p = 0.4, alternative = "less",
        conf.level = 0.95, correct = FALSE)
#OR
```

prop.test(table(mtcars$am)[2],nrow(mtcars),p=0.4,alternative = "less",conf.level = 0.95,correct=FALSE)

The output of the R-Script (from Console window) is given as follows:

```
> prop.test(13, 32, p = 0.4, alternative = "less",
+           conf.level = 0.95, correct = FALSE)

        1-sample proportions test without continuity correction

data:  13 out of 32, null probability 0.4
X-squared = 0.0052083, df = 1, p-value = 0.5288
alternative hypothesis: true p is less than 0.4
95 percent confidence interval:
 0.0000000 0.5508812
sample estimates:
      p
0.40625

> #OR
>
> prop.test(table(mtcars$am)[2],nrow(mtcars),p=0.4,alternative =
"less",conf.level = 0.95,correct=FALSE)

        1-sample proportions test without continuity correction

data:  table(mtcars$am)[2] out of nrow(mtcars), null probability 0.4
X-squared = 0.0052083, df = 1, p-value = 0.5288
alternative hypothesis: true p is less than 0.4
95 percent confidence interval:
 0.0000000 0.5508812
sample estimates:
      p
0.40625
```

Conclusion/Interpretation:

Test Hypothesis: proportion of automatic cars is 40%.
At confidence level of 0.95, since p- value is greater than alpha, we fail to reject the null hypothesis