
MEAT & CANCER: A CRITICAL REVIEW BASED ON CAUSAL ANALYSIS (*work in progress*)

A PREPRINT

Any volunteer?*
Any Department
Anywhere

Enrique Otero
Madrid, Spain
@eoteromuras
eoteromuras@gmail.com

August 22, 2019

ABSTRACT

According to World Health Organization (WHO), processed meat has been declared Group 1 carcinogenic to humans. That means that according to epidemiological studies there is a convincing evidence that the agent causes cancer. However, reviewing some of the mainly referred studies with the lenses of causal inference analysis reveals possible flaws that would invalidate these conclusions. The author(s) intention is to discuss these studies with statistical rigor. By applying last accepted knowledge in the field of causal inference as diagrams and *do-calculus*. With main focus on transparent exposition of health domain assumptions. And the translation of these assumptions into explicit language, diagrams and formulas. So the veracity of domain assumptions can be refuted according to domain expertise. And any conclusion derived from these assumptions being validated or invalidated on the bases of axiomatic logic and maths.

Keywords Meat · Cancer · Causality

1 Introduction

In October 2015 IARC held an expert panel that considered the evidence for red and processed meats as possible human carcinogens. They classified processed meat as a Group 1 carcinogenic to humans, and red meat as Group 2A, probably carcinogenic [1]. A summary of the final evaluations were published online in The Lancet Oncology [2]. And the details of these conclusions were published later in a monograph in 2016 [3].

The consumption of processed meat was associated with small increases in the risk of cancer in the studies reviewed. In these studies, the risk generally increased with the amount of meat consumed [1].

In the next sections we will focus on three studies supporting IARC conclusions, and we will remark different flaws detected in them. Particularly:

- In Section 2, as starting point we will base on IARC monograph and Chan meta-analysis [4], as it's the main reference for IARC when they conclude "each 50 gram portion of processed meat eaten daily increases the risk of colorectal cancer by 18%". And we will remark potential problems related to heterogeneity. Though our focus will be in introducing possible problems related to not conditioning on missing confounders (Section 3), or conditioning on a collider (Section 4).
- In Section 3, as example of discarding a possible confounder based on a possible wrong procedure we present Sandhu et al meta-analysis [7], both referenced by Chan's [4] and IARC monograph [3].

*Use footnote for providing further information about author (webpage, alternative address)—*not* for acknowledging funding agencies.

Table 1: Relative risks of meta-analyses of processed meat, and colorectal cancer. Chan et al. meta-analysis

Pooled RR (95% CI)	n	Heterogeneity (I ²)
1.18 (1.10–1.28), P-value=0.00	9	12%, P-value=0.33

Table 2: *

RR – relative risk; CI – confidence interval; n – number of studies

- Finally in Section 4, as an example of generating strange and wrong conclusions based on the wrong procedure of conditioning on a collider, we'll discuss a study by Cross et al [12]. This study is particularly relevant as it's the one that contributes the most to results on Chan's meta-analysis.

For this purposes we will use different causal inference techniques as causal diagrams and *do-calculus*, as presented by Pearl's et al [5].

2 Processed Meat and Colorectal Cancer Incidence

In [3] the IARC Working Group analyzed both "20 large [...] cohort studies [...] extended from as early as the 1990s until the 2010s", and "a large number of case-control studies (approximately 150)". In the monograph authors describe also five criteria they applied in reviewing and interpreting the available literature in order to be considered for their meta-analysis. One of these criteria, which we will focus specifically on this paper is the "Adjustment for potential confounding factors"

Regarding case-control studies, they considered that "approximately 10% of all case-control studies reviewed were informative for the assessment of the consumption of processed meat in relation to incidence of cancer of the colorectum". Taking into account previous statement of approximately 150 control-studies, they should be $150/10 = 15$ informative studies. However they say: "Six of the nine studies considered showed positive associations with cancer of the colorectum." Could this difference (9 vs. 15) be a typo?

In relation to cohort studies, authors presented conclusions from a meta-analysis including data from 10 of these studies that "reported a statistically significant dose-response association between consumption of red meat and/or processed meat and cancer of the colorectum". More concretely, they refer to Chan et al [4], where "dose-response relationships were expressed per increment of intake of 100 grams per day for red and processed meat, and 50 grams per day for processed meat as in previous meta-analyses [6], [7]

2.1 Dose-response Analysis on Processed Meat. Heterogeneity and Confounders

In detail, in Chan's meta-analysis [4] 26 publications from 21 studies were included. Being 15 publications from 14 studies on processed meat. Results in Table 1.

Regarding heterogeneity, though I-squared level of 12% may seem no significant, specially with a p-value of 0.33, criticism has been done to I-squared as an adequate measure of heterogeneity, specially when the number of studies is small. [8].

In relation to confounders, authors claim: "we cannot rule out residual confounding". Though "In all studies, relative risk estimates were adjusted for age and sex, and all except two adjusted for total energy intake. More than half of the study results were adjusted for body mass index (BMI), smoking, alcohol consumption, or physical activity, close to half controlled for dairy food or calcium intake, social economic status, family history of colorectal cancer, or plant food or folate intake. In some studies, the estimates were controlled for use of nonsteroidal anti-inflammatory drugs, fish or white meat intake". Anyway, they also say: "several potential confounders were not included in the final statistical models in some studies because, as the authors reported, their inclusion in the model did not substantially modified the relative risk estimates." However, sometimes the decision of exclude a confounder could be flawed, not only based on domain assumptions, but also in the logical procedure itself, as we will discuss on the example of Section 3

Authors also refer to other studies with similar conclusions, like [6, 10] Though after their conclusions they also say: "In a more recent article on the NHS and the HPFS, the associations of red meat and processed meat and colon cancer were attenuated after better adjustment for confounders and longer followup [9]

The biggest study analyzed in [4] regarding number of people was Cross et al. [12] with 494036 men and women. In this study adjusts were made on "Age, sex, ethnicity, BMI, smoking habits, alcohol intake, physical activity, total energy

intake, fruit and vegetable intake, education level, marital status, family history of cancer" And according to weight and results Cross's study is the one that contributes the most to RR for colorectal cancer on the consumption of processed meat on Chan's meta-analysis. So in the next section 4 we'll try to proof that Cross study could be flawed because of a wrong experiment design on implicitly conditioning on a collider.

3 Don't Discard Fiber, Vegetables, Fruits or Life-style as Possible Confounders


In 2001 Sandhu et al published a meta-analysis on the relation between meat consumption and colorectal cancer [7]. Discussing about "Meat and Other Dietary and Associated Factors" they wrote: "the current prospective epidemiological data show only a weak negative association between vegetables and fruits consumption and risk of colorectal cancer. Four recent studies, two randomized trials on adenoma recurrence and two large prospective studies on colorectal cancer found no association among fiber, vegetables, and fruits consumption and risk of colorectal cancer."

But even if the total effect from fiber, vegetables, and fruits consumption on colorectal cancer is negligible, mathematically the direct effect even could be important. Under the assumption that consumption of vegetables was associated with consumption of meat via dietary lifestyle.

As an example, considering:

- X: meat
- Y: colorectal cancer
- V: vegetables
- U: some dietary lifestyle, confounder of V and X

with the following causal diagram:



If "no association among fiber, vegetables, and fruits consumption and risk of colorectal cancer"

$$P(Y|do(V)) = P(Y) \quad (1)$$

Otherwise, according to the provided causal diagram, V blocks all back-door paths between X and Y. Thus:

$$P(Y|do(X)) = \sum_{i=1}^N P(Y|X, V_i)P(V_i) \neq P(Y|X) \quad (2)$$

On the basis of the previous analysis we still have no evidences to ignore direct effect of fiber, vegetables and fruits consumption with dietary lifestyle as possible confounder.

Other of the biggest studies like English et al [11] included also fish as possible cause.

4 Leukemia versus Life-style Cancers. The Collider Bias

As previously said, Cross study on 2007 [12] was the second biggest included on Chan's meta-analysis. And the one that contributed more to results on relative risk for processed meat on colorectal cancer. Ignoring this study could reduce Chan's aggregated result from 1.18 to 1.10. Value that was included in the IARC monograph, and into the WHO claim. In this context, we will show in this section an example of bad control in this study leading to some strange results.

Extracted from Cross study: "Surprisingly, both leukemia and melanoma were inversely associated with processed meat intake; the inverse association for leukemia was mainly for lymphocytic leukemia (n = 534; HR = 0.70; 95% CI = 0.52–0.93; p for trend = 0.05) and not myeloid and monocytic leukemia (n = 457; HR = 0.88; 95% CI = 0.64–1.20; p for trend = 0.73). The associations between processed meat intake and cancer risk are summarized in Figure 2, in order of risk magnitude."

But according to the described cohort follow-up and case ascertainment: "these analyses was calculated from baseline (1995–1996) until censoring at the end of 2003, or when the participant moved out of one of the eight study areas, had a





Figure 1: HRs and 95% CIs for the 5th Versus 1st Quintile of Processed Meat Intake and Cancer Risk for Both Sexes Combined (Except for Sex-Specific Cancers). From Cross et al. [12]

cancer diagnosis, or died, whichever came first" [12]. Thus, analyzing several types of cancer as target while implicitly conditioning on not having any other cancer converts the variable "any cancer" in a collider.

Considering

- X: processed meat
- Y: leukemia
- C1: colorectal cancer
- C: any type of cancer

and assuming the following causal diagram:



So conditioning on the collider C opens the path $Y \rightarrow C \leftarrow C1 \leftarrow X$, invalidating the results of the study. This situation is typically known as *collider bias*. Being a wellknown example the *Berkson Paradox* described for instance in Pearl's et al [5].

And this could explain these strange results where cancers usually associated with dietary lifestyle as colorectal have relative risk ratios above 1 (HR=1.2), whereas leukemia shows apparently an inverse relation to processed meat, with an HR of 0.7. As we can see in the 1

To be continued...

5 Conclusions

To be finished...

References

- [1] Q&A on the carcinogenicity of the consumption of red meat and processed meat <https://www.who.int/features/qa/cancer-red-meat/en/> October 2015
- [2] Bouvard V. et al. Carcinogenicity of consumption of red and processed meat
- [3] IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Red Meat and Processed Meat. Vol 114 <https://monographs.iarc.fr/wp-content/uploads/2018/06/mono114.pdf>
- [4] Chan et al. Red and Processed Meat and Colorectal Cancer Incidence: Meta-Analysis of Prospective Studies <https://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0020456&type=printable>
- [5] Pearl et al. The Book of Why. The New Science of Cause and Effect
- [6] World Cancer Research Fund/American Institute for Cancer Research. (2007) Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective Washington DC: AICR.
- [7] Sandhu MS, White IR, McPherson K (2001) Systematic review of the prospective cohort studies on meat consumption and colorectal cancer risk: a meta-analytical approach. *Cancer Epidemiol Biomarkers Prev* 10: 439–446
- [8] Von Hippel P. The heterogeneity statistic I² can be biased in small meta-analyses <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4410499/> *BMC Med Res Methodol.* 2015; 15: 35.
- [9] Wei EK, Giovannucci E, Wu K, Rosner B, Fuchs CS, et al. (2004) Comparison of risk factors for colon and rectal cancer. *IntJCancer* 108: 433–442.
- [10] Wei EK, Colditz GA, Giovannucci EL, Fuchs CS, Rosner BA (2009) Cumulative risk of colon cancer up to age 70 years by risk factor status using data from the Nurses' Health Study. *AmJEpidemiol* 170: 863–872.
- [11] English DR, MacInnis RJ, Hodge AM, Hopper JL, Haydon AM, et al. (2004) Red meat, chicken, and fish consumption and risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev* 13: 1509–1514.
- [12] Cross et al. 2007 A prospective study of red and processed meat intake in relation to cancer risk.