

Student Information

Full Name : Murat Bolu

Id Number : 2521300

Answer 1

Basis: For $n = 1$, $6^{2n} - 1 = 6^2 - 1 = 35$, and $5 \mid 35$, and $7 \mid 35$.

Inductive Step: Assuming that $6^{2n} - 1$ is divisible by both 5 and 7, one can show that $6^{2(n+1)} - 1 = 6^{2n+2} - 1$ is divisible by 5 and 7 for $n \in \{1, 2, 3, \dots\}$. Since $5 \mid 6^{2n} - 1$ and $7 \mid 6^{2n} - 1$, that means $35 \mid 6^{2n} - 1$ which is shown below.

$$\begin{aligned} 6^{2n} - 1 &= 5 \cdot x && \text{for some } x \in \mathbb{N} \\ 6^{2n} - 1 &= 7 \cdot y && \text{for some } y \in \mathbb{N} \\ 5 \cdot x &= 7 \cdot y \end{aligned}$$

Since x and y are natural numbers, x must be divisible by 7 and y must be divisible by 5, and we can rewrite them as $x = 7 \cdot k$ and $y = 5 \cdot l$. The other possibility is that they are both zero, which is obviously not possible since $6^{2n} - 1$ is at least 35.

$$\begin{aligned} 6^{2n} - 1 &= 5 \cdot 7 \cdot k && \text{for some } k \in \mathbb{N} \\ 6^{2n} - 1 &= 7 \cdot 5 \cdot l && \text{for some } l \in \mathbb{N} \\ 6^{2n} - 1 &= 35 \cdot k \\ 6^{2n} - 1 &= 35 \cdot l \end{aligned}$$

There exists k or l such that they are natural numbers, and that means $6^{2n} - 1$ is divisible by 35.

$$\begin{aligned} 6^{2n} - 1 &\equiv 0 && (\text{mod } 35) \\ 6^{2n} &\equiv 1 && (\text{mod } 35) \\ 6^{2n+2} &\equiv 36 && (\text{mod } 35) \\ 6^{2n+2} &\equiv 1 && (\text{mod } 35) \\ 6^{2n+2} - 1 &\equiv 0 && (\text{mod } 35) \end{aligned}$$

Therefore, there exists a natural number $a \in \mathbb{N}$ such that $6^{2n+2} - 1 = 35 \cdot a$. It is obvious that $6^{2n+2} - 1$ is divisible by both 5 and 7 since $5 \mid 35a$ and $7 \mid 35a$.

■

Answer 2

Basis:

- For $n = 0$, $H_0 = 1 \leq 1 = 9^0$.
- For $n = 1$, $H_1 = 5 \leq 9 = 9^1$.
- For $n = 2$, $H_2 = 7 \leq 81 = 9^2$.
- For $n = 3$, $H_3 = 8H_2 + 8H_1 + 9H_0 = 8 \cdot 7 + 8 \cdot 5 + 9 \cdot 1 = 105 \leq 729 = 9^3$.

Inductive Step: Assuming that $H_n \leq 9^n$, $\forall n \leq k$, one can show that H_{k+1} is less than or equal to 9^{k+1} . We can replace every H_n with something greater than or equal to itself.

$$H_{k+1} = 8H_k + 8H_{k-1} + 9H_{k-2}$$

$$H_{k+1} \leq 8 \cdot 9^k + 8H_{k-1} + 9H_{k-2}$$

$$H_{k+1} \leq 8 \cdot 9^k + 8 \cdot 9^{k-1} + 9H_{k-2}$$

$$H_{k+1} \leq 8 \cdot 9^k + 8 \cdot 9^{k-1} + 9 \cdot 9^{k-2}$$

$$H_{k+1} \leq 8 \cdot 9^k + 8 \cdot 9^{k-1} + 9^{k-1}$$

$$H_{k+1} \leq 8 \cdot 9^k + 9 \cdot 9^{k-1}$$

$$H_{k+1} \leq 8 \cdot 9^k + 9^k$$

$$H_{k+1} \leq 9 \cdot 9^k$$

$$H_{k+1} \leq 9^{k+1}$$

■

Answer 3

Answer 4

Answer 5

a)

b)

c)