# **Student Information**

Full Name : Murat Bolu Id Number : 2521300

## Answer 1

The graph G can be represented by a pair (V, E) where V is the set of vertices, and E is the set of edges.  $V = \{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}\}$  and  $E = \{\{\mathbf{a}, \mathbf{b}\}, \{\mathbf{a}, \mathbf{c}\}, \{\mathbf{b}, \mathbf{c}\}, \{\mathbf{b}, \mathbf{e}\}, \{\mathbf{c}, \mathbf{d}\}, \{\mathbf{d}, \mathbf{e}\}\}$ . Therefore, |V| = 5, and |E| = 7.

- a)  $\sum_{v \in V} \deg(v) = 2 \cdot |E| = 14$  by handshaking lemma, since all edges increase the degree of vertices they connect by one. The sum of degrees of all vertices is twice the number of edges since all edges are counted twice.
- **b)** Adjacency matrix is  $M = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 \end{bmatrix}$  where the item  $M_{ij}$  represents the existence of an

edge between the nodes i and j. The vertices are enumerated in the order they are written above. There are 14 non-zero entries in the adjacency matrix representation of G.

**c)** Incidence matrix is  $M = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 \end{bmatrix}$  where the item  $M_{ij}$  represents the incidence

of the vertex i with the edge j. The rows represent vertices and columns represent edges. The edges are enumerated in the order they are written above. There are 21 zero entries in the incidence matrix representation of G.

- d) There is not, since  $\mathbf{c}$  and  $\mathbf{e}$  are not connected, only one can be chosen, and only two choices are  $\{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}\}$  and  $\{\mathbf{a}, \mathbf{b}, \mathbf{d}, \mathbf{e}\}$ , both of which are obviously not connected since  $\mathbf{d}$  is not connected to  $\mathbf{b}$ .
- e) No, it is not, since  $\{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$  are all connected, it is not possible to place them in two disjoint and independent sets such that there is no edge connecting between two elements in the same set. One can exhaust all possibilities by trying  $\{\{\mathbf{a}, \mathbf{b}\}, \{\mathbf{c}\}\}, \{\{\mathbf{a}, \mathbf{c}\}, \{\mathbf{b}\}\}, \{\{\mathbf{b}, \mathbf{c}\}, \{\mathbf{a}\}\}$ . Since a subgraph of G is not bipartite, adding new nodes to that subgraph would not be able to fix that and therefore G is not bipartite.
- f) For every undirected edge, we can choose two directions, and therefore we have  $2^{|E|} = 2^7 = 128$  different directed graphs that have G as their underlying graph.

- g) The vertices {a, b, c, e} have odd degrees, therefore we cannot visit all edges, i.e. create a Euler path, since when we enter a vertex, we have to leave it. The only exception to this rule is for the first and the last vertices, which there are two of. A simple path can be represented by the sequence of vertices such as (a, b, c, d, e, a, c) in order. The path contains 6 edges in total. Therefore, the longest simple path is of length 6.
- h) There is only one, and it is the graph itself, since all vertices are connected, i.e. there exists a path between every pair of vertices. The are  $\binom{5}{2} = 10$  pairs, and the path between each pair is as follows:  $\{(\mathbf{a}, \mathbf{b}), (\mathbf{a}, \mathbf{c}), (\mathbf{a}, \mathbf{c}, \mathbf{d}), (\mathbf{a}, \mathbf{e}), (\mathbf{b}, \mathbf{c}), (\mathbf{b}, \mathbf{c}, \mathbf{d}), (\mathbf{b}, \mathbf{e}), (\mathbf{c}, \mathbf{d}), (\mathbf{c}, \mathbf{d}, \mathbf{e}), (\mathbf{d}, \mathbf{e})\}$ .
- i) Since the graph does not have an Euler path as explained in item j), it does not have an Euler circuit since Euler circuits are a subset of Euler paths.
- j) The graph G does not have an Euler path, since there are 4 vertices with odd degrees,  $\{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{e}\}$ . There must be either no vertices with odd degrees or exactly two vertices with an odd degree to create an Euler path.
- k) Yes, it has many, one example being (a, b, c, d, e, a).
- 1) Yes, it has many, one example being  $(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e})$ .

## Answer 2

#### Answer 3

## Answer 4