

# Your Lady-Microbiome



*Health begins with us.*



Sample-ID  
Sample receipt

## Table of contents

Summary of results	3
What are bacteria doing in my vagina?	4
MICROBIAL COMPOSITION	5
Diversity of the vaginal microbiome	5
Dysbiosis of the vaginal microbiome	6
HEALTH	7
Is there a connection between the vaginal microbiome and endometriosis?	7
How does the vaginal microbiome affect female fertility?	8
How does the microbiome change with HPV infections?	9
Does my microbiome promote the growth of fungi in the vagina?	10
RECOMMENDATIONS	10
How can I support the health of my vagina?	11-12
What is recommended for suspected endometriosis or for those already diagnosed with endometriosis to relieve symptoms?	13
How can fertility be promoted?	14-15
What is recommended if there is suspicion of an HPV infection?	16
What can I do about a yeast infection and prevent it in the future?	17
Literature	18
Notes	19
Miscellaneous	19

### **Bacterial Composition**

Diversity of the microbiome (Shannon-Index)

Balance of the microbiome (Dysbiosis-Index)

### **Health**

Endometriosis

Fertility

HPV Infection

### **Fungal Profile**

Fungal Infection

## What are bacteria doing in my vagina?

The vaginal microbiome consists of a variety of bacteria, viruses, fungi, and other microorganisms that reside in the vagina\*. This community of microbes, particularly lactic acid bacteria, protects the mucous membranes and prevents the spread of harmful bacteria <sup>[1]</sup>.


While microorganisms coexist with us humans for mutual benefit on and in our bodies, the microbiome only becomes noticeable in cases of health issues. Many of these microorganisms, especially bacteria, promote a healthy balance of the vaginal mucosa, while some „bad“ bacteria can lead to discomfort <sup>[2]</sup>.

The composition of the bacterial community can vary from woman to woman and depends on various factors such as hormones, environment, diet, and hygiene habits. A balanced microbiome contributes to protective mucous membranes by inhibiting the spread of potentially harmful microorganisms. Just like in other parts of the body, microorganisms live in a balance that can be disrupted by external and internal influences <sup>[2][3]</sup>.

While research is still in its early stages, some studies suggest that targeted support of the vaginal microbiome may play a significant role in a healthy pregnancy and the protection of internal reproductive organs in the future. Additionally, a disrupted microbiome can affect various health issues, such as the transmission of infectious diseases. In the future, personalized prevention approaches may help protect and strengthen the bacterial community <sup>[3]</sup>.

The vaginal mycobiome is a component of the vaginal microbiome composed of fungi. In addition to the bacterial microbiome, the mycobiome also plays a crucial role in the health and microbial balance of the vagina. [33][34]

In the following, we provide you with insights into the composition of the microorganisms in your microbiome. Over 2900 groups of bacteria in your microbiome are analyzed for this purpose. We also provide recommendations for improving vaginal health to support women's health.“



\* A muscular hollow organ that connects the uterus to the vulva, which is the externally visible part of the female genitalia.

## MICROBIAL COMPOSITION

### Diversity of the vaginal microbiome

The diversity of the microbiome is an important indicator of its health. In most cases (e.g., in the gut), higher diversity is associated with a stable microbiome. This is because a greater variety of bacteria can help maintain the balance of the microbiome and limit the growth of harmful bacteria. However, the vaginal microbiome is primarily composed of lactic acid bacteria, so diversity in the vagina is naturally lower than in other parts of the body.


The Shannon Index is used to assess diversity and provides information about the bacterial diversity in your vaginal microbiome. A high value indicates that the microbiome community is relatively diverse. In the case of the vaginal microbiome, this often suggests a decrease in beneficial lactic acid bacteria.

## MICROBIAL COMPOSITION

### Dysbiosis of the vaginal microbiome

The Dysbiosis Index is another way to assess the balance of the microbiome. Dysbiosis refers to an imbalance in the bacterial groups in the vagina. The microbiome is compared to the average composition of reference individuals to determine dysbiosis <sup>[1]</sup>.


A low Dysbiosis Index value (green range) indicates a balanced microbiome, while high values (red range) are associated with dysbiosis. Vaginal microbiome dysbiosis is not a disease but increases the risk of bacterial vaginosis, where harmful bacteria that typically do not thrive in the vagina can overpopulate <sup>[2]</sup>.



### Is there a connection between the vaginal microbiome and endometriosis?

Endometriosis is a painful and complex condition in which tissue that normally lines the uterus grows outside the uterus. This abnormal growth can lead to inflammation, severe menstrual pain, pain during intercourse, and other severe symptoms, including infertility. Estimates suggest that about 10% of women of childbearing age are affected by endometriosis. While the exact cause of endometriosis is not fully understood, there is evidence to suggest that the vaginal microbiome could play a role in this condition <sup>[4]</sup>.

Studies have shown that the bacterial community in the vagina of women with endometriosis may differ in composition and diversity compared to healthy women. It is suspected that these changes may exacerbate inflammatory reactions in the pelvic area and contribute to the development or worsening of endometriosis, including the growth and spread of uterine tissue. Below, we provide an overview of the bacteria that are present in altered amounts in the vaginal microbiome of women with endometriosis. These bacteria include *Gemella* and *Atopobium* <sup>[5][6]</sup>.




## How does the vaginal microbiome affect female fertility?

Fertility, the ability to conceive or receive offspring, is an important aspect of women's reproductive health and a central theme in family planning. While many couples have no fertility issues, fertility problems can be a distressing challenge for others. Common causes of fertility problems include hormonal disorders, often associated with irregular or absent periods and issues with the reproductive organs, such as endometriosis. Male factors, such as reduced sperm quality or quantity, can also lead to infertility. Age also plays a role, as fertility in women tends to decrease after the age of 30 <sup>[7][8]</sup>.

In recent years, research has begun to investigate the potential link between vaginal bacteria and fertility more closely. The microbiome plays an important role in maintaining vaginal health. It is believed that a healthy vaginal microbiome increases the likelihood of successful implantation of a fertilized egg in the uterus. A balanced community of bacteria provides protection against infections that could affect the reproductive organs or implantation. Additionally, early studies suggest that the vaginal microbiome and female hormones may influence each other <sup>[8][9][10]</sup>.

Below, we provide an overview of the bacteria associated with infertility in women. These bacteria include *Staphylococcus* and *Enterococcus*, among others <sup>[8][10]</sup>.





## How does the microbiome change with HPV infections?

Human papillomaviruses (HPV) are a group of viruses that can infect the skin and mucous membranes, including the genital areas of both men and women. HPV infections are widespread worldwide, with most people becoming infected with HPV at least once in their lifetime. In women, HPV infections are particularly relevant as some strains of the virus are associated with various health issues, including cervical cancer <sup>[1][12]</sup>.

HPV is highly contagious and is mainly transmitted through sexual contact. There are many different strains of HPV, some of which are harmless and do not cause symptoms. Other strains can cause genital warts, while still others are considered high-risk strains because they can increase the risk of cervical cancer <sup>[11]</sup>. HPV infections, especially by high-risk strains like HPV-16 and HPV-18, can increase the risk of cervical cancer. This occurs when the virus penetrates the cells of the cervix and causes changes in the cells that can ultimately lead to cancer <sup>[13]</sup>.

It is increasingly being researched whether the vaginal microbiome could play a role in HPV infections. Some studies have suggested that women with a balanced microbiome may have a lower risk of HPV infections. The beneficial bacteria in the vagina could strengthen the vagina's defense mechanisms and make it more difficult for the virus to penetrate. Additionally, it is possible that an HPV infection itself may influence the bacterial community in the vagina <sup>[1][2][11]</sup>.

Below, we provide an overview of the bacteria associated with HPV infections. It has already been shown that, among others, *Prevotella* and *Sneathia* are elevated in women with HPV infections <sup>[12][13][14]</sup>.

## Does my microbiome promote the growth of fungi in the vagina?

*Candida albicans* is the most common fungus found in the vagina and is a natural component of the vaginal microbiome. Normally, *Candida* does not cause issues. However, under certain circumstances such as hormonal changes during pregnancy, breastfeeding, or menopause, antibiotic use, or a weakened immune system, *Candida* can overgrow and lead to a fungal infection.

A fungal infection in the vagina, also known as vaginal yeast infection or vaginal candidiasis, can cause unpleasant symptoms. The most common symptoms include itching in the vaginal area and a burning sensation, especially during urination or intercourse. Additionally, the vaginal discharge may be altered, often becoming thick and white during an infection.

Some women experience recurrent yeast infections, which may also indicate other health issues such as diabetes and a weakened immune system. In pregnant women, fungal infections can lead to complications. Untreated fungal infections also increase the risk of sexually transmitted infections. Women with diabetes or a weakened immune system are at a higher risk of more severe fungal infections. <sup>[33][34]</sup>

Below, we provide an overview of the most common fungi in the vaginal microbiome and whether they occur in elevated amounts in your case. <sup>[35][36]</sup>

### Your Result:

*Candida albicans*

*Candida glabrata*

*Candida krusei*

*Candida parapsilosis*

*Candida tropicalis*

A result in the green range indicates that no pathogenic fungi were found in elevated amounts, while results in the red range mean that elevated levels of the respective fungus are present in your vagina. If your result is not in the green range, it should be clarified by a doctor.

## RECOMMENDATIONS

### How can I support the health of my vagina?

A healthy microbiome plays a crucial role in maintaining vaginal health and protecting against infections by creating an acidic environment in the genital area that restricts the growth of harmful microorganisms <sup>[1]</sup>.

Vaginal health can be promoted through a healthy lifestyle and specific measures to maintain it. A balanced diet rich in fiber, fruits, and vegetables can help promote a healthy microbiome. Hormones play a significant role in regulating the vaginal microbiome <sup>[15][16]</sup>. During the menstrual cycle, the bacterial composition naturally undergoes changes. Estrogen levels rise before ovulation, promoting the growth of lactic acid bacteria, the „good“ bacteria that maintain the acidic environment. After menopause, estrogen levels decrease, which can lead to changes in the vaginal microbiome <sup>[17][18]</sup>. Some studies suggest that hormones in meat products may have an impact on the vaginal microbiome. In particular, hormones used in livestock farming to promote growth could potentially enter meat and affect hormonal balance in the body <sup>[19]</sup>. This could potentially result in changes in bacterial composition. Choosing meat products from organic or hormone-free sources can minimize the hormonal influence on the vaginal microbiome. Antibiotics can disrupt the microbiome, so they should only be taken as prescribed by a doctor, in the full prescribed dose <sup>[20]</sup>.

Aggressive intimate care products can disrupt the balance. Therefore, it is preferable to use mild, pH-neutral soaps. Using a condom during sexual intercourse reduces the risk of sexually transmitted infections that can negatively affect the vaginal microbiome. Regular check-ups are essential to monitor the health of your vagina and vaginal microbiome <sup>[21]</sup>. Additionally, avoiding chronic stress is important, as it weakens the immune system and can thus influence the vaginal microbiome.



## RECOMMENDATIONS

### How can I support the health of my vagina?

Furthermore, choosing the right underwear can support vaginal health. Synthetic fabrics like nylon, polyester, and spandex are widespread in many types of underwear. These materials are often chosen for their durability and low cost but are not recommended for a healthy vagina<sup>[22]</sup>. The vaginal environment is naturally moist and warm and benefits from good airflow. Synthetic fabrics tend to trap moisture and reduce airflow, creating an ideal environment for bacterial growth. This can lead to an imbalance in the vaginal microbiome and promote infections, including bacterial vaginosis. There is evidence that wearing synthetic underwear, especially in women already prone to vaginal infections, can increase the risk of bacterial vaginosis. Women who experience recurrent infections may consider switching to underwear made of natural materials like cotton, as these can contribute to better ventilation and moisture regulation. Additionally, especially in cases of increased susceptibility to vaginal infections, it is important to avoid wearing thongs and G-strings, as these types of underwear can transfer bacteria from the anal area to the vaginal area. Therefore, it is also important to wipe from front to back after using the toilet to prevent the spread of intestinal bacteria to the vagina, especially if there is already an increased susceptibility to bacterial vaginosis. In addition, the thin straps of thongs and G-strings can cause friction and irritation during movement. These irritations can compromise the vaginal barrier and facilitate the penetration of harmful bacteria.



## RECOMMENDATIONS

### What is recommended for suspected endometriosis or for those already diagnosed with endometriosis to relieve symptoms?

If you suspect endometriosis, especially due to severe menstrual pain, you should consult a specialist in gynecology who specializes in the diagnosis and treatment of endometriosis. It is helpful for the treating physician to note the symptoms, their duration, and their intensity in advance <sup>[4][5]</sup>. Imaging techniques such as ultrasound or magnetic resonance imaging (MRI) can identify anomalies in the pelvic area indicative of endometriosis. The definitive diagnosis of endometriosis often requires a laparoscopy, a minimally invasive surgery <sup>[23]</sup>. During this procedure, the doctor can take tissue samples and assess the extent of the disease. Treatment options following a diagnosis include pain medication and hormonal therapies. Hormonal preparations can be used to regulate menstrual cycles and reduce the growth activity of endometriosis tissue. This can alleviate pain and inflammation. In some cases, surgery may be required to remove the endometriosis tissue. This can often be done minimally invasively and depends on the severity of endometriosis and other individual factors.

A healthy diet rich in fruits, vegetables, whole grains, and lean protein sources can reduce inflammation in the body. Some women experience an improvement in their symptoms by avoiding foods that may promote inflammation, especially sugary and processed foods <sup>[24]</sup>. Simultaneously, anti-inflammatory foods such as fatty fish (e.g., salmon), berries, leafy greens, nuts, and seeds can support symptom relief. Some studies suggest that the consumption of red meat, especially processed meat, may be associated with a higher risk of endometriosis. It is advisable to reduce red meat consumption or switch to lean meat and plant-based protein sources. Stress can worsen the symptoms of endometriosis, so stress management techniques such as regular exercise, adequate sleep, and relaxation exercises can help reduce stress.

## RECOMMENDATIONS

### How can fertility be promoted?

If there is suspicion of infertility due to symptoms during the menstrual cycle or difficulties in conceiving, a doctor who specializes in fertility should be consulted. Through various examinations, fertility and potential issues can be determined. These examinations include blood tests to check hormone levels, ultrasound examinations to assess the ovaries and uterus, and semen analyses for the partner to assess sperm quality. Fertility problems can often be treated with professional assistance, thanks to the current state of research. Various treatment options include hormonal therapies and in vitro fertilization (IVF) <sup>[25]</sup>.

Fertility can be promoted through a combination of a healthy lifestyle, a balanced diet, and stress-reducing measures. In addition to an adequate intake of fruits, vegetables, whole grains, low-fat protein sources, and unsaturated fats in the diet, folic acid, iron, vitamin D, and omega-3 fatty acids are particularly important for fertility <sup>[26]</sup>. If necessary, it is advisable to take these nutrients as supplements in consultation with a doctor. Maintaining a healthy weight within the normal range is important, as overweight can affect fertility <sup>[27]</sup>. It is recommended to avoid or significantly reduce alcohol consumption and quit smoking, as excessive alcohol consumption and smoking can negatively affect fertility. Additionally, stress should be reduced, as chronic stress can negatively impact fertility. Regular physical activity is important for a healthy lifestyle. However, excessive exercise may have the opposite effect under certain circumstances <sup>[28]</sup>.

The fertile phase in a woman's menstrual cycle can vary, but it typically falls in the middle of the cycle. The average menstrual cycle is about 28 days, with the first day of menstruation considered as day 1. In a 28-day cycle, ovulation usually occurs around day 14. Sperm can survive in the female reproductive tract for up to five days, and the egg is fertile for about 12-24 hours after ovulation <sup>[29]</sup>. Therefore, a period of about 6 days is often considered the fertile phase - the 5 days before ovulation and the day of ovulation itself. Therefore, timing intercourse correctly on a woman's most fertile days is important to increase the chances of pregnancy.



## RECOMMENDATIONS

### How can fertility be promoted?

Tracking the menstrual cycle can be supported by the basal temperature method. Basal temperature is the body's resting temperature. This method involves daily measurement of basal temperature every morning at the same time before getting out of bed. The temperature usually rises slightly after ovulation has occurred. Once the menstrual cycle has been tracked over several months, the day of ovulation can be more accurately predicted. Additionally, the consistency and appearance of cervical mucus change during the menstrual cycle. During ovulation, the mucus typically becomes clearer and more stretchy. In some cases, fertility tests can assist in determining ovulation. An ovulation test detects the rise in LH hormone in the urine, which typically occurs about 24-48 hours before ovulation, so positive test results can indicate impending ovulation<sup>[30]</sup>.

Infertility in women can be influenced by a variety of factors. Irregular or absent menstrual cycles may indicate hormonal problems that can affect fertility. Blocked or damaged fallopian tubes can prevent the egg from reaching the uterus, while uterine abnormalities can hinder the implantation of a fertilized egg.



### What is recommended if there is suspicion of an HPV infection?

If there is suspicion of an HPV infection due to changes in the bacterial community, it is advisable to consult a gynecologist who will then make a diagnosis. Most HPV infections resolve on their own and do not require treatment. A healthy lifestyle, including a balanced diet, regular exercise, and adequate sleep, can help strengthen the immune system in fighting the HPV infection. If action is needed, your doctor will recommend the best steps for treating or monitoring the HPV infection based on your individual circumstances. If you have not already been vaccinated against HPV, vaccination may be considered depending on your health status. When sexual activities, especially sexual intercourse, are involved, barrier methods such as condoms (for both women and men) should be used to reduce the risk of HPV transmission <sup>[31]</sup>.

Preventing HPV infections is important to reduce the risk of serious health problems, including cervical cancer. HPV vaccination can protect against the most common high-risk HPV types that can cause cancer, among other conditions. Vaccination is particularly effective when administered at a young age, before sexual activity begins. While condoms do not offer one hundred percent protection, they can reduce the risk of an HPV infection. Using condoms during sexual activity, especially with multiple sexual partners, can help minimize the transmission of HPV. Regular gynecological examinations and Pap smears are part of the early detection of HPV infections and related conditions. Therefore, regular check-ups should be attended. A healthy lifestyle that promotes immune system strength can make it more difficult for HPV to spread in the body. Smoking, in particular, is associated with an increased risk of HPV-related cancers <sup>[32]</sup>.



### What can I do about a yeast infection and prevent it in the future?

If you have signs or symptoms of a yeast infection in the vagina, it is important to consult a doctor to receive an accurate diagnosis and the proper treatment. Vaginal yeast infections, in particular, are typically treated with antifungals in the form of vaginal creams, tablets, or suppositories. Ensure that you use the prescribed medications according to your doctor's instructions or the instructions on the packaging. Most antifungals are applied over a specific period. During treatment, it's advisable to avoid irritating substances, such as perfumed soaps, intimate sprays, or tight synthetic underwear that could worsen the symptoms. <sup>[37]</sup>

A balanced diet rich in vegetables, whole grains, and about 200g of fruit daily can strengthen the immune system and contribute to prevention. Particularly, reducing excessive sugar consumption can decrease the risk of yeast infections, as fungal growth is promoted by sugar.

It is also highly beneficial if underwear is washed at a minimum of 60°C. Additionally, breathable cotton underwear supports a healthy microclimate by wicking moisture away from the vaginal area. Loose-fitting clothing also contributes to air circulation. If the intimate area is regularly shaved, the razor should always be disinfected before use.

Aggressive hygiene products and the unnecessary use of antibiotics can disrupt the balance of the vaginal microbiome, promoting the proliferation of yeast fungi. <sup>[37][38]</sup>

- [1] Verstraelen, Hans, et al. „The vaginal microbiome: I. research development, lexicon, defining “normal” and the dynamics throughout women’s lives.” *Journal of Lower Genital Tract Disease* 26.1 (2022): 73.
- [2] Barrientos-Durán, Antonio, et al. „Reviewing the composition of vaginal microbiota: inclusion of nutrition and probiotic factors in the maintenance of eubiosis.” *Nutrients* 12.2 (2020): 419.
- [3] Baud, Agnes, et al. „Microbial diversity in the vaginal microbiota and its link to pregnancy outcomes.” *Scientific Reports* 13.1 (2023): 9061.
- [4] Ser, Hooi-Leng, et al. „Current Updates on the Role of Microbiome in Endometriosis: A Narrative Review.” *Microorganisms* 11.2 (2023): 360.
- [5] Chao, Xiaopei, et al. „The role of the vaginal microbiome in distinguishing female chronic pelvic pain caused by endometriosis/adenomyosis.” *Annals of Translational Medicine* 9.9 (2021).
- [6] Ata, Baris, et al. „The endobiota study: comparison of vaginal, cervical and gut microbiota between women with stage 3/4 endometriosis and healthy controls.” *Scientific reports* 9.1 (2019): 2204.
- [7] Vitale, Salvatore Giovanni, et al. „The role of genital tract microbiome in fertility: a systematic review.” *International Journal of Molecular Sciences* 23.1 (2021): 180.
- [8] Uledo, B., et al. „Identification of vaginal microbiome associated with IVF pregnancy.” *Scientific Reports* 12.1 (2022): 6807.
- [9] Kaur, Harisham, et al. „Crosstalk between female gonadal hormones and vaginal microbiota across various phases of women’s gynecological lifecycle.” *Frontiers in microbiology* 11 (2020): 551.
- [10] Babu, Geethavani, et al. „Comparative study on the vaginal flora and incidence of asymptomatic vaginosis among healthy women and in women with infertility problems of reproductive age.” *Journal of clinical and diagnostic research: JCDR* 11.8 (2017): DC18.
- [11] Wang, Yuanyuan, et al. „Influences of vaginal microbiota on human papillomavirus infection and host immune regulation: What we have learned?”. *Decoding Infection and Transmission* 1 (2023): 10002.
- [12] Chao, Xiaopei, et al. „Research of the potential biomarkers in vaginal microbiome for persistent high-risk human papillomavirus infection.” *Annals of Translational Medicine* 8.4 (2020).
- [13] Chen, Yulian, et al. „Human papillomavirus infection and cervical intraepithelial neoplasia progression are associated with increased vaginal microbiome diversity in a Chinese cohort.” *BMC infectious diseases* 20 (2020): 1-12.
- [14] Lin, Wenyu, et al. „Changes of the vaginal microbiota in HPV infection and cervical intraepithelial neoplasia: a cross-sectional analysis.” *Scientific Reports* 12.1 (2022): 2812.
- [15] YH, Neggers, Nansel TR, and W. W. Andrews. „Dietary intake of selected nutrients affects bacterial vaginosis in women.” *J Nutr* 137 (2007): 2128-2133.
- [16] Thoma, Marie E., et al. „Bacterial vaginosis is associated with variation in dietary indices.” *The Journal of nutrition* 141.9 (2011): 1698-1704.
- [17] Yoshikata, Remi, et al. „Age-Related Changes, Influencing Factors, and Crosstalk Between Vaginal and Gut Microbiota: A Cross-Sectional Comparative Study of Pre-and Postmenopausal Women.” *Journal of Women’s Health* 31.12 (2022): 1763-1772.
- [18] Park, Min Gu, Seok Cho, and Mi Mi Oh. „Menopausal Changes in the Microbiome—A Review Focused on the Genitourinary Microbiome.” *Diagnostics* 13.6 (2023): 1193.
- [19] Yamamoto, Ayae, et al. „A prospective cohort study of meat and fish consumption and endometriosis risk.” *American journal of obstetrics and gynecology* 219.2 (2018): 178-e1.
- [20] Ahrens, Peter, et al. „Changes in the vaginal microbiota following antibiotic treatment for Mycoplasma genitalium, Chlamydia trachomatis and bacterial vaginosis.” *PloS one* 15.7 (2020): e0236036.
- [21] Daher, Alaa, et al. „Intimate hygiene practices and reproductive tract infections: A systematic review.” *Gynecology and Obstetrics Clinical Medicine* (2022).
- [22] Chen, Ying, et al. „Role of female intimate hygiene in vulvovaginal health: Global hygiene practices and product usage.” *Women’s Health* 13.3 (2017): 58-67.
- [23] Parasar, Parveen, Pinar Ozcan, and Kathryn L. Terry. „Endometriosis: epidemiology, diagnosis and clinical management.” *Current obstetrics and gynecology reports* 6 (2017): 34-41.
- [24] Barnard, Neal D., et al. „Nutrition in the prevention and treatment of endometriosis: A review.” *Frontiers in Nutrition* 10 (2023): 1089891.
- [25] Singh, Kadambari, and Deepika Dewani. „Recent Advancements in In Vitro Fertilisation.” *Cureus* 14.10 (2022).
- [26] Skoracka, Kinga, et al. „Female fertility and the nutritional approach: the most essential aspects.” *Advances in nutrition* 12.6 (2021): 2372-2386.
- [27] Silvestris, Erica, et al. „Obesity as disruptor of the female fertility.” *Reproductive Biology and Endocrinology* 16 (2018): 1-13.
- [28] Brinson, Alison K., et al. „Impact of Physical Activity and Sedentary Behavior on Spontaneous Female and Male Fertility: A Systematic Review.” *Journal of Physical Activity and Health* 20.7 (2023): 600-615.
- [29] Wilcox, Allen J., David Dunson, and Donna Day Baird. „The timing of the “fertile window” in the menstrual cycle: day specific estimates from a prospective study.” *Bmj* 321.7271 (2000): 1259-1262.
- [30] Kumar, Pratap, and Sameer Farouk Sait. „Luteinizing hormone and its dilemma in ovulation induction.” *Journal of human reproductive sciences* 4.1 (2011): 2-7.
- [31] Soheili, Maryam, et al. „Human papilloma virus: A review study of epidemiology, carcinogenesis, diagnostic methods, and treatment of all HPV-related cancers.” *Medical journal of the Islamic Republic of Iran* 35 (2021): 65.
- [32] Castle, Philip E. „How does tobacco smoke contribute to cervical carcinogenesis?”. *Journal of virology* 82.12 (2008): 6084-6086.
- [33] Willems, Hubertine ME, et al. „Vulvovaginal candidiasis: a current understanding and burning questions.” *Journal of Fungi* 6.1 (2020): 27.
- [34] Bradford, L. Latéy, and Jacques Ravel. „The vaginal microbiome: A contemporary perspective on fungi in women’s health and diseases.” *Virulence* 8.3 (2017): 342-351.
- [35] Kalaiarasan, Krishnapriya, Rakesh Singh, and Latha Chaturvedula. „Fungal profile of vulvovaginal candidiasis in a tertiary care hospital.” *Journal of clinical and diagnostic research: JCDR* 11.3 (2017): DC06.
- [36] Zhao, Changying, et al. „Mycobiome Study Reveals Different Pathogens of Vulvovaginal Candidiasis Shape Characteristic Vaginal Bacteriome.” *Microbiology Spectrum* (2023): e03152-22.
- [37] Chen, Ying, et al. „Role of female intimate hygiene in vulvovaginal health: Global hygiene practices and product usage.” *Women’s Health* 13.3 (2017): 58-67.
- [38] Man, Adrian, et al. „New perspectives on the nutritional factors influencing growth rate of Candida albicans in diabetics. An in vitro study.” *Memórias do Instituto Oswaldo Cruz* 112 (2017): 587-592.

---

---

---

---

---

---

---

---

---

---

## Miscellaneous

### Results report prepared by:

**Procomcure Biotech GmbH**

Breitwies 1  
5303 Thalgau  
Österreich

### Measuring method:

**NGS**


Next-Generation-Sequencing (16S rRNA Gen)

### Primary sample or submitted material:

Vaginal sample

### Disclaimer:

The analysis is based on the sequencing of the 16S rRNA gene, which allows for the classification of bacterial strains in the microbiome. The results of the microbiome test and its interpretation may be incomplete. The number of detected microorganisms is not exhaustive, and there may be other microorganisms present that were not captured by the sequencing. The current interpretation of the microbiome test is based on data from adult reference individuals and may change in the future due to the publication of new scientific studies. Inaccurate or missing information can lead to misleading interpretation. This report is provided solely for informational and educational purposes and does not replace a visit to a doctor or the advice or services of a physician.



*Health begins with us.*

