

---

## Second homework assignment

---

PhD in Business Economics  
**Professor:** Hedibert Freitas Lopes

Course: Advanced Bayesian Econometrics  
Due date: 12h, February 11th, 2021.

---

Use, preferably, **Rmarkdown** (via **RStudio**) to produce your report in PDF or HTML.

---

### Fitting Gaussian and Student's $t$ ARMA(1,1) model

Let us assume that some observed time series data  $\{y_1, \dots, y_n\}$  follows an ARMA(1,1) model

$$y_t = \phi y_{t-1} + \varepsilon_t + \gamma \varepsilon_{t-1}$$

where  $\varepsilon_1, \dots, \varepsilon_n$  are i.i.d. either  $\mathcal{M}_0 : N(0, \sigma^2)$  or  $\mathcal{M}_1 : t_\nu(0, \tau^2)$ , where  $\tau^2 = (\nu - 2)/\nu\sigma^2$ . We will keep  $\nu$  fixed and known throughout. In addition, in order to simplify the homework, we will assume that  $y_0 = \varepsilon_0 = 0$ . Therefore, it is easy to see that  $\{\varepsilon_1, \dots, \varepsilon_n\}$  are deterministically obtained from  $\theta = (\phi, \gamma, \sigma)$  and the data  $y^n = \{y_1, \dots, y_n\}$ :  $\varepsilon_1 = y_1$  and  $\varepsilon_t = y_t - \phi y_{t-1} - \gamma \varepsilon_{t-1}$ , for  $t = 2, \dots, n$ .

**Likelihood functions.** To avoid overloading the notation, let us drop  $(\varepsilon_0, y_0)$  in what follows. The likelihood functions are, therefore,

$$\begin{aligned}\mathcal{L}(\theta|y^n, \mathcal{M}_0) &= (2\pi\sigma^2)^{-n/2} \exp\left\{-\frac{\sum_{t=1}^n \varepsilon_t^2}{2\sigma^2}\right\} \\ \mathcal{L}(\theta|y^n, \mathcal{M}_1) &= \left(\frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})\sqrt{\pi\nu\tau^2}}\right)^n \prod_{t=1}^n \left(1 + \frac{1}{\nu} \frac{\varepsilon_t^2}{\tau^2}\right)^{-\frac{\nu+1}{2}}.\end{aligned}$$

**Prior distribution.** Let us assume that

$$p(\theta) = p(\phi, \gamma, \sigma^2) = p(\phi)p(\gamma)p(\sigma^2),$$

for

$$\begin{aligned}\phi &\sim U(-1, 1) \\ \gamma &\sim U(-1, 1) \\ \sigma^2 &\sim IG(5/2, 5(1.4)/2).\end{aligned}$$

Hence, we are constraining our inference to the class of stationary and invertible ARMA(1,1) models. In addition, prior mean, mode and standard deviation for  $\sigma^2$  is around 2.33, 1 and 3.3, respectively. Also,  $Pr(\sigma^2 \in (0.33, 26.3)) \approx 99.9\%$ , so  $\sigma < 0.5$  or  $\sigma > 5$  are essentially ruled out as well.

**Simulating some data.** You should simulate two datasets of size  $n = 400$ , one with Gaussian errors and the other with Student's  $t$  errors where  $\sigma = 1$ ,  $\nu = 4$ ,  $\phi = 0.98$  and  $\gamma = -0.64$ . Feel free to use the following R script:

```
set.seed(12345)
n = 400
sig = 1.0
nu = 4
phi = 0.98
theta = -0.64
tau = sqrt((nu-2)/nu)*sig
e.n = sig*rnorm(n)
e.t = tau*rt(n,df=nu)
y.n = rep(0,n)
y.t = rep(0,n)
y.n[1] = e.n[1]
y.t[1] = e.t[1]
for (t in 2:n){
  y.n[t] = phi*y.n[t-1]+e.n[t]+theta*e.n[t-1]
  y.t[t] = phi*y.t[t-1]+e.t[t]+theta*e.t[t-1]
}
par(mfrow=c(1,1))
ts.plot(cbind(y.n,y.t),col=1:2,main="ARMA(1,1) data")
legend("bottomleft",legend=c("Gaussian","Student's t"),col=1:2,lty=1,bty="n")
```

**Questions:** Answer the following questions for each one of the two datasets generated by the previous script.

1. **Maximum likelihood inference.**

What are the maximum likelihood estimates (MLE) of  $\theta$  under both models? Use the R function `nlm` to minimize the negative of the likelihood functions. Are the results similar to the ones from the R function `arima(y,order=c(1,0,1))`?

2. **Bayesian inference via Monte Carlo methods.**

- (a) Use sampling importance resampling (SIR) to sample from both posterior distributions of  $\theta$ :

$$\begin{aligned} p(\theta|y^n, \mathcal{M}_0) &\propto \mathcal{L}(\theta|y^n, \mathcal{M}_0)p(\theta) \\ p(\theta|y^n, \mathcal{M}_1) &\propto \mathcal{L}(\theta|y^n, \mathcal{M}_1)p(\theta). \end{aligned}$$

Use these draws whenever necessary in the next several questions.

- (b) Compute posterior means, medians and 95% credibility interval for  $\phi$ ,  $\gamma$  and  $\sigma^2$ . Are posterior means (and medians) similar to their MLE counterparts?

- (c) Plot the contours of the posterior density. What are the posterior probabilities that  $\phi > 0.9$  under both models?

### 3. Prior predictive, Bayes factor and posterior model probability.

- (a) Compute both **prior predictive**  $p(y^n|\mathcal{M}_0)$  and  $p(y^n|\mathcal{M}_1)$ . We can approximate the prior predictive densities, for  $j = 0, 1$

$$p(y^n|\mathcal{M}_j) = \int_0^\infty \int_{-1}^1 \int_{-1}^1 \prod_{t=1}^n p(y_t|y_{t-1}, \theta, \mathcal{M}_j) d\phi d\gamma d\sigma^2,$$

via Monte Carlo by

$$\hat{p}(y^n|\mathcal{M}_j) = \frac{1}{M} \sum_{i=1}^M p(y_t|y_{t-1}, \phi^{(i)}, \gamma^{(i)}, \sigma^{2(i)}|\mathcal{M}_j),$$

where  $\theta^{(1)}, \dots, \theta^{(M)}$  are draws from the prior  $p(\theta)$ . Let us use  $M = 100,000$ .

- (b) Compute a MC approximation to the **Bayes factor**:

$$B_{01} = \frac{p(y^n|\mathcal{M}_0)}{p(y^n|\mathcal{M}_1)}.$$

- (c) Finally, the posterior model odds can be computed as

$$\frac{Pr(\mathcal{M}_0|y^n)}{Pr(\mathcal{M}_1|y^n)} = \frac{Pr(\mathcal{M}_0)}{Pr(\mathcal{M}_1)} \times B_{01},$$

where  $Pr(\mathcal{M}_0)$  and  $Pr(\mathcal{M}_1)$  are the prior probabilities assigned to models  $\mathcal{M}_0$  and  $\mathcal{M}_1$ , respectively. Assuming  $Pr(\mathcal{M}_0) = Pr(\mathcal{M}_1)$ , obtain a MC approximation to  $Pr(\mathcal{M}_0|y^n)$ , the **posterior model probability** of the Gaussian model.

Discuss your findings.