

# Arithmetic: A Programmatic Approach

Murisi Tarusenga

Thursday 16<sup>th</sup> January, 2020 15:52

# ReadMe

What follows is a reformulation of the elementary parts of number theory, hard analysis, calculus, and linear algebra in terms of a system of self-evidently correct procedures for achieving particular objectives, objectives like showing that modular exponentiation of a specific integer with specific properties yields a stated result. So, while formal mathematics usually takes the format of definition-theorem-proof, this project has the format of declaration-procedure objective-procedure implementation. So where there usually would have been a statement and proof of Euler's totient theorem, [procedure I:75](#) is provided, and where there would have been a definition of Euler's totient function, [declaration I:27](#) is provided.

At this point the natural question is whether it is always possible to write a procedure in such a way that its correctness is self-evident. While I have not been able to come up with an theoretical argument that that should be the case, actually writing out numerous procedures to achieve a range of objectives has convinced me that this indeed is the case. And as I refine and extend the following procedures, it is becoming clearer to me that a mathematical proof does not necessarily have to be some sort of argument. Rather, what is turning out to be important is the granularity of the writing's subdivisions (i.e. sub-procedures in programming and lemmas in mathematics) and the communication of intent (i.e. comments in programming and theorem statements in mathematics).

For the purposes of storage and transmission of knowledge pertaining to the elementary parts of number theory, hard analysis, calculus, and linear algebra, the following procedures are interchangeable with their analogous proofs in the sense that, assuming equal competence in programming and proving, if you have the procedure objective and implementation, you can trivially generate the analogous theorem and proof, and if you are in possession of the theorem and proof, then you can trivially generate the analogous procedure objective and implementation.

# Contents

<b>I</b>	<b>Integer Arithmetic</b>	<b>7</b>
1	Integer Arithmetic	8
2	Modular Arithmetic	16
3	Congruence Equations	30
4	Permutations and Combinations	38
<b>II</b>	<b>Rational Arithmetic</b>	<b>41</b>
5	Rational Arithmetic	42
6	Polynomial Arithmetic	51
7	Polynomial Sign Changes	60
<b>III</b>	<b>Complex Arithmetic</b>	<b>68</b>
8	Complex Arithmetic	69
9	Exponential and Trigonometric Functions	77
10	Binomial and Mercator Series	85
11	Gregory-Leibniz Series	97
<b>IV</b>	<b>Differential Arithmetic</b>	<b>109</b>
12	Differential Arithmetic	110
13	Common Derivatives	117
14	Integral Arithmetic	133
<b>V</b>	<b>Matrix Arithmetic</b>	<b>137</b>
15	Matrix Arithmetic	138

<b>16 Compound Matrices</b>	<b>148</b>
<b>17 Polynomials and Normal Forms</b>	<b>156</b>

# Declarations

**integer** . 8

$\text{po}(a)$  positive part of  $a$ . 8

$\text{ne}(a)$  negative part of  $a$ . 8

$a = b$  integer equality. 8

$a + b$  integer addition. 9

$a$  . 9

$-a$  integer negation. 10

$ab$  integer multiplication. 10

$a < b$  integer less than. 12

$\|a\|$  absolute value. 14

$\text{sgn}(a)$  sign function. 15

$a \text{ div } b$  integer division. 16

$a \bmod b$  integer modulus. 16

$a \equiv b \pmod{c}$  modular equality. 16

$(a, b)$  . 19

$(a_0, a_1, \dots, a_{n-1})$  . 22

**prime number** . 23

$|a|$  length of list. 24

$a \frown b$  list concatenation. 24

$f(R)$  elementwise operation. 24

$a_*$  product of list. 24

$\prod_r^R f(r)$  pi product notation. 24

$[a : b]$  integer range. 25

$[a, b]$  . 27

$[a_0, a_1, \dots, a_{n-1}]$  . 28

$\chi_{b,d}(a, c)$  . 30

$\chi_{b_0, b_1, \dots, b_{n-1}}(a_0, a_1, \dots, a_{n-1})$  . 32

$\phi(n)$  Euler's phi function. 32

$a \times b$  Cartesian product. 34

$[P]$  Iverson bracket. 36

$a_+$  sum of list. 36

$\sum_r^R f(r)$  sigma summation notation. 36

$a^b$  falling power. 38

$a^{\bar{b}}$  rising power. 38

$\binom{n}{r}$  binomial coefficient. 38

**rational number** . 42

$\text{nu}(a)$  numerator of  $a$ . 42

$\text{de}(a)$  denominator of  $a$ . 42

$a = b$  rational equality. 42

$a + b$  rational addition. 43

$a$  . 44

$-a$  rational negation. 44

$ab$  rational multiplication. 44

$\frac{1}{a}$  rational reciprocal. 45

$a < b$  rational less than. 47

$\lfloor a \rfloor$  floor function. 49

$\lceil a \rceil$  ceiling function. 49

$\min(c)$  minimum of list. 51

$\min_r^R c(r)$  minimum notation. 51

$\max(c)$  maximum of list. 51

$\max_r^R c(r)$  maximum notation. 51

**polynomial** . 51

$a_i$  polynomial coefficient. 51

$a = b$  polynomial equality. 51

$\Lambda(a, b)$  polynomial evaluation. 51

$\langle f(j) \text{ for } j \in R \rangle$  list comprehension. 52

$a + b$  polynomial addition. 52

$a$  . 53

$-a$  polynomial negation. 54

$ab$  polynomial multiplication. 54

$\lambda$  . 56

$\deg(a)$  polynomial degree. 57

**monic polynomial** . 58

$\text{mon}(p)$  . 58

$a \text{ div } b$  polynomial division. 59

$a \bmod b$  polynomial modulus. 59

$J_s(x)$  . 63

**complex number** . 69

$\text{re}(a)$  real part of  $a$ . 69

$\text{im}(a)$  imaginary part of  $a$ . 69

$a = b$  complex equality. 69

$a + b$  complex addition. 70

$a$  . 70

$-a$  complex negation. 71

$ab$  complex multiplication. 71

$\bar{a}$  complex conjugate. 72

$\|a\|^2$  Euclidean length squared. 72

$\frac{1}{a}$  complex reciprocal. 74

$i$  imaginary number. 75

$a \equiv b$  (err  $c_1$ ) (err  $c_2$ )  $\cdots$  (err  $c_n$ ) approximate equality. 75

$\exp_n(a)$  complex exponential function. 77

$\cos_n(z)$  cosine. 82

$\sin_n(z)$  sine. 82

$(1+x)_n^a$  binomial series. 85

$\omega(r)$  . 94

$\ln_k(1+x)$  natural logarithm. 94

$\tau_n$  tau. 97

**complex polynomial** . 105

$\{x\}$  taxicab length. 110

$\Delta_{x=y}^z f(x)$  difference quotient. 112

$\ln_n(x)$  natural logarithm. 122

$x_n^a$  exponentiation. 127

$\int_r^R f(r, \delta_r)$  Riemann sum. 133

$\Delta X$  first difference. 134

**matrix** . 138

$A_{I,J}$  submatrix. 138

$A = B$  matrix equality. 138

$A + B$  matrix addition. 139

$0_{m \times n}$   $m \times n$  zero matrix. 139

$-A$  matrix negation. 139

$AB$  matrix multiplication. 140

$a_{m \times m}$  scalar matrix. 141

$A_{i,*}$  matrix row. 141

$A_{*,i}$  matrix column. 141

**matrix diagonal** . 142

**diagonal matrix** . 142

**tilt matrix** . 143

$A^{-1}$  . 143

$\text{rows}(A)$  number of rows of  $A$ . 146

$\text{cols}(A)$  number of columns of  $A$ . 146

$\text{diag}(C)$  block diagonal matrix. 146

$\det(A)$  matrix determinant. 148

$C_k(A)$   $k^{\text{th}}$  compound matrix. 151

$A_{I,J}$  labelled matrix entry. 151

$A^T$  matrix transpose. 154

$A \setminus B$  matrix left division. 156  
 $A / B$  matrix right division. 156  
 $(e_i)_{k \times 1}$  standard unit vector. 160  
 $\text{mat}_t(p)$  . 160  
 $\text{comp}(p)$  companion matrix. 160  
 $\text{last}_A$  last polynomial. 163  
 $\text{pows}(A)$  . 165  
 $\text{tr}(A)$  matrix trace. 166  
**symmetric matrix** . 167  
 $\text{sel}_A$  selector polynomial. 168

## Part I

# Integer Arithmetic



# Chapter 1

## Integer Arithmetic

### Declaration I:0

The phrase "integer" will be used as a shorthand for an ordered pair of natural numbers.

### Declaration I:1

The phrase "the positive part of  $a$ " and the notation  $\text{po}(a)$ , where  $a$  is an integer, will be used as a shorthand for the first entry of  $a$ .

### Declaration I:2

The phrase "the negative part of  $a$ " and the notation  $\text{ne}(a)$ , where  $a$  is an integer, will be used as a shorthand for the second entry of  $a$ .

### Declaration I:3

The phrase " $a = b$ ", where  $a, b$  are integers, will be used as a shorthand for " $\text{po}(a) + \text{ne}(b) = \text{ne}(a) + \text{po}(b)$ ".

### Procedure I:0(1.65)

#### Objective

Choose an integer  $a$ . The objective of the following instructions is to show that  $a = a$ .

### Implementation

1. **Show that  $a = a$  given that  $\text{po}(a) + \text{ne}(a) = \text{ne}(a) + \text{po}(a)$ .**

### Procedure I:1(1.66)

#### Objective

Choose two integers  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $b = a$ .

### Implementation

1. Using **declaration I:3**, show that  $b = a$ 
  - (a) given that  $\text{po}(b) + \text{ne}(a) = \text{ne}(b) + \text{po}(a)$
  - (b) given that  $\text{po}(a) + \text{ne}(b) = \text{ne}(a) + \text{po}(b)$
  - (c) given that  $a = b$ .

### Procedure I:2(1.67)

#### Objective

Choose three integers  $a, b, c$  such that  $a = b$  and  $b = c$ . The objective of the following instructions is to show that  $a = c$ .

### Implementation

1. Show that  $\text{po}(a) + \text{ne}(b) = \text{ne}(a) + \text{po}(b)$  using **declaration I:3**.
2. Show that  $\text{po}(b) + \text{ne}(c) = \text{ne}(b) + \text{po}(c)$  using **declaration I:3**.
3. Hence show that  $a = c$ 
  - (a) given that  $\text{po}(a) + \text{ne}(c) = \text{ne}(a) + \text{po}(c)$
  - (b) given that  $\text{po}(a) + \text{ne}(b) + \text{po}(b) + \text{ne}(c) = \text{ne}(a) + \text{po}(b) + \text{ne}(b) + \text{po}(c)$ .

### Declaration I:4

The notation  $a + b$ , where  $a, b$  are integers, will be used as a shorthand for the pair  $\langle \text{po}(a) + \text{po}(b), \text{ne}(a) + \text{ne}(b) \rangle$ .

### Procedure I:3(1.68)

#### Objective

Choose four integers  $a, b, c, d$  such that  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $a + b = c + d$ .

### Implementation

1. Show that  $\text{po}(a) + \text{ne}(c) = \text{ne}(a) + \text{po}(c)$  using **declaration I:3**.
2. Show that  $\text{po}(b) + \text{ne}(d) = \text{ne}(b) + \text{po}(d)$  using **declaration I:3**.
3. Hence show that  $a + b$ 
  - (a)  $= \langle \text{po}(a), \text{ne}(a) \rangle + \langle \text{po}(b), \text{ne}(b) \rangle$
  - (b)  $= \langle \text{po}(a) + \text{po}(b), \text{ne}(a) + \text{ne}(b) \rangle$
  - (c)  $= \langle \text{po}(a) + \text{po}(b) + \text{ne}(c) + \text{ne}(d), \text{ne}(a) + \text{ne}(b) + \text{ne}(c) + \text{ne}(d) \rangle$
  - (d)  $= \langle (\text{po}(a) + \text{ne}(c)) + (\text{po}(b) + \text{ne}(d)), \text{ne}(a) + \text{ne}(b) + \text{ne}(c) + \text{ne}(d) \rangle$
  - (e)  $= \langle (\text{ne}(a) + \text{po}(c)) + (\text{ne}(b) + \text{po}(d)), \text{ne}(a) + \text{ne}(b) + \text{ne}(c) + \text{ne}(d) \rangle$
  - (f)  $= \langle \text{ne}(a) + \text{ne}(b) + \text{po}(c) + \text{po}(d), \text{ne}(a) + \text{ne}(b) + \text{ne}(c) + \text{ne}(d) \rangle$
  - (g)  $= \langle \text{po}(c) + \text{po}(d), \text{ne}(c) + \text{ne}(d) \rangle$

$$(h) = \langle \text{po}(c), \text{ne}(c) \rangle + \langle \text{po}(d), \text{ne}(d) \rangle$$

$$(i) = c + d.$$

### Procedure I:4(1.69)

#### Objective

Choose three integers  $a, b, c$ . The objective of the following instructions is to show that  $(a + b) + c = a + (b + c)$ .

### Implementation

1. Show that  $(a + b) + c$ 
  - (a)  $= \langle \text{po}(a) + \text{po}(b), \text{ne}(a) + \text{ne}(b) \rangle + \langle \text{po}(c), \text{ne}(c) \rangle$
  - (b)  $= \langle (\text{po}(a) + \text{po}(b)) + \text{po}(c), (\text{ne}(a) + \text{ne}(b)) + \text{ne}(c) \rangle$
  - (c)  $= \langle \text{po}(a) + (\text{po}(b) + \text{po}(c)), \text{ne}(a) + (\text{ne}(b) + \text{ne}(c)) \rangle$
  - (d)  $= \langle \text{po}(a), \text{ne}(a) \rangle + \langle \text{po}(b) + \text{po}(c), \text{ne}(b) + \text{ne}(c) \rangle$
  - (e)  $= a + (b + c)$ .

### Procedure I:5(1.70)

#### Objective

Choose two integers  $a, b$ . The objective of the following instructions is to show that  $a + b = b + a$ .

### Implementation

1. Show that  $a + b$ 
  - (a)  $= \langle \text{po}(a) + \text{po}(b), \text{ne}(a) + \text{ne}(b) \rangle$
  - (b)  $= \langle \text{po}(b) + \text{po}(a), \text{ne}(b) + \text{ne}(a) \rangle$
  - (c)  $= b + a$ .

### Declaration I:5

The notation  $a$ , where  $a$  is a natural number, will contextually be used as a shorthand for the pair  $\langle a, 0 \rangle$ .

### Procedure I:6(1.71)

#### Objective

Choose an integer  $a$ . The objective of the following instructions is to show that  $0 + a = a$ .

#### Implementation

1. Show that  $0 + a$

$$(a) = \langle 0, 0 \rangle + \langle \text{po}(a), \text{ne}(a) \rangle$$

$$(b) = \langle 0 + \text{po}(a), 0 + \text{ne}(a) \rangle$$

$$(c) = \langle \text{po}(a), \text{ne}(a) \rangle$$

$$(d) = a.$$

#### Declaration I:6

The notation  $-a$ , where  $a$  is an integer, will be used as a shorthand for the pair  $\langle \text{ne}(a), \text{po}(a) \rangle$ .

### Procedure I:7(1.72)

#### Objective

Choose two integers  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $-a = -b$ .

#### Implementation

1. Show that  $\text{po}(a) + \text{ne}(b) = \text{ne}(a) + \text{po}(b)$  using **declaration I:3**.

2. Hence show that  $-a$

$$(a) = \langle \text{ne}(a), \text{po}(a) \rangle$$

$$(b) = \langle \text{ne}(a) + \text{po}(b), \text{po}(a) + \text{po}(b) \rangle$$

$$(c) = \langle \text{po}(a) + \text{ne}(b), \text{po}(a) + \text{po}(b) \rangle$$

$$(d) = \langle \text{ne}(b), \text{po}(b) \rangle$$

$$(e) = -b.$$

### Procedure I:8(1.73)

#### Objective

Choose an integer  $a$ . The objective of the following instructions is to show that  $-a + a = 0$ .

#### Implementation

1. Show that  $-a + a$

$$(a) = (-a) + a$$

$$(b) = \langle \text{ne}(a), \text{po}(a) \rangle + \langle \text{po}(a), \text{ne}(a) \rangle$$

$$(c) = \langle \text{ne}(a) + \text{po}(a), \text{po}(a) + \text{ne}(a) \rangle$$

$$(d) = \langle 0, 0 \rangle$$

$$(e) = 0.$$

#### Declaration I:7

The notation  $ab$ , where  $a, b$  are integers, will be used as a shorthand for the pair  $\langle \text{po}(a) \text{po}(b) + \text{ne}(a) \text{ne}(b), \text{po}(a) \text{ne}(b) + \text{ne}(a) \text{po}(b) \rangle$ .

### Procedure I:9(1.74)

#### Objective

Choose four integers  $a, b, c, d$  such that  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $ab = cd$ .

#### Implementation

1. Show that  $\text{po}(a) + \text{ne}(c) = \text{ne}(a) + \text{po}(c)$  using **declaration I:3**.

2. Show that  $\text{po}(b) + \text{ne}(d) = \text{ne}(b) + \text{po}(d)$  using **declaration I:3**.

3. Hence show that  $ab$

$$(a) = \langle \text{po}(a) \text{po}(b) + \text{ne}(a) \text{ne}(b), \text{po}(a) \text{ne}(b) + \text{ne}(a) \text{po}(b) \rangle$$

$$(b) = \langle \text{po}(a) \text{po}(b) + \text{ne}(a) \text{ne}(b) + \text{po}(a) \text{ne}(d) + \text{ne}(c) \text{po}(d) + \text{po}(c) \text{ne}(d), \text{po}(a) \text{ne}(b) + \text{ne}(a) \text{po}(b) + \text{po}(a) \text{ne}(d) + \text{ne}(c) \text{po}(d) + \text{po}(c) \text{ne}(d) \rangle$$

$$(c) = \langle \text{po}(a)(\text{po}(b) + \text{ne}(d)) + \text{ne}(a)\text{ne}(b) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d), \text{po}(a)\text{ne}(b) + \text{ne}(a)\text{po}(b) + \text{po}(a)\text{ne}(d) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(d) = \langle \text{po}(a)(\text{ne}(b) + \text{po}(d)) + \text{ne}(a)\text{ne}(b) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d), \text{po}(a)\text{ne}(b) + \text{ne}(a)\text{po}(b) + \text{po}(a)\text{ne}(d) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(e) = \langle (\text{po}(a) + \text{ne}(c))\text{po}(d) + \text{ne}(a)\text{ne}(b) + \text{po}(c)\text{ne}(d), \text{ne}(a)\text{po}(b) + \text{po}(a)\text{ne}(d) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(f) = \langle (\text{ne}(a) + \text{po}(c))\text{po}(d) + \text{ne}(a)\text{ne}(b) + \text{po}(c)\text{ne}(d), \text{ne}(a)\text{po}(b) + \text{po}(a)\text{ne}(d) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(g) = \langle \text{ne}(a)(\text{po}(d) + \text{ne}(b)) + \text{po}(c)\text{po}(d) + \text{po}(c)\text{ne}(d), \text{ne}(a)\text{po}(b) + \text{po}(a)\text{ne}(d) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(h) = \langle \text{ne}(a)(\text{po}(b) + \text{ne}(d)) + \text{po}(c)\text{po}(d) + \text{po}(c)\text{ne}(d), \text{ne}(a)\text{po}(b) + \text{po}(a)\text{ne}(d) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(i) = \langle (\text{ne}(a) + \text{po}(c))\text{ne}(d) + \text{po}(c)\text{po}(d), \text{po}(a)\text{ne}(d) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(j) = \langle (\text{po}(a) + \text{ne}(c))\text{ne}(d) + \text{po}(c)\text{po}(d), \text{po}(a)\text{ne}(d) + \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(k) = \langle \text{ne}(c)\text{ne}(d) + \text{po}(c)\text{po}(d), \text{ne}(c)\text{po}(d) + \text{po}(c)\text{ne}(d) \rangle$$

$$(l) = cd.$$

## Procedure I:10(1.75)

### Objective

Choose three integers  $a, b, c$ . The objective of the following instructions is to show that  $(ab)c = a(bc)$ .

### Implementation

1. Show that  $(ab)c$

$$(a) = \langle \text{po}(a)\text{po}(b) + \text{ne}(a)\text{ne}(b), \text{po}(a)\text{ne}(b) + \text{ne}(a)\text{po}(b) \rangle \langle \text{po}(c), \text{ne}(c) \rangle$$

$$(b) = \langle (\text{po}(a)\text{po}(b) + \text{ne}(a)\text{ne}(b))\text{po}(c) + (\text{po}(a)\text{ne}(b) + \text{ne}(a)\text{po}(b))\text{ne}(c), (\text{po}(a)\text{po}(b) + \text{ne}(a)\text{ne}(b))\text{ne}(c) + (\text{po}(a)\text{ne}(b) + \text{ne}(a)\text{po}(b))\text{po}(c) \rangle$$

$$(c) = \langle \text{po}(a)(\text{po}(b)\text{po}(c) + \text{ne}(b)\text{ne}(c)) + \text{ne}(a)(\text{po}(b)\text{ne}(c) + \text{ne}(b)\text{po}(c)), \text{po}(a)(\text{po}(b)\text{ne}(c) + \text{ne}(b)\text{po}(c)) + \text{ne}(a)(\text{po}(b)\text{po}(c) + \text{ne}(b)\text{ne}(c)) \rangle$$

$$(d) = \langle \text{po}(a), \text{ne}(a) \rangle \langle \text{po}(b)\text{po}(c) + \text{ne}(b)\text{ne}(c), \text{po}(b)\text{ne}(c) + \text{ne}(b)\text{po}(c) \rangle$$

$$(e) = a(bc).$$

## Procedure I:11(1.76)

### Objective

Choose two integers  $a, b$ . The objective of the following instructions is to show that  $ab = ba$ .

### Implementation

1. Show that  $ab$

$$(a) = \langle \text{po}(a)\text{po}(b) + \text{ne}(a)\text{ne}(b), \text{po}(a)\text{ne}(b) + \text{ne}(a)\text{po}(b) \rangle$$

$$(b) = \langle \text{po}(b)\text{po}(a) + \text{ne}(b)\text{ne}(a), \text{po}(b)\text{ne}(a) + \text{ne}(b)\text{po}(a) \rangle$$

$$(c) = ba.$$

## Procedure I:12(1.77)

### Objective

Choose an integer  $a$ . The objective of the following instructions is to show that  $1a = a$ .

### Implementation

1. Show that  $1a$

$$(a) = \langle 1, 0 \rangle \langle \text{po}(a), \text{ne}(a) \rangle$$

$$(b) = \langle 1\text{po}(a) + 0\text{ne}(a), 1\text{ne}(a) + 0\text{po}(a) \rangle$$

$$(c) = \langle \text{po}(a), \text{ne}(a) \rangle$$

$$(d) = a.$$

### Procedure I:13(1.78)

#### Objective

Choose three integers  $a, b, c$ . The objective of the following instructions is to show that  $a(b + c) = ab + ac$ .

#### Implementation

1. Show that  $a(b + c)$ 
  - (a)  $= \langle \text{po}(a), \text{ne}(a) \rangle \langle \text{po}(b) + \text{po}(c), \text{ne}(b) + \text{ne}(c) \rangle$
  - (b)  $= \langle \text{po}(a)(\text{po}(b) + \text{po}(c)) + \text{ne}(a)(\text{ne}(b) + \text{ne}(c)), \text{po}(a)(\text{ne}(b) + \text{ne}(c)) + \text{ne}(a)(\text{po}(b) + \text{po}(c)) \rangle$
  - (c)  $= \langle (\text{po}(a)\text{po}(b) + \text{ne}(a)\text{ne}(b)) + (\text{po}(a)\text{po}(c) + \text{ne}(a)\text{ne}(c)), (\text{po}(a)\text{ne}(b) + \text{ne}(a)\text{po}(b)) + (\text{po}(a)\text{ne}(c) + \text{ne}(a)\text{po}(c)) \rangle$
  - (d)  $= \langle \text{po}(a)\text{po}(b) + \text{ne}(a)\text{ne}(b), \text{po}(a)\text{ne}(b) + \text{ne}(a)\text{po}(b) \rangle + \langle \text{po}(a)\text{po}(c) + \text{ne}(a)\text{ne}(c), \text{po}(a)\text{ne}(c) + \text{ne}(a)\text{po}(c) \rangle$
  - (e)  $= ab + ac$ .

### Procedure I:14(1.91)

#### Objective

Choose an integer  $a$ . The objective of the following instructions is to show that  $(-1)^{2a} = 1$  and  $(-1)^{2a+1} = -1$ .

#### Implementation

1. **Show that**  $(-1)^{2a+1} = (-1)^{2a}(-1) = 1(-1) = -1$ 
  - (a) given that  $(-1)^{2a} = ((-1)^2)^a = 1^a = 1$
  - (b) given that  $(-1)^2 = (-1)(-1) + 1 + (-1) = (-1)((-1) + 1) + 1 = (-1)0 + 1 = 1$ .

#### Declaration I:8

The phrase " $a < b$ ", where  $a, b$  are rational numbers, will be used as a shorthand for " $\text{po}(a) + \text{ne}(b) < \text{ne}(a) + \text{po}(b)$ ".

### Procedure I:15(1.79)

#### Objective

Choose four integers  $a, b, c, d$  such that  $a < b$ ,  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $c < d$ .

#### Implementation

1. Show that  $\text{po}(a) + \text{ne}(c) = \text{ne}(a) + \text{po}(c)$  using **declaration I:3**.
2. Show that  $\text{po}(b) + \text{ne}(d) = \text{ne}(b) + \text{po}(d)$  using **declaration I:3**.
3. Show that  $\text{po}(a) + \text{ne}(b) < \text{ne}(a) + \text{po}(b)$  using **declaration I:8**.
4. Hence show that  $\text{po}(c) + \text{ne}(d)$ 
  - (a)  $= (\text{ne}(a) + \text{po}(c)) + (\text{po}(b) + \text{ne}(d)) - \text{ne}(a) - \text{po}(b)$
  - (b)  $= (\text{po}(a) + \text{ne}(c)) + (\text{ne}(b) + \text{po}(d)) - \text{ne}(a) - \text{po}(b)$
  - (c)  $= (\text{po}(a) + \text{ne}(b)) + \text{ne}(c) + \text{po}(d) - \text{ne}(a) - \text{po}(b)$
  - (d)  $< (\text{ne}(a) + \text{po}(b)) + \text{ne}(c) + \text{po}(d) - \text{ne}(a) - \text{po}(b)$
  - (e)  $= \text{ne}(c) + \text{po}(d)$ .
5. **Hence show that**  $c < d$ .

### Procedure I:16(1.80)

#### Objective

Choose three integers  $a, b, c$  such that  $a < b$ . The objective of the following instructions is to show that  $a + c < b + c$ .

### Implementation

1. Show that  $\text{po}(a) + \text{ne}(b) < \text{ne}(a) + \text{po}(b)$  using **declaration I:8**.
2. Hence show that  $\text{po}(a + c) + \text{ne}(b + c)$   
(a)  $= \text{po}(a) + \text{po}(c) + \text{ne}(b) + \text{ne}(c)$   
(b)  $= (\text{po}(a) + \text{ne}(b)) + \text{po}(c) + \text{ne}(c)$   
(c)  $= (\text{ne}(a) + \text{po}(b)) + \text{po}(c) + \text{ne}(c)$   
(d)  $= \text{ne}(a) + \text{ne}(c) + \text{po}(b) + \text{po}(c)$   
(e)  $= \text{ne}(a + c) + \text{po}(b + c)$ .
3. **Hence show that  $a + c < b + c$ .**

### Procedure I:17(1.81)

#### Objective

Choose two integers  $a, b$  such that  $a < b$ . The objective of the following instructions is to show that  $a \neq b$  and  $b \not< a$ .

#### Implementation

1. Show that  $\text{po}(a) + \text{ne}(b) < \text{ne}(a) + \text{po}(b)$ .
2. **Hence show that  $a \neq b$  given that  $\text{po}(a) + \text{ne}(b) \neq \text{ne}(a) + \text{po}(b)$ .**
3. **Also show that  $b \not< a$  given that  $\text{ne}(a) + \text{po}(b) \not< \text{po}(a) + \text{ne}(b)$ .**

### Procedure I:18(1.82)

#### Objective

Choose two integers  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $a \not< b$  and  $b \not< a$ .

#### Implementation

Implementation is analogous to that of **procedure I:17**.

### Procedure I:19(1.83)

#### Objective

Choose two integers  $a, b$  such that  $a \neq b$ . The objective of the following instructions is to show that  $a < b$  or  $b < a$ .

#### Implementation

1. Show that  $\text{po}(a) + \text{ne}(b) \neq \text{ne}(a) + \text{po}(b)$ .
2. If  $\text{po}(a) + \text{ne}(b) < \text{ne}(a) + \text{po}(b)$ , then do the following:  
(a) **Show that  $a < b$ .**
3. Otherwise do the following:  
(a) **Show that  $b < a$  given that  $\text{ne}(a) + \text{po}(b) < \text{po}(a) + \text{ne}(b)$ .**

### Procedure I:20(1.84)

#### Objective

Choose two integers  $a, b$  such that  $a \not< b$ . The objective of the following instructions is to show that  $a = b$  or  $b < a$ .

#### Implementation

Implementation is analogous to that of **procedure I:19**.

### Procedure I:21(1.85)

#### Objective

Choose two integers  $a, b$  such that  $0 < a$  and  $0 < b$ . The objective of the following instructions is to show that  $0 < a + b$ .

#### Implementation

1. Show that  $\text{ne}(a) = \text{po}(0) + \text{ne}(a) < \text{ne}(0) + \text{po}(a) = \text{po}(a)$  using **declaration I:8**.
2. Show that  $\text{ne}(b) = \text{po}(0) + \text{ne}(b) < \text{ne}(0) + \text{po}(b) = \text{po}(b)$  using **declaration I:8**.

3. Hence show that  $0 < a + b$  given that  
 $\text{po}(0) + \text{ne}(a+b) = \text{ne}(a+b) = \text{ne}(a) + \text{ne}(b) <$   
 $\text{po}(a) + \text{po}(b) = \text{po}(a+b) = \text{ne}(0) + \text{po}(a+b).$

## Procedure I:22(1.86)

### Objective

Choose two integers  $a, b$  such that  $0 < a$  and  $0 < b$ .  
The objective of the following instructions is to show  
that  $0 < ab$ .

### Implementation

1. Show that  $\text{ne}(a) = \text{po}(0) + \text{ne}(a) < \text{ne}(0) + \text{po}(a) = \text{po}(a)$  using **declaration I:8**.
2. Hence show that  $0 < \text{po}(a) - \text{ne}(a)$ .
3. Show that  $\text{ne}(b) = \text{po}(0) + \text{ne}(b) < \text{ne}(0) + \text{po}(b) = \text{po}(b)$  using **declaration I:8**.
4. Hence show that  $0 < \text{po}(b) - \text{ne}(b)$ .
5. Hence show that  $0 < ab$ 
  - (a) given that  $\text{po}(0) + \text{ne}(ab) = \text{ne}(a)\text{po}(b) + \text{po}(a)\text{ne}(b) < \text{po}(a)\text{po}(b) + \text{ne}(a)\text{ne}(b) = \text{ne}(0) + \text{po}(ab)$
  - (b) given that  $\text{ne}(a)(\text{po}(b) - \text{ne}(b)) < \text{po}(a)(\text{po}(b) - \text{ne}(b))$
  - (c) given that  $0 < (\text{po}(a) - \text{ne}(a))(\text{po}(b) - \text{ne}(b))$ .

### Declaration I:9

The notation  $\|a\|$  will be used as a shorthand for the following expression:

1.  $-a$  if  $a < 0$
2.  $a$  if  $a \geq 0$

## Procedure I:23(1.87)

### Objective

Choose two integers  $a, b$ . The objective of the following instructions is to show that  $\|ab\| = \|a\|\|b\|$ .

### Implementation

1. If  $a \geq 0$  and  $b \geq 0$ , then do the following:
  - (a) Show that  $\|ab\| = ab = \|a\|\|b\|$  given that  $ab \geq 0$ .
2. Otherwise if  $a < 0$  and  $b \geq 0$ , then do the following:
  - (a) Show that  $\|ab\| = -(ab) = (-a)b = \|a\|\|b\|$  given that  $ab < 0$ .
3. Otherwise if  $a \geq 0$  and  $b < 0$ , then do the following:
  - (a) Show that  $\|ab\| = -(ab) = a(-b) = \|a\|\|b\|$  given that  $ab < 0$ .
4. Otherwise do the following:
  - (a) Show that  $\|ab\| = ab = (-a)(-b) = \|a\|\|b\|$ .
    - i. given that  $ab > 0$
    - ii. given that  $a < 0$  and  $b < 0$ .

## Procedure I:24(1.88)

### Objective

Choose two integers  $a, b$ . The objective of the following instructions is to show that  $\|a+b\| \leq \|a\| + \|b\|$ .

### Implementation

1. If  $a + b \geq 0$ , then do the following:
  - (a) Show that  $\|a+b\| = a+b \leq \|a\| + \|b\|$ 
    - i. given that  $a \leq \|a\|$
    - ii. and  $b \leq \|b\|$ .
2. Otherwise do the following:
  - (a) Show that  $\|a+b\| = -(a+b) = (-a) + (-b) \leq \|a\| + \|b\|$ 
    - i. given that  $-a \leq \|a\|$
    - ii. and  $-b \leq \|b\|$
    - iii. and  $a+b < 0$ .

## Procedure I:25(1.89)

### Objective

Choose two integers  $a, b$ . The objective of the following instructions is to show that  $\|a\| - \|b\| \leq \|a - b\|$ .

### Implementation

1. Show that  $\|a\| = \|b + (a - b)\| \leq \|b\| + \|a - b\|$  using **procedure I:24**.
2. **Hence show that**  $\|a\| - \|b\| \leq \|a - b\|$ .

### Declaration I:10

The notation **sgn( $a$ )** will be used as a shorthand for the following expression:

1.  $-1$  if  $a < 0$
2.  $0$  if  $a = 0$
3.  $1$  if  $a > 0$

## Procedure I:26(1.90)

### Objective

Choose an integer  $a$ . The objective of the following instructions is to show that  $a = \text{sgn}(a)\|a\|$ .

### Implementation

1. If  $a > 0$ , then do the following:
  - (a) **Show that**  $a = 1a = \text{sgn}(a)\|a\|$ 
    - i. given that  $\|a\| = a$
    - ii. and  $\text{sgn}(a) = 1$ .
2. If  $a = 0$ , then do the following:
  - (a) **Show that**  $a = 0 = \text{sgn}(a)0 = \text{sgn}(a)\|a\|$   
**given that**  $\|a\| = a = 0$ .
3. Otherwise if  $a < 0$ , then do the following:
  - (a) **Show that**  $a = (-1)(-a) = \text{sgn}(a)\|a\|$ 
    - i. given that  $\|a\| = -a$
    - ii. and  $\text{sgn}(a) = -1$ .



## Chapter 2

# Modular Arithmetic

### Procedure I:27(1.00)

#### Objective

Choose an integer  $a$  and a positive integer  $b$ . The objective of the following instructions is to construct integers  $n$  and  $m$  such that  $a = nb + m$  and  $0 \leq m < b$ .

#### Implementation

1. Let  $n = 0$ .
2. While  $(n + 1)b \leq a$ , do the following:
  - (a) Let  $n$  receive  $n + 1$ .
  - (b) Show that  $nb \leq a$ .
3. While  $nb > a$ , do the following:
  - (a) Let  $n$  receive  $n - 1$ .
  - (b) Show that  $(n + 1)b > a$ .
4. Hence show that  $nb \leq a$  and  $(n + 1)b > a$ .
5. Let  $m = a - nb$ .
6. **Now show that**  $b > a - nb = m \geq 0$  **and**  
 $a = bn + a - nb = nb + m$ .
7. **Yield**  $\langle n, m \rangle$ .

#### Declaration I:11

The notation  $a \text{ div } b$  will be used to refer to the first part of the pair yielded by executing **procedure I:27** on  $\langle a, b \rangle$ .

#### Declaration I:12

The notation  $a \bmod b$  will be used to refer to the second part of the pair yielded by executing **procedure I:27** on  $\langle a, b \rangle$ .

#### Declaration I:13

The notation  $a \equiv b \pmod{c}$  will be used as a shorthand for " $a \bmod c = b \bmod c$ ".

### Procedure I:28(1.01)

#### Objective

Choose four integers  $a, b, c, d$  and a positive integer  $e$  in such a way that  $a \equiv c \pmod{e}$  and  $b \equiv d \pmod{e}$ . The objective of the following instructions is to show that  $a + b \equiv c + d \pmod{e}$ .

#### Implementation

1. Show that  $a + b$ 
  - (a)  $\equiv (a \text{ div } e)e + (a \bmod e) + (b \text{ div } e)e + (b \bmod e)$
  - (b)  $\equiv (a \bmod e) + (b \bmod e)$
  - (c)  $\equiv (c \bmod e) + (d \bmod e)$
  - (d)  $\equiv (c \text{ div } e)e + (c \bmod e) + (d \text{ div } e)e + (d \bmod e)$
  - (e)  $\equiv c + d \pmod{e}$ .

### Procedure I:29(1.02)

#### Objective

Choose four integers  $a, b, c, d$  and a positive integer  $e$  in such a way that  $a \equiv c \pmod{e}$  and  $b \equiv d \pmod{e}$ . The objective of the following instructions is to show that  $ab \equiv cd \pmod{e}$ .

#### Implementation

1. Show that  $ab$

- (a)  $\equiv ((a \operatorname{div} e)e + (a \operatorname{mod} e))((b \operatorname{div} e)e + (b \operatorname{mod} e))$
- (b)  $\equiv (a \operatorname{div} e)(b \operatorname{div} e)e^2 + (a \operatorname{div} e)(b \operatorname{mod} e)e + (a \operatorname{mod} e)(b \operatorname{div} e)e + (a \operatorname{mod} e)(b \operatorname{mod} e)$
- (c)  $\equiv (a \operatorname{mod} e)(b \operatorname{mod} e)$
- (d)  $\equiv (c \operatorname{mod} e)(d \operatorname{mod} e)$
- (e)  $\equiv (c \operatorname{div} e)(d \operatorname{div} e)e^2 + (c \operatorname{div} e)(d \operatorname{mod} e)e + (c \operatorname{mod} e)(d \operatorname{div} e)e + (c \operatorname{mod} e)(d \operatorname{mod} e)$
- (f)  $\equiv cd \pmod{e}$ .

### Procedure I:30(1.03)

#### Objective

Choose an integer  $a$  and two positive integers  $b, c$ . The objective of the following instructions is to show that  $(a \operatorname{mod} bc) \operatorname{mod} b = a \operatorname{mod} b$ .

#### Implementation

- 1. **Show that**  $(a \operatorname{mod} bc) \operatorname{mod} b = (a - (a \operatorname{div} bc)bc) \operatorname{mod} b = a \operatorname{mod} b$ .

### Procedure I:31(1.04)

#### Objective

Choose a positive integer  $a$  and four integers  $b_1, b_0, c_1, c_0$  such that  $0 \leq b_0 < a$ ,  $0 \leq c_0 < a$ , and  $b_1a + b_0 = c_1a + c_0$ . The objective of the following instructions is to show that  $b_1 = c_1$  and  $b_0 = c_0$ .

#### Implementation

- 1. **Show that**  $b_0 = b_0 \operatorname{mod} a = (b_1a + b_0) \operatorname{mod} a = (c_1a + c_0) \operatorname{mod} a = c_0 \operatorname{mod} a = c_0$ .
- 2. **Therefore show that**  $b_1 = c_1$  **given that**  $b_1a = c_1a$ .

### Procedure I:32(1.05)

#### Objective

Choose an integer  $a$  and two positive integers  $b, c$ . The objective of the following instructions is to show that  $ca \operatorname{mod} cb = c(a \operatorname{mod} b)$  and that  $ca \operatorname{div} cb = a \operatorname{div} b$ .

#### Implementation

- 1. Show that  $bc(a \operatorname{div} b) + c(a \operatorname{mod} b) = c(b(a \operatorname{div} b) + a \operatorname{mod} b) = ca = cb(ca \operatorname{div} cb) + ca \operatorname{mod} cb$ .
- 2. Show that  $0 \leq a \operatorname{mod} b < b$ .
- 3. Show that  $0 \leq c(a \operatorname{mod} b) < cb$ .
- 4. Show that  $0 \leq ca \operatorname{mod} cb < cb$ .
- 5. **Hence show that**  $c(a \operatorname{mod} b) = ca \operatorname{mod} cb$  **and**  $a \operatorname{div} b = ca \operatorname{div} cb$  **using procedure I:31.**

### Procedure I:33(1.06)

#### Objective

Choose two integers  $a, b$  and a positive integer  $c$  such that  $a \operatorname{mod} c + b \operatorname{mod} c < c$ . The objective of the following instructions is to show that  $a \operatorname{div} c + b \operatorname{div} c = (a + b) \operatorname{div} c$  and  $a \operatorname{mod} c + b \operatorname{mod} c = (a + b) \operatorname{mod} c$ .

#### Implementation

- 1. Show that  $a = c(a \operatorname{div} c) + a \operatorname{mod} c$ .
- 2. Show that  $b = c(b \operatorname{div} c) + b \operatorname{mod} c$ .
- 3. Therefore show that  $a + b = c(a \operatorname{div} c + b \operatorname{div} c) + (a \operatorname{mod} c + b \operatorname{mod} c)$ .
- 4. Show that  $0 \leq a \operatorname{mod} c + b \operatorname{mod} c < c$ .

5. Also show that  $a + b = ((a + b) \text{ div } c)c + (a + b) \bmod c$ .
6. Show that  $0 \leq (a + b) \bmod c < c$ .
7. **Hence show that**  $a \text{ div } c + b \text{ div } c = (a + b) \text{ div } c$  **and**  $a \bmod c + b \bmod c = (a + b) \bmod c$  **using procedure I:31.**

### Procedure I:34(1.07)

#### Objective

Choose two integers  $a, b$  and a positive integer  $c$  such that  $a \bmod c + b \bmod c \geq c$ . The objective of the following instructions is to show that  $1 + a \text{ div } c + b \text{ div } c = (a + b) \text{ div } c$  and  $a \bmod c + b \bmod c - c = (a + b) \bmod c$ .

#### Implementation

1. Show that  $a = c(a \text{ div } c) + a \bmod c$ .
2. Show that  $b = c(b \text{ div } c) + b \bmod c$ .
3. Therefore show that  $a + b = c(a \text{ div } c + b \text{ div } c) + a \bmod c + b \bmod c = c(1 + a \text{ div } c + b \text{ div } c) + (a \bmod c + b \bmod c - c)$ .
4. Show that  $c \leq a \bmod c + b \bmod c < 2c$ .
5. Therefore show that  $0 \leq a \bmod c + b \bmod c - c < c$ .
6. Also show that  $a + b = c((a + b) \text{ div } c) + (a + b) \bmod c$ .
7. Show that  $0 \leq (a + b) \bmod c < c$ .
8. **Therefore show that**  $1 + a \text{ div } c + b \text{ div } c = (a + b) \text{ div } c$  **and**  $a \bmod c + b \bmod c - c = (a + b) \bmod c$  **using procedure I:31.**

### Procedure I:35(1.08)

#### Objective

Choose an integer  $a$  and two positive integers  $b, c$ . The objective of the following instructions is to show that  $a \text{ div } bc = (a \text{ div } b) \text{ div } c$  and  $a \bmod bc = ((a \text{ div } b) \bmod c)b + a \bmod b$ .

#### Implementation

1. Show that  $a = (((a \text{ div } b) \text{ div } c)c + (a \text{ div } b) \bmod c)b + a \bmod b = ((a \text{ div } b) \text{ div } c)bc + ((a \text{ div } b) \bmod c)b + a \bmod b$ 
  - (a) given that  $a = (a \text{ div } b)b + a \bmod b$
  - (b) given that  $a \text{ div } b = ((a \text{ div } b) \text{ div } c)c + (a \text{ div } b) \bmod c$ .
2. Show that  $0 \leq ((a \text{ div } b) \bmod c)b \leq cb - b$  given that  $0 \leq (a \text{ div } b) \bmod c \leq c - 1$ .
3. Therefore show that  $0 \leq ((a \text{ div } b) \bmod c)b + a \bmod b < cb$  given that  $0 \leq a \bmod b < b$ .
4. Now show that  $a = (a \text{ div } bc)bc + a \bmod bc$  and  $0 \leq a \bmod bc < bc$ .
5. **Therefore show that**  $(a \text{ div } b) \text{ div } c = a \text{ div } bc$  **and**  $((a \text{ div } b) \bmod c)b + a \bmod b = a \bmod bc$  **using procedure I:31.**

### Procedure I:36(1.09)

#### Objective

Choose an integer  $a$  and a non-negative integer  $b$ . The objective of the following instructions is to construct integers  $c, d, e, f, g$  such that  $a = cd$ ,  $b = ce$ ,  $fa + gb = c$ , and if  $b = 0$ , then  $c = |a|$ , otherwise  $0 < c \leq b$ .

#### Implementation

1. If  $b = 0$ , then do the following:
  - (a) **Show that**  $a = \text{sgn}(a)|a|$ .
  - (b) **Show that**  $b = 0|a|$ .
  - (c) **Show that**  $|a| = \text{sgn}(a)a + 0b$ .
  - (d) **Yield**  $\langle |a|, \text{sgn}(a), 0, \text{sgn}(a), 0 \rangle$ .
2. Otherwise do the following:
  - (a) Show that  $0 \leq a \bmod b < b$ .
  - (b) Use **procedure I:36** on  $\langle b, a \bmod b \rangle$  to construct  $\langle c, d, e, f, g \rangle$  and show that:
    - i.  $b = cd$
    - ii.  $a \bmod b = ce$

- iii.  $c = \|b\|$  if  $a \bmod b = 0$ , otherwise  $0 < c \leq a \bmod b$
- iv.  $fb + g(a \bmod b) = c$ .
- (c) **Hence show that**  $a = (a \operatorname{div} b)b + (a \bmod b) = c(d(a \operatorname{div} b) + e)$ .
- (d) **Also show that**  $(f - g(a \operatorname{div} b))b + ga = fb + g(a - (a \operatorname{div} b)b) = fb + g(a \bmod b) = c$ .
- (e) If  $a \bmod b = 0$ , then do the following:
  - i. **Show that**  $0 < b = c \leq b$  **given that**  $b \geq 0$ ,  $b \neq 0$ , **and**  $c = \|b\| = b$ .
- (f) Otherwise do the following:
  - i. **Show that**  $0 < c \leq a \bmod b < b$  **given**  $0 < c \leq a \bmod b$ .
- (g) **Therefore yield**  $\langle c, d(a \operatorname{div} b) + e, d, g, f - g(a \operatorname{div} b) \rangle$ .

#### Declaration I:14

The notation  $(a, b)$  will be used to refer to the first part of the quintuple constructed by using **procedure I:36** on the pair  $\langle a, b \rangle$ .

#### Procedure I:37(1.10)

##### Objective

Choose an integer  $a$  and a positive integer  $b$ . Let  $1 \leq c \leq b$  be the largest integer such that  $a \bmod c = 0$  and  $b \bmod c = 0$ . The objective of the following instructions is to either show that  $0 \neq 0$  or  $(a, b) = c$ .

##### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle d, e, f, g, h \rangle$  and show that:
  - (a)  $a = ed$
  - (b)  $b = fd$
  - (c)  $ga + hb = d$
  - (d)  $0 < d \leq b$ .
2. If  $d > c$ , then do the following:

- (a) Show that  $a \bmod d \neq 0$  or  $b \bmod d \neq 0$  given that  $0 < d \leq b$  is larger than the largest integer such that  $a \bmod c = 0$  and  $b \bmod c = 0$ .
- (b) If  $a \bmod d \neq 0$ , then do the following:
  - i. Show that  $a \bmod d = 0$  given that  $a = ed$ .
  - ii. **Hence show that**  $0 \neq 0$  **given that**  $a \bmod d \neq 0$  **and**  $a \bmod d = 0$ .
  - iii. **Abort procedure.**
- (c) Otherwise if  $b \bmod d \neq 0$ , then do the following:
  - i. Show that  $b \bmod d = 0$  given that  $b = fd$ .
  - ii. **Hence show that**  $0 \neq 0$  **given that**  $b \bmod d \neq 0$  **and**  $b \bmod d = 0$ .
  - iii. **Abort procedure.**
- 3. Otherwise if  $d < c$ , then do the following:
  - (a) Show that  $0 \equiv gc(a \operatorname{div} c) + hc(b \operatorname{div} c) = g(c(a \operatorname{div} c) + a \bmod c) + h(c(b \operatorname{div} c) + b \bmod c) = ga + hb = d \not\equiv 0 \pmod{c}$  given that:
    - i.  $ga + hb = d$
    - ii.  $a \bmod c = 0$
    - iii.  $b \bmod c = 0$ .
  - (b) **Hence show that**  $0 \neq 0$ .
  - (c) **Abort procedure.**
- 4. **Otherwise show that**  $(a, b) = d = c$ .

#### Procedure I:38(1.11)

##### Objective

Choose integers  $a, c, d, j$  and a non-negative integer  $b$ . Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle e, f, g, h, i \rangle$ . The objective of the following instructions is to show that  $ca + db = (c + gj)a + (d - fj)b$ .

##### Implementation

1. **Show that**  $(c + gj)a + (d - fj)b = ca + db + gja - fjb = ca + db + gje - fje = ca + db$ .

### Procedure I:39(1.12)

#### Objective

Choose integers  $a, c, d$  and a non-negative integer  $b$  such that  $ca + db = (a, b)$ . Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle e, f, g, h, i \rangle$ . The objective of the following instructions is to construct a  $j$  such that  $c = h + gj$  and  $d = i - fj$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to show that:
  - (a)  $a = ef$
  - (b)  $b = eg$
  - (c)  $ha + ib = e$ .
2. Show that  $cf + dg = 1$ 
  - (a) given that  $cef + deg = ca + db = (a, b) = e$
  - (b) given that  $a = ef$  and  $b = eg$ .
3. Show that  $hf + ig = 1$ 
  - (a) given that  $hef + ieg = ha + ib = e$
  - (b) given that  $a = ef$  and  $b = eg$ .
4. Let  $j = ci - hd$ .
5. **Show that**  $c = h + cig - hdg = h + g(ci - hd) = h + gj$ 
  - (a) given that  $c - cig = c(1 - ig) = chf = h(1 - dg) = h - hdg$
  - (b) given that  $cf = 1 - dg$ .
6. **Show that**  $d = i - icf + dhf = i - f(ic - dh) = i - fj$ 
  - (a) given that  $d - dhf = d(1 - hf) = dig = i(1 - cf) = i - icf$
  - (b) given that  $dg = 1 - cf$ .
7. **Yield**  $\langle j \rangle$ .

### Procedure I:40(1.13)

#### Objective

Choose an integer  $a$  and a positive integer  $b$  such that  $0 < (a, b) < b$ . The objective of the following

instructions is to show that  $0 \neq 0$  or  $a \bmod b \neq 0$ .

#### Implementation

1. If  $a \bmod b = 0$ , then do the following:
  - (a) Show that  $af \equiv 0f \equiv 0 \pmod{b}$  given that  $a \bmod b = 0$ .
  - (b) Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle c, d, e, f, g \rangle$  and show that:
    - i.  $fa + gb = c = (a, b)$
    - ii.  $0 < c = (a, b) \leq b$ .
  - (c) Hence show that  $fa \equiv (a, b) \not\equiv 0 \pmod{b}$  given that  $0 < (a, b) < b$ .
  - (d) Hence show that  $0 \neq 0$  given that  $0 \equiv af \not\equiv 0 \pmod{b}$ .
  - (e) **Abort procedure.**
2. **Otherwise show that**  $a \bmod b \neq 0$ .

### Procedure I:41(1.14)

#### Objective

Choose five integers  $a, d, e, f, g$  and two non-negative integers  $b, c$  such that  $a = cd$ ,  $b = ce$ , and  $fa + gb = c$ . The objective of the following instructions is to show that  $0 < 0$  or  $(a, b) = c$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle u, v, x, y, z \rangle$  and show that:
  - (a)  $u \geq 0$
  - (b)  $a = uv$
  - (c)  $b = xu$
  - (d)  $u = ya + zb$ .
2. Hence show that  $c = fa + gb = (fv + gx)u$ .
3. If  $u = 0$ , then do the following:
  - (a) **Show that**  $c = (fv + gx)u = 0 = u = (a, b)$ .
  - (b) **Yield.**
4. Show that  $u = ya + zb = (yd + ze)c$  given that  $u = ya + zb$ ,  $a = cd$ , and  $b = ce$ .

5. If  $c = 0$ , then do the following:
  - (a) **Show that**  $(a, b) = u = (yd + ze)c = 0 = c$ .
  - (b) **Yield.**
6. Show that  $fv + gx = yd + ze = \pm 1$ 
  - (a) given that  $(fv + gx)(yd + ze) = 1$
  - (b) given that  $c = (fv + gx)u = (fv + gx)(yd + ze)c$  and  $c > 0$ .
7. If  $fv + gx = yd + ze = -1$ , then do the following:
  - (a) Show that  $u = (yd + ze)c = (-1)c < 0$  given that  $u = (yd + ze)c$  and  $c > 0$ .
  - (b) **Hence show that**  $0 \leq u < 0$  **given that**  $u \geq 0$ .
  - (c) **Abort procedure.**
8. Otherwise, do the following:
  - (a) Show that  $fv + gx = yd + ze = 1$ .
  - (b) **Hence show that**  $c = (fv + gx)u = (1)u = (a, b)$  **given that**  $c = (fv + gx)u$ .

### Procedure I:42(1.15)

#### Objective

Choose an integer  $a$  and a non-negative integer  $b$ . The objective of the following instructions is to show that  $0 < 0$  or  $(a, b) = (-a, b)$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle c, d, e, f, g \rangle$  and show that:
  - (a)  $a = dc$
  - (b)  $b = ec$
  - (c)  $fa + gb = c$ .
2. Hence show that  $-a = (-d)c$ .
3. Also show that  $(-f)(-a) + gb = c$ .
4. Use **procedure I:41** on  $\langle -a, b, c, -d, e, -f, g \rangle$  to show that  $(-a, b) = c = (a, b)$ .

### Procedure I:43(1.16)

#### Objective

Choose two non-negative integers  $a, b$ . The objective of the following instructions is to show that  $0 < 0$  or  $(a, b) = (b, a)$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle c, d, e, f, g \rangle$  and show that:
  - (a)  $b = ec$
  - (b)  $a = dc$
  - (c)  $gb + fa = c$ .
2. Use **procedure I:41** on  $\langle b, a, c, e, d, g, f \rangle$  to show that  $(b, a) = c = (a, b)$ .

### Procedure I:44(1.17)

#### Objective

Choose two integers  $a, b$  and a positive integer  $c$  such that  $a \equiv b \pmod{c}$ . The objective of the following instructions is to show that  $0 < 0$  or  $(a, c) = (b, c)$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, c \rangle$  to construct  $\langle d, e, f, g, h \rangle$  and show that:
  - (a)  $a = ed$
  - (b)  $c = fd$
  - (c)  $ga + hc = d$ .
2. Let  $j = b \text{ div } c - a \text{ div } c$ .
3. Hence show that  $b = a + jc = ed + jfd = (e + jf)d$ .
4. Also show that  $gb + (h - gj)c = g(a + jc) + (h - gj)c = ga + hc = d$  given that  $b = a + jc$ .
5. Use **procedure I:41** on  $\langle b, c, d, e + jf, f, g, h - gj \rangle$  to show that  $(b, c) = d = (a, c)$ .

## Procedure I:45(1.18)

### Objective

Choose an integer  $a$  and two non-negative integers  $b, c$ . The objective of the following instructions is to show that either  $0 < 0$  or  $(ca, cb) = c(a, b)$ .

### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle d, e, f, g, h \rangle$  and show that:
  - (a)  $a = ed$
  - (b)  $b = df$
  - (c)  $ga + hb = d$ .
2. Hence show that  $ca = e(cd)$ ,  $cb = f(cd)$ , and  $g(ca) + h(cb) = cd$ .
3. Use **procedure I:41** on  $\langle ca, cb, cd, e, f, g, h \rangle$  to show that  $(ca, cb) = cd = c(a, b)$ .

## Procedure I:46(1.19)

### Objective

Choose an integer  $a$  and two non-negative integers  $b, c$ . The objective of the following instructions is to show that either  $0 < 0$  or  $(a, (b, c)) = ((a, b), c)$ .

### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle d_0, e_0, f_0, g_0, h_0 \rangle$  and show that:
  - (a)  $a = d_0e_0$
  - (b)  $b = d_0f_0$
  - (c)  $g_0a + h_0b = d_0$ .
2. Use **procedure I:36** on  $\langle b, c \rangle$  to construct  $\langle d_1, e_1, f_1, g_1, h_1 \rangle$  and show that:
  - (a)  $b = d_1e_1$
  - (b)  $c = d_1f_1$
  - (c)  $g_1b + h_1c = d_1$ .
3. Use **procedure I:36** on  $\langle (a, b), c \rangle$  to construct  $\langle d_2, e_2, f_2, g_2, h_2 \rangle$  and show that:

$$(a) \ (a, b) = d_2e_2$$

$$(b) \ c = d_2f_2$$

$$(c) \ g_2(a, b) + h_2c = d_2.$$

$$4. \text{ Show that } a = d_0e_0 = e_0(a, b) = e_0d_2e_2 = e_0e_2((a, b), c).$$

$$5. \text{ Also show that } (b, c)$$

$$(a) \ = g_1b + h_1c$$

$$(b) \ = g_1d_0f_0 + h_1d_2f_2$$

$$(c) \ = g_1f_0(a, b) + h_1f_2((a, b), c)$$

$$(d) \ = g_1f_0d_2e_2 + h_1f_2((a, b), c)$$

$$(e) \ = g_1f_0e_2((a, b), c) + h_1f_2((a, b), c)$$

$$(f) \ = (g_1f_0e_2 + h_1f_2)((a, b), c).$$

$$6. \text{ Also show that } ((a, b), c)$$

$$(a) \ = d_2$$

$$(b) \ = g_2(a, b) + h_2c$$

$$(c) \ = g_2d_0 + h_2d_1f_1$$

$$(d) \ = g_2(g_0a + h_0b) + h_2f_1(b, c)$$

$$(e) \ = g_2g_0a + g_2h_0d_1e_1 + h_2f_1(b, c)$$

$$(f) \ = g_2g_0a + g_2h_0e_1(b, c) + h_2f_1(b, c)$$

$$(g) \ = g_2g_0a + (g_2h_0e_1 + h_2f_1)(b, c).$$

$$7. \text{ Use **procedure I:41** on } \langle a, (b, c), ((a, b), c), e_0e_2, g_1f_0e_2 + h_1f_2, g_2g_0, g_2h_0e_1 + h_2f_1 \rangle \text{ to show that } ((a, b), c) = (a, (b, c)).$$

### Declaration I:15

The notation  $(a_0, a_1, \dots, a_{n-1})$  will be used to contextually refer to one of the following integers:

1.  $((a_0), (a_1, a_2, \dots, a_{n-1}))$
2.  $((a_0, a_1), (a_2, a_3, \dots, a_{n-1}))$
3.  $\vdots$
4.  $((a_0, a_1, \dots, a_{n-2}), (a_{n-1}))$

### Procedure I:47(1.20)

#### Objective

Choose two integers  $a, b$  and a non-negative integer  $c$  such that  $(a, c) = 1$  and  $(b, c) = 1$ . The objective of the following instructions is to show that either  $0 < 0$  or  $(ab, c) = 1$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, c \rangle$  to construct  $\langle d, e, f, g, h \rangle$  and show that  $ga + hc = d = (a, c) = 1$ .
2. Use **procedure I:36** on  $\langle b, c \rangle$  to construct  $\langle t, u, v, w, x \rangle$  and show that  $wb + xc = t = (b, c) = 1$ .
3. Hence show that  $(gw)(ab) + (gax + wbh + hxc)c = (ga + hc)(wb + xc) = 1$ .
4. Use **procedure I:41** on  $\langle ab, c, 1, ab, c, gw, gax + wbh + hxc \rangle$  to show that  $(ab, c) = 1$ .

### Procedure I:48(1.21)

#### Objective

Choose an integer  $a$  and two non-negative integers  $b, c$  such that  $(a, bc) = 1$ . The objective of the following instructions is to show that either  $0 < 0$  or  $(a, b) = 1$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, bc \rangle$  to construct  $\langle d, e, f, g, h \rangle$  and show that  $ga + (hc)b = ga + h(bc) = d = (a, bc) = 1$ .
2. Now use **procedure I:41** on  $\langle a, b, 1, a, b, g, hc \rangle$  to show that  $(a, b) = 1$ .

#### Declaration I:16

The phrase "**prime number**" will be used to refer to integers  $a$  such that  $a > 1$  and  $a \bmod k \neq 0$  for  $1 < k < a$ .

### Procedure I:49(1.22)

#### Objective

Choose an integer  $a$  and a prime  $b$  such that  $a \bmod b \neq 0$ . The objective of the following instructions is to show that either  $0 \neq 0$  or  $(a, b) = 1$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle c, d, e, f, g \rangle$  and show that:
  - (a)  $a = cd$
  - (b)  $b = ce$
  - (c)  $0 < c \leq b$ .
2. If  $c = b$ , then do the following:
  - (a) Show that  $a \bmod b = 0$  given that  $a = cd = bd$ .
  - (b) **Hence show that  $0 \neq 0$  given that  $a \bmod b \neq 0$ .**
  - (c) **Abort procedure.**
3. Otherwise if  $1 < c < b$ , then do the following:
  - (a) Show that  $b \bmod c = 0$  given that  $b = ce$ .
  - (b) **Hence show that  $0 \neq 0$  given that  $b$  is prime.**
  - (c) **Abort procedure.**
4. Otherwise, do the following:
  - (a) **Show that  $(a, b) = c = 1$ .**

### Procedure I:50(1.23)

#### Objective

Choose two integers  $a, b$  and a prime  $c$  such that  $a \bmod c \neq 0$  and  $b \bmod c \neq 0$ . The objective of the following instructions is to show that either  $0 \neq 0$  or  $ab \bmod c \neq 0$ .



## Implementation

1. Use **procedure I:49** on  $\langle a, c \rangle$  to show that  $(a, c) = 1$ .
2. Use **procedure I:49** on  $\langle b, c \rangle$  to show that  $(b, c) = 1$ .
3. Use **procedure I:47** on  $\langle a, b, c \rangle$  to show that  $0 < (ab, c) = 1 < c$ .
4. Use **procedure I:40** on  $\langle ab, c \rangle$  to show that  $ab \bmod c \neq 0$ .

## Declaration I:17

The notation  $|a|$  will be used to refer to the number of items in the list  $a$ .

## Declaration I:18

The notation  $a \frown b$  will be used to refer to the list formed by concatenating  $a$  and  $b$ .

## Declaration I:19

The notation  $f(R)$ , where  $R$  is a list and  $f[r]$  is a function of  $r$ , will contextually be used as a shorthand for the list  $\langle f(R_0), f(R_1), \dots, f(R_{|R|-1}) \rangle$ .

## Declaration I:20

The notation  $a_*$ , where  $a$  is a list, will be used as a shorthand for 1 if  $a$  is empty, otherwise it will be a shorthand for the product of the entries of  $a$ .

## Declaration I:21

The notation  $\prod_r^R f(r)$ , where  $R$  is a list and  $f[r]$  is a function of  $r$ , will be used as a shorthand for  $f(R)_*$ .

## Procedure I:51(1.24)

### Objective

Choose a positive integer  $a$ . The objective of the following instructions is to construct a list of prime numbers  $b$  such that  $a = b_*$ .

## Implementation

1. If  $a = 1$ , then do the following:
  - (a) Show that  $a = 1 = \langle \rangle_*$ .
  - (b) Hence yield  $\langle \rangle$ .
2. Otherwise, do the following:
  - (a) Show that  $a > 1$ .
  - (b) If there is a  $c \in [2 : a]$  such that  $a \bmod c = 0$ , then do the following:
    - i. Show that  $a = (a \operatorname{div} c)c$ .
    - ii. Hence show that  $1 < a \operatorname{div} c < a$ .
    - iii. Use **procedure I:51** on  $\langle a \operatorname{div} c \rangle$  to construct  $\langle d \rangle$  and show that:
      - A. every element of  $d$  is prime.
      - B.  $a \operatorname{div} c = d_*$ .
    - iv. Hence show that  $d$  is non-empty given that  $1 < a \operatorname{div} c = d_*$ .
    - v. Use **procedure I:51** on  $\langle c \rangle$  to construct  $\langle e \rangle$  and show that:
      - A. every element of  $e$  is prime.
      - B.  $c = e_*$ .
    - vi. Hence show that  $e$  is non-empty given that  $1 < c = e_*$ .
    - vii. Hence show that  $d \frown e$  is a non-empty list of prime numbers such that  $a = (a \operatorname{div} c)c = d_*e_* = (d \frown e)_*$ .
    - viii. Yield  $\langle d \frown e \rangle$ .
  - (c) Otherwise do the following:
    - i. Show that  $a$  is prime.
    - ii. Yield  $\langle a \rangle$ .

## Procedure I:52(1.25)

### Objective

Choose a prime  $a$  and a list of primes  $b$  such that  $b_* \equiv 0 \pmod{a}$ . The objective of the following instructions is to either show that  $0 = 1$  or to construct a  $k$  such that  $a = b_k$ .

## Implementation

1. Show that  $a > 1$  given that  $a$  is prime.
2. If  $|b| = 0$ , then do the following:
  - (a) Show that  $1 = b_* \equiv 0 \pmod{a}$ .
  - (b) **Hence show that  $0 = 1$  given that  $a > 1$ .**
  - (c) **Abort procedure.**
3. Otherwise if  $0 \not\equiv b \pmod{a}$ , then do the following:
  - (a) Show that  $b_* \not\equiv 0 \pmod{a}$  using **procedure I:50**.
  - (b) **Hence show that  $0 \neq 0$  given that  $b_* \equiv 0 \pmod{a}$ .**
  - (c) **Abort procedure.**
4. Otherwise do the following:
  - (a) Let  $k$  be such that  $b_k \pmod{a} = 0$ .
  - (b) Show that  $b_k = (b_k \text{ div } a)a$ .
  - (c) Hence show that  $b_k \text{ div } a \geq 1$ .
  - (d) If  $b_k \text{ div } a > 1$ , then do the following:
    - i. Show that  $1 < a < b_k$  given that:
      - A.  $a > 1$
      - B.  $b_k \text{ div } a > 1$
      - C.  $b_k = (b_k \text{ div } a)a$ .
    - ii. Hence show that  $b_k \pmod{a} \neq 0$  given that  $b_k$  is prime and  $1 < a < b_k$ .
    - iii. **Hence show that  $0 \neq b_k \pmod{a} = 0$  given that  $b_k \pmod{a} = 0$ .**
    - iv. **Abort procedure.**
  - (e) Otherwise do the following:
    - i. **Show that  $b_k = a$  given that  $b_k \text{ div } a = 1$ .**
    - ii. **Yield  $\langle k \rangle$ .**

## Declaration I:22

The notation  $[a : b]$  will be used as a shorthand for the list:

1.  $\langle a, a + 1, \dots, b - 1 \rangle$ , if  $b > a$

2.  $\langle \rangle$ , if  $b = a$
3.  $\langle a - 1, a - 2, \dots, b \rangle$ , if  $b < a$

## Procedure I:53(1.26)

### Objective

Choose two lists of primes  $a, b$  such that  $a_* = b_*$ . The objective of the following instructions is to show that either  $1 > 1$  or  $a$  is included in  $b$ .

### Implementation

1. If  $|a| = 0$ , then do the following:
  - (a) **Show that  $a$  is included in  $b$ .**
2. Otherwise, do the following:
  - (a) Show that  $|a| > 0$ .
  - (b) Show that  $b_* \equiv a_* \equiv 0 \pmod{a_0}$ .
  - (c) Use **procedure I:52** on  $\langle a_0, b \rangle$  to construct  $\langle k \rangle$  and show that  $b_k = a_0$ .
  - (d) Now show that  $(a_{[1:|a|]})_* = (b_{[0:k] \cap [k+1:|b|]})_*$ .
  - (e) Now use **procedure I:53** on  $\langle a_{[1:|a|]}, b_{[0:k] \cap [k+1:|b|]} \rangle$  to show that  $a_{[1:|a|]}$  is included in  $b_{[0:k] \cap [k+1:|b|]}$ .
  - (f) **Hence show that  $a$  is included in  $b$ .**

## Procedure I:54(1.27)

### Objective

Choose two lists of primes  $a, b$  such that  $a_* = b_*$ . The objective of the following instructions is to show that either  $1 > 1$  or  $a$  is a rearrangement of  $b$ .

### Implementation

1. Use **procedure I:53** on  $\langle a, b \rangle$  to show that  $a$  is included in  $b$ .
2. Use **procedure I:53** on  $\langle b, a \rangle$  to show that  $b$  is included in  $a$ .
3. **Hence show that  $a$  is a rearrangement of  $b$ .**

## Procedure I:55(1.28)

### Objective

Choose a positive integer  $a$ . The objective of the following instructions is to either show that  $0 = 1$  or to construct a prime  $b$  such that  $b > a$  and  $[a+1 : b]$  does not contain a prime.

### Implementation

1. Show that  $a! + 1 > 1$ .
2. Use **procedure I:51** on  $\langle a! + 1 \rangle$  to construct  $\langle d \rangle$  and show that:
  - (a)  $a! + 1 = d_*$
  - (b) every element of  $d$  is prime.
3. Hence show that  $|d| > 0$  given that  $a! + 1 > 1$ .
4. Hence show that  $(a! + 1) \bmod d_0 = 0$ .
5. If  $d_0 \in [2 : a+1]$ , then do the following:
  - (a) Show that  $a! + 1 \equiv 1 \pmod{d_0}$ 
    - i. given that  $a! \pmod{d_0} \equiv 0$
    - ii. given that  $d_0 \in [2 : a+1]$ .
  - (b) Show that  $0 \equiv a! + 1 \pmod{d_0}$ 
    - i. given that  $(a! + 1) \bmod d_0 = 0$
    - ii. given that  $a! + 1 = d_*$ .
  - (c) **Hence show that  $0 = 1$ .**
  - (d) **Abort procedure.**
6. Otherwise do the following:
  - (a) **Show that  $d_0$  is prime given that every element of  $d$  is prime.**
  - (b) **Hence show that  $d_0 > a$  given that  $d_0 > 1$  and  $d_0 \notin [2 : a+1]$ .**
  - (c) **Let  $b$  be the least prime in  $[a+1 : d_0+1]$ .**
  - (d) **Yield  $\langle b \rangle$ .**

## Procedure I:56(1.29)

### Objective

Choose a positive integer  $a$ . The objective of the following instructions is to construct a positive integer  $b$  such that  $[b+1 : b+a]$  does not contain a prime.

### Implementation

1. Let  $b = a! + 1$ .
2. For  $i \in [1 : a]$ , do the following:
  - (a) Show that  $b + i = a! + 1 + i = i!(i+1)(i+2) \cdots (a+1+i) = (1+i)(i!)(i+2)(i+3) \cdots (a+1)$ .
  - (b) Therefore show that  $b + i \equiv 0 \pmod{i+1}$ .
  - (c) Also show that  $b + i = a! + 1 + i > a! \geq a \geq i+1 > 1$ .
  - (d) **Hence show that  $b + i$  is not prime.**
3. **Yield  $\langle b \rangle$ .**

## Procedure I:57(1.30)

### Objective

Choose two lists of primes  $a, b$  in such a way that their intersection is empty. The objective of the following instructions is to show that  $0 = 1$  or  $(a_*, b_*) = 1$ .

### Implementation

1. Use **procedure I:36** on  $\langle a_*, b_* \rangle$  to construct  $\langle c, d, e, f, g \rangle$  and show that:
  - (a)  $0 < c \leq b_*$
  - (b)  $a_* = cd$
  - (c)  $b_* = ce$ .
2. If  $c > 1$ , then do the following:
  - (a) Use **procedure I:51** on  $\langle c \rangle$  to construct  $\langle h \rangle$  and show that  $c = h_*$ .
  - (b) Hence show that  $|h| > 0$  given that  $h_* = c > 1$ .

- (c) Now show that  $a_* = dc = dh_* = dh_0(h_{[1:|h|]})_* \equiv 0 \pmod{h_0}$ .
  - (d) Use **procedure I:52** on  $\langle h_0, a \rangle$  to construct  $\langle k \rangle$  and show that  $h_0 = a_k$ .
  - (e) Now show that  $b_* = ec = eh_* = eh_0(h_{[1:|h|]})_* \equiv 0 \pmod{h_0}$ .
  - (f) Use **procedure I:52** on  $\langle h_0, b \rangle$  to construct  $\langle m \rangle$  and show that  $h_0 = b_m$ .
  - (g) **Hence show that  $a$  and  $b$  intersect given that  $a_k = h_0 = b_m$ .**
  - (h) **Abort procedure.**
3. Otherwise do the following:
- (a) **Show that  $(a_*, b_*) = c = 1$  given that  $0 < c \leq b_*$  and  $c \leq 1$ .**

### Procedure I:58(1.31)

#### Objective

Choose two lists of primes  $a, b$ . Let  $c$  be the common sublist with multiplicity of  $a$  and  $b$ . The objective of the following instructions is to show that either  $0 < 0$  or  $(a_*, b_*) = c_*$ .

#### Implementation

1. Let  $d$  be the result of removing with multiplicity elements of  $c$  from  $a$ .
2. Show that  $a_* = c_*d_*$ .
3. Let  $e$  be the result of removing with multiplicity elements of  $c$  from  $b$ .
4. Show that  $b_* = c_*e_*$ .
5. Show that  $d$  and  $e$  share no common elements.
6. **Therefore show that  $(a_*, b_*) = (c_*d_*, c_*e_*) = c_*(d_*, e_*) = c_*$  using **procedure I:45** and **procedure I:57**.**

### Procedure I:59(1.32)

#### Objective

Choose an integer  $a$  and a positive integer  $b$ . The objective of the following instructions is to construct

integers  $c, f, e$  such that  $c = af$ ,  $c = be$ ,  $c(a, b) = ab$ , and  $|a| \leq |c| \leq |a|b$ .

#### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle d, e, f, g, h \rangle$  and show that:
  - (a)  $a = de$
  - (b)  $b = df$
  - (c)  $d > 0$ .
2. **Let  $c = af$ .**
3. **Show that  $c = af = def = be$ .**
4. **Show that  $c(a, b) = cd = afd = ab$ .**
5. Show that  $1 \leq f \leq b$ 
  - (a) given that  $0 < b = df$
  - (b) and  $d > 0$ .
6. Therefore show that  $|a| \leq |a|f \leq |a|b$ .
7. **Therefore show that  $|a| \leq |c| \leq |a|b$ .**
8. **Yield the tuple  $\langle c, f, e \rangle$ .**

#### Declaration I:23

The notation  $[a, b]$  will be used to refer to the first part of the triple yielded by executing **procedure I:59** on  $\langle a, b \rangle$ .

### Procedure I:60(1.33)

#### Objective

Choose two positive integers  $a, b$ . The objective of the following instructions is to show that either  $0 < 0$  or  $[a, b] = [b, a]$ .

#### Implementation

1. Show that  $(a, b) > 0$ .
2. Show that  $[a, b](a, b) = ab = ba = [b, a](b, a) = [b, a](a, b)$  using **procedure I:43**.
3. **Therefore show that  $[a, b] = [b, a]$ .**

### Procedure I:61(1.34)

#### Objective

Choose an integer  $a$  and two positive integers  $b, c$ . The objective of the following instructions is to show that either  $0 < 0$  or  $[ca, cb] = c[a, b]$ .

#### Implementation

1. Show that  $(ca, cb) > 0$ .
2. Show that  $[ca, cb](ca, cb) = cacb = c^2ab = c^2[a, b](a, b) = c[a, b](ca, cb)$  using **procedure I:45**.
3. **Therefore show that**  $[ca, cb] = c[a, b]$ .

### Procedure I:62(1.35)

#### Objective

Choose an integer  $a$  and two positive integers  $b, c$ . The objective of the following instructions is to show that either  $0 < 0$  or  $[[a, b], c] = [a, [b, c]]$ .

#### Implementation

1. Using **procedure I:46**, show that  $(a, b)(ab, (ac, bc))(b, c)[[a, b], c]$ 
  - (a)  $= (ab, (ac, bc))(b, c)[(a, b)[a, b], (a, b)c]$
  - (b)  $= (ab, (ac, bc))(b, c)[ab, (ac, bc)]$
  - (c)  $= ab(ac, bc)(b, c)$
  - (d)  $= abc(a, b)(b, c)$
  - (e)  $= bc(a, b)(ab, ac)$
  - (f)  $= (a, b)((ab, ac), bc)[(ab, ac), bc]$
  - (g)  $= (a, b)(ab, (ac, bc))[(ab, ac), bc]$
  - (h)  $= (a, b)(ab, (ac, bc))[a(b, c), [b, c](b, c)]$
  - (i)  $= (a, b)(ab, (ac, bc))(b, c)[a, [b, c]]$ .
2. Show that  $(a, b)(ab, (ac, bc))(b, c) > 0$ .
3. **Therefore show that**  $[[a, b], c] = [a, [b, c]]$ .

### Declaration I:24

The notation  $[a_0, a_1, \dots, a_{n-1}]$  will be used to contextually refer to one of the following integers:

1.  $[[a_0], [a_1, a_2, \dots, a_{n-1}]]$
2.  $[[a_0, a_1], [a_2, a_3, \dots, a_{n-1}]]$
3.  $\vdots$
4.  $[[a_0, a_1, \dots, a_{n-2}], [a_{n-1}]]$

### Procedure I:63(1.36)

#### Objective

Choose three positive integers  $a, b, c$ . The objective of the following instructions is to show that either  $0 < 0$  or  $((a, b), c) = [(a, c), (b, c)]$ .

#### Implementation

1. Using **procedure I:59**, **procedure I:45**, **procedure I:46**, **procedure I:43**, and **procedure I:37**, show that  $(a, b)((a, c), (b, c))([a, b], c)$ 
  - (a)  $= ((a, c), (b, c))((a, b)[a, b], (a, b)c)$
  - (b)  $= ((a, c), (b, c))(ab, (ac, bc))$
  - (c)  $= (a^2b, a^2c, c^2a, c^2b, b^2a, bac, b^2c)$
  - (d)  $= (a, b)(ab, ac, bc, c^2)$
  - (e)  $= (a, b)(a, c)(b, c)$
  - (f)  $= (a, b)((a, c), (b, c))[(a, c), (b, c)]$ .
2. Show that  $(a, b)((a, c), (b, c)) > 0$ .
3. **Therefore show that**  $[(a, b), c] = [(a, c), (b, c)]$ .

### Procedure I:64(1.37)

#### Objective

Choose three positive integers  $a, b, c$ . The objective of the following instructions is to show that either  $0 < 0$  or  $[(a, b), c] = ([a, c], [b, c])$ .

## Implementation

1. Using [procedure I:59](#), [procedure I:45](#), [procedure I:46](#), [procedure I:43](#), and [procedure I:37](#), show that  $((a, b), c)(a, c)(b, c)[(a, b), c]$ 
  - (a)  $= (a, c)(b, c)(a, b)c$
  - (b)  $= (ab, ac, cb, c^2)(a, b)c$
  - (c)  $= (a^2b, a^2c, ac^2, ab^2, abc, cb^2, bc^2)c$
  - (d)  $= (a, b, c)(ab, ac, bc)c$
  - (e)  $= ((a, b), c)(ac(b, c), bc(a, c))$
  - (f)  $= ((a, b), c)(a, c)(b, c)([a, c], [b, c])$ .
2. Show that  $((a, b), c)(a, c)(b, c) > 0$ .
3. **Therefore show that**  $[(a, b), c] = ([a, c], [b, c])$ .

## Chapter 3

# Congruence Equations

### Declaration I:25

The notation  $\chi_{b,d}(a, c)$ , where  $a, c$  are two integers and  $b, d$  are two positive integers such that  $a \equiv c \pmod{(b, d)}$ , will be used to refer to the result yielded by executing the following instructions:

1. Use **procedure I:36** on  $\langle b, d \rangle$  to construct  $\langle f, g, h, i, j \rangle$ .
2. **Yield the tuple**  $\langle (a + ((c - a) \text{div}(b, d))ib) \bmod [b, d] \rangle$ .

### Procedure I:65(1.39)

#### Objective

Choose three integers  $x, a, c$  and two positive integers  $b, d$  such that  $x \equiv a \pmod{b}$  and  $x \equiv c \pmod{d}$ . The objective of the following instructions is to show that  $0 \neq 0$  if  $a \not\equiv c \pmod{(b, d)}$ , otherwise  $x \equiv \chi_{b,d}(a, c) \pmod{[b, d]}$ .

#### Implementation

1. Use **procedure I:36** on  $\langle b, d \rangle$  to construct  $\langle e, f, g, h, i \rangle$  and show that:
  - (a)  $b = ef$
  - (b)  $d = eg$
  - (c)  $hb + id = e$ .
2. Let  $j = x \text{div } b - a \text{div } b$ .
3. Show that  $x = a + jb$  given that  $x \equiv a \pmod{b}$ .

4. Let  $k = x \text{div } d - c \text{div } d$ .
5. Show that  $x = c + kd$  given that  $x \equiv c \pmod{d}$ .
6. Therefore show that  $c - a = jb - kd$ .
7. If  $a \not\equiv c \pmod{(b, d)}$ , then do the following:
  - (a) Show that  $0 \neq c - a = jb - kd = jef - keg \equiv 0 \pmod{e}$ .
  - (b) **Therefore show that**  $0 \neq 0$ .
  - (c) **Abort procedure.**
8. Otherwise do the following:
  - (a) Let  $l = (c - a) \text{div}(b, d)$ .
  - (b) Show that  $l(b, d) = le = c - a = jb - kd = jef - keg$  given that  $c - a \equiv 0 \pmod{(b, d)}$ .
  - (c) Hence show that  $l \equiv jf \pmod{g}$  given that  $l = jf - kg$ .
  - (d) Hence show that  $fh \equiv 1 \pmod{g}$ 
    - i. given that  $fh + gi = 1$
    - ii. given that  $efh + egi = bh + di = e$
    - iii. given that  $b = ef, d = eg$ , and  $hb + id = e$ .
  - (e) Hence show that  $lh \equiv jfh \equiv j \pmod{g}$ 
    - i. given that  $l \equiv jf \pmod{g}$
    - ii. and  $fh \equiv 1 \pmod{g}$ .
  - (f) Hence show that  $lhb \equiv jb \pmod{bg = [b, d]}$  using **procedure I:32**.
  - (g) **Hence show that**  $x = a + jb \equiv a + lhb \equiv \chi_{b,d}(a, c) \pmod{[b, d]}$ .

## Procedure I:66(1.40)

### Objective

Choose two integers  $a, c$  and two positive integers  $b, d$  in such a way that  $a \equiv c \pmod{(b, d)}$ . The objective of the following instructions is to show that either  $0 < 0$  or  $\chi_{b,d}(a, c) = \chi_{d,b}(c, a)$ .

### Implementation

1. Use **procedure I:36** on  $\langle b, d \rangle$  to construct  $\langle f, g, h, i, j \rangle$  and show that  $ib + jd = f = (b, d)$ .
2. Use **procedure I:36** on  $\langle d, b \rangle$  to construct  $\langle k, l, m, n, p \rangle$  and show that  $pb + nd = k = (d, b) = (b, d)$ .
3. Use **procedure I:39** on  $\langle b, p, n, d \rangle$  to construct  $\langle q \rangle$  and show that  $n = j - qg$ .
4. Now using **procedure I:60**, show that  $\chi_{b,d}(a, c)$ 
  - (a)  $= (a + ((c - a) \operatorname{div}(b, d))ib) \bmod [b, d]$
  - (b)  $= (a + ((c - a) \operatorname{div}(b, d))(f - jd)) \bmod [b, d]$
  - (c)  $= (a + ((c - a) \operatorname{div}(b, d))f + ((a - c) \operatorname{div}(b, d))jd) \bmod [b, d]$
  - (d)  $= (a + (c - a) + ((a - c) \operatorname{div}(b, d))jd) \bmod [b, d]$
  - (e)  $= (c + ((a - c) \operatorname{div}(d, b))(n + qg)d) \bmod [b, d]$
  - (f)  $= (c + ((a - c) \operatorname{div}(d, b))dn + ((a - c) \operatorname{div}(d, b))q[b, d]) \bmod [b, d]$
  - (g)  $= (c + ((a - c) \operatorname{div}(d, b))dn) \bmod [b, d]$
  - (h)  $= (c + ((a - c) \operatorname{div}(d, b))dn) \bmod [d, b]$
  - (i)  $= \chi_{d,b}(c, a)$ .

## Procedure I:67(1.41)

### Objective

Choose three integers  $x, a, c$  and two positive integers  $b, d$  such that  $a \equiv c \pmod{(b, d)}$  and  $x \equiv \chi_{b,d}(a, c) \pmod{[b, d]}$ . The objective of the following instructions is to show that  $x \equiv a \pmod{b}$ .

### Implementation

1. Use **procedure I:36** on  $\langle b, d \rangle$  to construct  $\langle e, f, g, h, i \rangle$ .
2. Show that  $[b, d] = bg$ .
3. Hence show that  $(x \bmod (bg)) \bmod b = (\chi_{b,d}(a, c) \bmod (bg)) \bmod b$ 
  - (a) given that  $x \bmod (bg) = \chi_{b,d}(a, c) \bmod (bg)$
  - (b) given that  $x \bmod [b, d] = \chi_{b,d}(a, c) \bmod [b, d]$ .
4. **Therefore using procedure I:30, show that**  $x \bmod b = \chi_{b,d}(a, c) \bmod b = (a + ((c - a) \operatorname{div}(b, d))hb) \bmod b = a \bmod b$ .

## Procedure I:68(1.42)

### Objective

Choose three integers  $x, a, c$  and two positive integers  $b, d$  such that  $a \equiv c \pmod{(b, d)}$  and  $x \equiv \chi_{b,d}(a, c) \pmod{[b, d]}$ . The objective of the following instructions is to either show that  $0 < 0$  or to show that  $x \equiv a \pmod{b}$  and  $x \equiv c \pmod{d}$ .

### Implementation

1. Use **procedure I:67** on  $\langle x, a, c, b, d \rangle$  to show that  $x \equiv a \pmod{b}$ .
2. Show that  $x \equiv \chi_{b,d}(a, c) \equiv \chi_{d,b}(c, a) \pmod{[d, b]}$  using **procedure I:66**.
3. Use **procedure I:67** on  $\langle x, c, a, d, b \rangle$  to show that  $x \equiv c \pmod{d}$ .

## Procedure I:69(1.43)

### Objective

Choose two integers  $a, c$  and three positive integers  $b, d, e$  such that  $a \equiv c \pmod{(b, d)}$ . The objective of the following instructions is to show that  $\chi_{b,d}(ea, ec) = e\chi_{b,d}(a, c)$ .



## Implementation

1. Use **procedure I:68** on  $\langle \chi_{b,d}(a, c), a, c, b, d \rangle$  to show that:
  - (a)  $\chi_{b,d}(a, c) \equiv a \pmod{b}$
  - (b)  $\chi_{b,d}(a, c) \equiv c \pmod{d}$ .
2. Hence show that  $e\chi_{b,d}(a, c) \equiv ea \pmod{b}$  using **procedure I:32**.
3. Also show that  $e\chi_{b,d}(a, c) \equiv ec \pmod{d}$  using **procedure I:32**.
4. Also show that  $ea \equiv ec \pmod{(b, d)}$  using **procedure I:29** given that  $a \equiv c \pmod{(b, d)}$ .
5. **Hence show that**  $e\chi_{b,d}(a, c) \equiv \chi_{b,d}(ea, ec) \pmod{[b, d]}$  using **procedure I:65**.

## Procedure I:70(1.44)

### Objective

Choose two integers  $a, c$  and three positive integers  $b, d, e$  such that  $a \equiv c \pmod{(eb, ed)}$ . The objective of the following instructions is to show that  $\chi_{eb,ed}(a, c) \pmod{[b, d]} = \chi_{b,d}(a, c)$ .

### Implementation

1. Use **procedure I:68** on  $\langle \chi_{eb,ed}(a, c), a, c, eb, ed \rangle$  to show that:
  - (a)  $\chi_{eb,ed}(a, c) \equiv a \pmod{eb}$
  - (b)  $\chi_{eb,ed}(a, c) \equiv c \pmod{ed}$ .
2. Show that  $\chi_{eb,ed}(a, c) \equiv a \pmod{b}$  using **procedure I:30**.
3. Show that  $\chi_{eb,ed}(a, c) \equiv c \pmod{d}$  using **procedure I:30**.
4. Show that  $a \equiv c \pmod{(b, d)}$  using **procedure I:30** given that  $a \equiv c \pmod{e(b, d)}$ .
5. Hence show that  $\chi_{eb,ed}(a, c) \equiv \chi_{b,d}(a, c) \pmod{[b, d]}$  using **procedure I:65**.
6. **Hence show that**  $\chi_{eb,ed}(a, c) \pmod{[b, d]} = \chi_{b,d}(a, c)$ .

## Procedure I:71(1.46)

### Objective

Choose three integers  $a, c, e$  and three positive integers  $b, d, f$  such that  $a \equiv c \pmod{(b, d)}$  and  $\chi_{b,d}(a, c) \equiv e \pmod{([b, d], f)}$ . The objective of the following instructions is to show that  $0 \neq 0$  if  $c \not\equiv e \pmod{(d, f)}$  or  $a \not\equiv \chi_{d,f}(c, e) \pmod{(b, [d, f])}$ , otherwise  $\chi_{[b,d],f}(\chi_{b,d}(a, c), e) = \chi_{b,[d,f]}(a, \chi_{d,f}(c, e))$ .

### Implementation

1. Show that  $\chi_{[b,d],f}(\chi_{b,d}(a, c), e) \equiv e \pmod{f}$  using **procedure I:68**.
2. Show that  $\chi_{[b,d],f}(\chi_{b,d}(a, c), e) \equiv \chi_{b,d}(a, c) \pmod{[b, d] = gb = hd}$  using **procedure I:68**.
3. Show that  $\chi_{[b,d],f}(\chi_{b,d}(a, c), e) \equiv \chi_{b,d}(a, c) \equiv a \pmod{b}$  using **procedure I:30** and **procedure I:68**.
4. Show that  $\chi_{[b,d],f}(\chi_{b,d}(a, c), e) \equiv \chi_{b,d}(a, c) \equiv c \pmod{d}$  using **procedure I:30** and **procedure I:68**.
5. Use **procedure I:65** on  $\langle \chi_{[b,d],f}(\chi_{b,d}(a, c), e), c, e, d, f \rangle$  to show that  $0 \neq 0$  if  $c \not\equiv e \pmod{(d, f)}$ , otherwise  $\chi_{[b,d],f}(\chi_{b,d}(a, c), e) \equiv \chi_{d,f}(c, e) \pmod{[d, f]}$ .
6. Use **procedure I:65** on  $\langle \chi_{[b,d],f}(\chi_{b,d}(a, c), e), a, \chi_{d,f}(c, e), b, [d, f] \rangle$  to show that  $0 \neq 0$  if  $a \not\equiv \chi_{d,f}(c, e) \pmod{(b, [d, f])}$ , otherwise  $\chi_{[b,d],f}(\chi_{b,d}(a, c), e) \equiv \chi_{b,[d,f]}(a, \chi_{d,f}(c, e)) \pmod{[b, [d, f]] = [[b, d], f]}$ .
7. **Hence show that**  $\chi_{[b,d],f}(\chi_{b,d}(a, c), e) = \chi_{b,[d,f]}(a, \chi_{d,f}(c, e))$ .

### Declaration I:26

The notation  $\chi_{b_0, b_1, \dots, b_{n-1}}(a_0, a_1, \dots, a_{n-1})$  will be used to contextually refer to one of the following integers:

1.  $\chi_{b_0, [b_1, b_2, \dots, b_{n-1}]}(a_0, \chi_{b_1, b_2, \dots, b_{n-1}}(a_1, a_2, \dots, a_{n-1}))$
2.  $\chi_{[b_0, b_1], [b_2, b_3, \dots, b_{n-1}]}(\chi_{b_0, b_1}(a_0, a_1), \chi_{b_2, b_3, \dots, b_{n-1}}(a_2, a_3, \dots, a_{n-1}))$
3.  $\vdots$

4.  $\chi_{[b_0, b_1, \dots, b_{n-2}], b_{n-1}}(\chi_{b_0, b_1, \dots, b_{n-2}}(a_0, a_1, \dots, a_{n-2}), a_{n-1})$

### Declaration I:27

The notation  $\phi(n)$  will be used as a shorthand for the sublist of  $[0 : n]$  where each entry  $x$  is such that  $(x, n) = 1$ .

### Procedure I:72(1.47)

#### Objective

Choose an integer  $a$  and a positive integer  $b$  such that  $(a, b) = 1$ . The objective of the following instructions is to either show that  $0 < 0$  or to show that each element of  $a\phi(b) \bmod b$  is in  $\phi(b)$ .

#### Implementation

1. Show that  $(a, b) = 1$ .
2. For  $i$  in  $[0 : |\phi(b)|]$ , do the following:
  - (a) Show that  $(\phi(b)_i, b) = 1$  using **declaration I:27**.
  - (b) Use **procedure I:47** on  $\langle a, \phi(b)_i, b \rangle$  to show that  $(a\phi(b)_i, b) = 1$ .
  - (c) Use **procedure I:44** on  $\langle a\phi(b)_i \bmod b, a\phi(b)_i, b \rangle$  to show that  $(a\phi(b)_i \bmod b, b) = (a\phi(b)_i, b) = 1$ .
  - (d) Hence show that  $a\phi(b)_i \bmod b$  is contained in the list  $\phi(b)$  given that  $0 \leq a\phi(b)_i \bmod b < b$ .
3. **Hence show that each element of  $a\phi(b) \bmod b$  is in  $\phi(b)$ .**

### Procedure I:73(1.48)

#### Objective

Choose an integer  $a$  and a positive integer  $b$  such that  $(a, b) = 1$ . The objective of the following instructions is to either show that  $0 \neq 0$  or to show that each element of  $a\phi(b) \bmod b$  is distinct.

#### Implementation

1. Use **procedure I:36** on  $\langle a, b \rangle$  to construct  $\langle r, t, u, v, w \rangle$  and show that  $va + wb = r = (a, b) = 1$ .
2. Hence show that  $va \equiv 1 \pmod{b}$ .
3. Now for  $i$  in  $[0 : |\phi(b)|]$ , do the following:
  - (a) For  $j$  in  $[i + 1 : |\phi(b)|]$ , do the following:
    - i. If  $a\phi(b)_i \equiv a\phi(b)_j \pmod{b}$ , then do the following:
      - A. Show that  $\phi(b)_i \equiv va\phi(b)_i \equiv va\phi(b)_j \equiv \phi(b)_j \pmod{b}$ .
      - B. Hence show that  $\phi(b)_i = \phi(b)_j$ .
      - C. Show that  $\phi(b)_i \neq \phi(b)_j$  using **declaration I:27** given that  $i \neq j$ .
      - D. **Hence show that  $\phi(b)_i \neq \phi(b)_i$  given that  $\phi(b)_i = \phi(b)_j$  and  $\phi(b)_i \neq \phi(b)_j$ .**
      - E. **Abort procedure.**
    - ii. Otherwise, do the following:
      - A. Show that  $a\phi(b)_i \not\equiv a\phi(b)_j \pmod{b}$ .
4. **Therefore show that  $a\phi(b) \bmod b$  is composed of distinct elements.**

### Procedure I:74(1.49)

#### Objective

Choose an integer  $a$  and a positive integer  $b$  such that  $(a, b) = 1$ . The objective of the following instructions is to either show that  $0 < 0$  or to show that  $a\phi(b) \bmod b$  is a rearrangement of  $\phi(b)$ .

#### Implementation

1. Use **procedure I:72** on  $\langle a, b \rangle$  to show that each element of  $a\phi(b) \bmod b$  is in  $\phi(b)$ .
2. Show that  $|a\phi(b) \bmod b| = |\phi(b)|$ .
3. Use **procedure I:73** on  $\langle a, b \rangle$  to show that  $a\phi(b) \bmod b$  is composed of distinct elements.
4. **Hence show that  $a\phi(b) \bmod b$  is a rearrangement of  $\phi(b)$ .**

## Procedure I:75(1.50)

### Objective

Choose an integer  $a$  and a positive integer  $b$  such that  $(a, b) = 1$ . The objective of the following instructions is to show that either  $0 < 0$  or  $a^{|\phi(b)|} \equiv 1 \pmod{b}$ .

### Implementation

1. For  $i$  in  $[0 : |\phi(b)|]$ , do the following:
  - (a) Use **procedure I:36** on  $\langle \phi(b)_i, b \rangle$  to construct  $\langle r_i, t_i, u_i, v_i, w_i \rangle$  and show that  $v_i \phi(b)_i + w_i b = r_i = \langle \phi(b)_i, b \rangle$ .
  - (b) Show that  $v_i \phi(b)_i + w_i b = \langle \phi(b)_i, b \rangle = 1$  using **declaration I:27**.
  - (c) Hence show that  $v_i \phi(b)_i \equiv 1 \pmod{b}$ .
2. Hence using **procedure I:74**, show that  $\prod_i^{[0:|\phi(b)|]} \phi(b)_i$ 
  - (a)  $\equiv \prod_i^{[0:|\phi(b)|]} a \phi(b)_i$
  - (b)  $\equiv a^{|\phi(b)|} \prod_i^{[0:|\phi(b)|]} \phi(b)_i \pmod{b}$ .
3. **Hence show that 1**
  - (a)  $\equiv \prod_i^{[0:|\phi(b)|]} (v_i \phi(b)_i)$
  - (b)  $= \prod_i^{[0:|\phi(b)|]} v_i \prod_i^{[0:|\phi(b)|]} \phi(b)_i$
  - (c)  $\equiv a^{|\phi(b)|} \prod_i^{[0:|\phi(b)|]} \phi(b)_i \prod_i^{[0:|\phi(b)|]} v_i$
  - (d)  $\equiv a^{|\phi(b)|} \pmod{b}$ .

### Declaration I:28

The notation  $a \times b$  as a shorthand for the  $|a| \times |b|$  matrix such that for  $i$  in  $[0 : |a|]$ , for  $j$  in  $[0 : |b|]$ ,  $(a \times b)_{i,j} = \langle a_i, b_j \rangle$ .

## Procedure I:76(1.52)

### Objective

Choose two positive integers  $a, b$  such that  $(a, b) = 1$ . The objective of the following instructions is to show that each entry of  $\chi_{a,b}([0 : a] \times [0 : b])$  is in  $[0 : ab]$ .

### Implementation

1. Let  $h = \chi_{a,b}([0 : a] \times [0 : b])$ .
2. Show that  $0 \leq h_{i,j} < [a, b] = [a, b](a, b) = ab$  for  $i$  in  $[0 : a]$ , for  $j$  in  $[0 : b]$ .
3. **Hence show that each entry of  $h$  is in  $[0 : ab]$ .**

## Procedure I:77(1.53)

### Objective

Choose two positive integers  $a, b$  such that  $(a, b) = 1$ . The objective of the following instructions is to either show that  $0 < 0$  or to show that each entry of  $\chi_{a,b}([0 : a] \times [0 : b])$  is distinct.

### Implementation

1. Let  $h = \chi_{a,b}([0 : a] \times [0 : b])$ .
2. For each distinct unordered pair of index pairs  $\langle i, j \rangle$  and  $\langle k, l \rangle$  of  $h$ , do the following:
  - (a) If  $h_{i,j} = h_{k,l}$ , then do the following:
    - i. Show that  $\chi_{a,b}(i, j) = \chi_{a,b}([0 : a]_i, [0 : b]_j) = h_{i,j} = h_{k,l} = \chi_{a,b}([0 : a]_k, [0 : b]_l) = \chi_{a,b}(k, l)$ .
    - ii. Show that  $i \equiv \chi_{a,b}(i, j) = \chi_{a,b}(k, l) \equiv k \pmod{a}$  using **procedure I:68** given that  $\chi_{a,b}(i, j) = \chi_{a,b}(k, l)$ .
    - iii. Hence show that  $i = k$ .
    - iv. Show that  $j \equiv \chi_{a,b}(i, j) = \chi_{a,b}(k, l) \equiv l \pmod{b}$  using **procedure I:68** given that  $\chi_{a,b}(i, j) = \chi_{a,b}(k, l)$ .
    - v. Hence show that  $j = l$ .
    - vi. Hence show that  $\langle i, j \rangle = \langle k, l \rangle$ .
    - vii. **Hence show that  $\langle i, j \rangle \neq \langle i, j \rangle$  given that  $\langle i, j \rangle$  and  $\langle k, l \rangle$  are distinct.**
    - viii. **Abort procedure.**
  - (b) Otherwise do the following:
    - i. Show that  $h_{i,j} \neq h_{k,l}$ .
3. **Hence show that each entry of  $h$  is distinct.**

### Procedure I:78(1.54)

#### Objective

Choose two positive integers  $a, b$  such that  $(a, b) = 1$ . The objective of the following instructions is to show that either  $0 < 0$  or  $\chi_{a,b}([0 : a] \times [0 : b])$  is a rearrangement  $[0 : ab]$ .

#### Implementation

1. Let  $h = \chi_{a,b}([0 : a] \times [0 : b])$ .
2. Use [procedure I:76](#) on  $\langle a, b \rangle$  to show that each element of  $h$  is in  $[0 : ab]$ .
3. Also show that  $h$  has the same number of entries as  $[0 : ab]$ .
4. Use [procedure I:77](#) on  $\langle a, b \rangle$  to show that  $h$  is composed of distinct elements.
5. Hence show that  $h$  is a rearrangement of  $[0 : ab]$ .

### Procedure I:79(1.55)

#### Objective

Choose two positive integers  $a, b$  such that  $(a, b) = 1$ . The objective of the following instructions is to either show that  $0 < 0$  or to show that each entry of  $\chi_{a,b}(\phi(a) \times \phi(b))$  is in  $\phi(ab)$ .

#### Implementation

1. Let  $h = \chi_{a,b}(\phi(a) \times \phi(b))$ .
2. Now, for each index pair  $\langle i, j \rangle$  of  $h$ , do the following:
  - (a) Show that  $0 \leq h_{i,j} < [a, b] = [a, b](a, b) = ab$ .
  - (b) Show that  $h_{i,j} = \chi_{a,b}(\phi(a)_i, \phi(b)_j) \equiv \phi(a)_i \pmod{a}$ .
  - (c) Hence use [procedure I:44](#) on  $\langle h_{i,j}, \phi(a)_i, a \rangle$  to show that  $(a, h_{i,j}) = (h_{i,j}, a) = (\phi(a)_i, a) = 1$ .
  - (d) Also show that  $h_{i,j} = \chi_{a,b}(\phi(a)_i, \phi(b)_j) \equiv \phi(b)_j \pmod{b}$ .

(e) Hence use [procedure I:44](#) on  $\langle h_{i,j}, \phi(b)_j, b \rangle$  to show that  $(b, h_{i,j}) = (h_{i,j}, b) = (\phi(b)_j, b) = 1$ .

(f) Hence show that  $(h_{i,j}, ab) = (ab, h_{i,j}) = 1$ .

(g) Hence show that  $h_{i,j}$  is in  $\phi(ab)$ .

3. Hence show that each entry of  $\chi_{a,b}(\phi(a) \times \phi(b))$  is in  $\phi(ab)$ .

### Procedure I:80(1.56)

#### Objective

Choose two positive integers  $a, b$  such that  $(a, b) = 1$ . The objective of the following instructions is to either show that  $0 < 0$  or to show that each entry of  $\phi(ab)$  is in  $\chi_{a,b}(\phi(a) \times \phi(b))$ .

#### Implementation

1. For  $i$  in  $[0 : |\phi(ab)|]$ , do the following:
  - (a) Show that  $(\phi(ab)_i, ab) = 1$ .
  - (b) Show that  $\phi(ab)_i \equiv \phi(ab)_i \pmod{a}$ .
  - (c) Hence show that  $(\phi(ab)_i \pmod{a}, a) = (\phi(ab)_i, a) = 1$  using [procedure I:44](#).
  - (d) Hence show that  $\phi(ab)_i \pmod{a}$  is amongst  $\phi(a)$  given that  $0 \leq \phi(ab)_i \pmod{a} < a$ .
  - (e) Show that  $\phi(ab)_i \equiv \phi(ab)_i \pmod{b}$ .
  - (f) Hence show that  $(\phi(ab)_i \pmod{b}, b) = (\phi(ab)_i, b) = 1$  using [procedure I:44](#).
  - (g) Hence show that  $\phi(ab)_i \pmod{b}$  is amongst  $\phi(b)$  given that  $0 \leq \phi(ab)_i \pmod{b} < b$ .
  - (h) Hence show that  $\langle \phi(ab)_i \pmod{a}, \phi(ab)_i \pmod{b} \rangle$  is amongst  $\phi(a) \times \phi(b)$ .
  - (i) Show that  $\phi(ab)_i \equiv \chi_{a,b}(\phi(ab)_i \pmod{a}, \phi(ab)_i \pmod{b}) \pmod{[a, b]} = [a, b](a, b) = ab$  using [procedure I:65](#) given that  $\phi(ab)_i \equiv \phi(ab)_i \pmod{a} \pmod{a}$  and  $\phi(ab)_i \equiv \phi(ab)_i \pmod{b} \pmod{b}$ .
  - (j) Hence show that  $\phi(ab)_i = \chi_{a,b}(\phi(ab)_i \pmod{a}, \phi(ab)_i \pmod{b})$ .

- (k) Hence show that  $\phi(ab)_i$  is amongst  $\chi_{a,b}(\phi(a) \times \phi(b))$  given that  $\langle \phi(ab)_i \bmod a, \phi(ab)_i \bmod b \rangle$  is amongst  $\phi(a) \times \phi(b)$  and  $\phi(ab)_i = \chi_{a,b}(\phi(ab)_i \bmod a, \phi(ab)_i \bmod b)$ .

2. Hence show that each entry of  $\phi(ab)$  is in  $\chi_{a,b}(\phi(a) \times \phi(b))$ .

## Procedure I:81(1.57)

### Objective

Choose two positive integers  $a, b$  such that  $(a, b) = 1$ . The objective of the following instructions is to either show that  $0 < 0$  or to show that  $\phi(ab)$  is a rearrangement of  $\chi_{a,b}(\phi(a) \times \phi(b))$  and that  $|\phi(ab)| = |\phi(a)||\phi(b)|$ .

### Implementation

1. Use **procedure I:78** on  $\langle a, b \rangle$  to show that  $\chi_{a,b}([0 : a] \times [0 : b])$  is a rearrangement of  $[0 : ab]$ .
2. Show that  $\chi_{a,b}(\phi(a) \times \phi(b))$  is a submatrix of  $\chi_{a,b}([0 : a] \times [0 : b])$ .
3. Hence show that the entries of  $\chi_{a,b}(\phi(a) \times \phi(b))$  are distinct.
4. Use **procedure I:79** on  $\langle a, b \rangle$  to show that the entries of  $\chi_{a,b}(\phi(a) \times \phi(b))$  are in  $\phi(ab)$ .
5. Show that the entries of  $\phi(ab)$  are distinct.
6. Use **procedure I:80** on  $\langle a, b \rangle$  to show that the entries of  $\phi(ab)$  are in  $\chi_{a,b}(\phi(a) \times \phi(b))$ .
7. Hence show that  $\phi(ab)$  is a rearrangement of  $\chi_{a,b}(\phi(a) \times \phi(b))$ .
8. Hence show that  $|\phi(ab)| = |\chi_{a,b}(\phi(a) \times \phi(b))| = |\phi(a) \times \phi(b)| = |\phi(a)||\phi(b)|$ .

## Declaration I:29

The notation  $[P]$ , where  $P$  is a condition, will be used as a shorthand for 1 if  $P$ , otherwise it will stand for 0.

## Declaration I:30

The notation  $a_+$ , where  $a$  is a list, will be used as a shorthand for 0 if  $a$  is empty, otherwise it will be a shorthand for the sum of the entries of  $a$ .

## Declaration I:31

The notation  $\sum_r^R f(r)$ , where  $R$  is a list and  $f[r]$  is a function of  $r$ , will be used as a shorthand for  $f(R)_+$ .

## Procedure I:82(1.58)

### Objective

Choose a positive integer  $a$  and a prime  $b$ . The objective of the following instructions is to show that either  $0 < 0$  or  $|\phi(b^a)| = b^a - b^{a-1}$ .

### Implementation

1. Show that  $\sum_r^{[0:b^a]} [(r, b^a) = 1] \leq \sum_r^{[0:b^a]} [(r, b) = 1]$  using **procedure I:48**.
2. Show that  $\sum_r^{[0:b^a]} [(r, b) = 1] \leq \sum_r^{[0:b^a]} [(r, b^a) = 1]$  using **procedure I:47**.
3. Hence show that  $\sum_r^{[0:b^a]} [(r, b^a) = 1] = \sum_r^{[0:b^a]} [(r, b) = 1]$ .
4. Show that  $\sum_r^{[0:b^a]} [(r, b) = 1] \leq \sum_r^{[0:b^a]} [r \bmod b \neq 0]$  using **procedure I:40**.
5. Show that  $\sum_r^{[0:b^a]} [r \bmod b \neq 0] \leq \sum_r^{[0:b^a]} [(r, b) = 1]$  using **procedure I:49**.
6. Hence show that  $\sum_r^{[0:b^a]} [(r, b) = 1] = \sum_r^{[0:b^a]} [r \bmod b \neq 0]$ .
7. Hence show that  $|\phi(b^a)| = \sum_r^{[0:b^a]} [(r, b^a) = 1] = \sum_r^{[0:b^a]} [(r, b) = 1] = \sum_r^{[0:b^a]} [r \bmod b \neq 0] = \sum_r^{[0:b^a]} (1 - [r \bmod b = 0]) = b^a - b^{a-1}$ .

## Procedure I:83(1.59)

### Objective

Choose a list of primes  $a$ . Let  $b$  be the list of distinct primes in  $a$ . Let  $c$  be a list such that  $c_i$  is the mul-

tiplicity of  $b_i$  in  $a$  for  $i = 1$  to  $i = |b|$ . The objective of the following instructions is to show that either  $0 < 0$  or  $|\phi(a_*)| = \prod_i^{[0:|b|]} (b_i^{c_i} - b_i^{c_i-1})$ .

### Implementation

1. If  $a = \langle \rangle$ , then do the following:
  - (a) Show that  $|b| = |a| = 0$ .
  - (b) **Hence show that**  $\phi(a_*) = \phi(1) = 1 = \prod_i^{[0:|b|]} (b_i^{c_i} - b_i^{c_i-1})$ .
2. Otherwise, do the following:
  - (a) Show that  $a_* = \prod_i^{[0:|b|]} b_i^{c_i}$ .
  - (b) Show that  $|a| > 0$ .
  - (c) Hence show that  $|c| = |b| > 0$ .
  - (d) Hence show that  $(b_0^{c_0}, \prod_i^{[1:|b|]} b_i^{c_i}) = 1$  using **procedure I:57**.
  - (e) Let  $d$  be the list  $a$  with all instances of  $a_0$  removed.
  - (f) Verify that  $|d| < |a|$ .
  - (g) Now use **procedure I:83** on  $\langle d \rangle$  to show that  $\phi(d_*) = \phi(\prod_i^{[1:|b|]} b_i^{c_i}) = \prod_i^{[1:|b|]} (b_i^{c_i} - b_i^{c_i-1})$ .
  - (h) **Hence show that**

$$\begin{aligned}
|\phi(a_*)| &= |\phi(\prod_i^{[0:|b|]} b_i^{c_i})| = |\phi(b_0^{c_0} \prod_i^{[1:|b|]} b_i^{c_i})| = \\
&= |\phi(b_0^{c_0})| |\phi(\prod_i^{[1:|b|]} b_i^{c_i})| = (b_0^{c_0} - b_0^{c_0-1}) |\phi(\prod_i^{[1:|b|]} b_i^{c_i})| = \\
&= (b_0^{c_0} - b_0^{c_0-1}) \prod_i^{[1:|b|]} (b_i^{c_i} - b_i^{c_i-1}) = \prod_i^{[0:|b|]} (b_i^{c_i} - b_i^{c_i-1})
\end{aligned}$$
**using procedure I:81 and procedure I:82 given that**  $(b_0^{c_0}, \prod_i^{[1:|b|]} b_i^{c_i}) = 1$  **and**  $\phi(\prod_i^{[1:|b|]} b_i^{c_i}) = \prod_i^{[1:|b|]} (b_i^{c_i} - b_i^{c_i-1})$ .

## Chapter 4

# Permutations and Combinations

### Declaration I:32

The notation  $a^{\underline{b}}$  will be used as a shorthand for  $\prod_i^{[0:b]}(a-i)$ .

### Declaration I:33

The notation  $a^{\overline{b}}$  will be used as a shorthand for  $\prod_i^{[0:b]}(a+i)$ .

### Procedure I:84(1.60)

#### Objective

Choose a list of distinct elements  $a$  and a non-negative integer  $b$  such that  $b \leq |a|$ . Let  $c$  be a list of length- $b$  permutations of  $a$ . The objective of the following instructions is to show that  $|c| = |a|^{\underline{b}}$ .

#### Implementation

1. If  $|b| > 0$ , then do the following:
  - (a) For each entry  $d$  in  $a$ , do the following:
    - i. Let  $e$  be the list formed by removing  $d$  from  $a$ .
    - ii. Show that the entries of  $e$  are distinct given that the entries of  $a$  are distinct.
    - iii. Show that  $|e| = |a| - 1$ .
    - iv. Now use **procedure I:84** on  $\langle e, b-1 \rangle$  to show that the number of length- $b-1$  permutations of  $e$  is  $|e|^{\underline{b-1}}$ .

v. Hence show that the number of length- $b$  permutations of  $a$  beginning with  $d$  is  $|e|^{\underline{b-1}} = (|a|-1)^{\underline{b-1}}$ .

(b) Hence show that the number of length- $b$  permutations of  $a$  beginning with any entry of  $a$  is  $|a|(|a|-1)^{\underline{b-1}} = |a|^{\underline{b}}$ .

(c) Hence show that the number of length- $b$  permutations of  $a$  are  $|a|^{\underline{b}}$ .

(d) **Hence show that**  $|c| = |a|^{\underline{b}}$ .

2. Otherwise do the following:

(a) Show that  $b = 0$ .

(b) Show that the number of length-0 permutations of  $a$  is 1.

(c) **Therefore show that**  $|c| = 1 = |a|^{\underline{0}} = |a|^{\underline{b}}$ .

### Declaration I:34

The notation  $\binom{n}{r}$  will be used as a shorthand for  $n^{\underline{r}} \text{div}(r!)$ .

### Procedure I:85(1.61)

#### Objective

Choose a list of distinct elements  $n$  and a non-negative integer  $r$  such that  $r \leq |n|$ . Let  $b$  be the largest list of length- $r$  sublists of  $n$  such that no two of them are permutations of each other. The objective of the following instructions is to either show that  $b$  contains at least two permutations of the same list, construct a list larger than  $b$  that is

also a list of length- $r$  sublists of  $n$  such that no two of them are permutations of each other, or to show that  $|b| = \binom{|n|}{r}$  and that  $|n|^x \bmod r! = 0$ .

### Implementation

1. Let  $a$  and  $f$  be a list of all the permutations of  $n$ .
2. Show that  $|a| = |n|^{\underline{|n|}}$  using **procedure I:84**.
3. For each list  $c$  in  $b$ , do the following:
  - (a) Show that the number of permutations of  $c$  is  $r!$  using **procedure I:84**.
  - (b) Let  $d$  be the list obtained by removing the elements of  $c$  from  $n$ .
  - (c) Show that the number of permutations of  $d$  is  $(n - r)!$  using **procedure I:84**.
  - (d) Let  $e$  be the list of permutations of  $n$  beginning with a permutation of  $c$ .
  - (e) Show that  $|e| = r!(|n| - r)!$  given that there are  $r!$  possible choices for the first part of  $e$  and  $(|n| - r)!$  possible choices for the second part of  $e$ .
  - (f) If  $e$  is not a sublist of  $a$ , then do the following:
    - i. Let  $g$  be a list in  $e$  that is not in  $a$ .
    - ii. Show that  $e$  is a sublist of  $f$ .
    - iii. Therefore show that  $g$  was in  $a$  but then was removed.
    - iv. Therefore show that the variable  $c$  was formerly equal to a permutation of the current  $c$ .
    - v. **Therefore show that  $b$  contains at least two permutations of  $c$ .**
    - vi. **Abort procedure.**
  - (g) Otherwise, do the following:
    - i. Show that  $e$  is a sublist of  $a$ .
    - ii. Remove the lists in  $e$  from  $a$ .
4. If  $a \neq \langle \rangle$ , then do the following:
  - (a) Let  $g$  be a list in  $a$ .
  - (b) Let  $h$  be the sublist of  $g$  corresponding to its first  $r$  elements.

- (c) Therefore show that the permutations of  $n$  beginning with a permutation of  $h$  were never removed from  $a$ .
  - (d) Therefore show that the variable  $c$  was never equal to a permutation of  $h$ .
  - (e) Therefore show that no permutation of  $h$  is in  $b$ .
  - (f) **Therefore show that  $b \setminus \langle h \rangle$  is larger than  $b$  and is also a list of length- $r$  sublists of  $n$  such that no two of them are permutations of each other.**
  - (g) **Abort procedure.**
5. Otherwise do the following:
    - (a) Show that  $|n|! \bmod (r!(|n| - r)!) = 0$ .
    - (b) **Therefore show that  $n^x \bmod r!$** 
      - i.  $= (|n|! \operatorname{div}(|n| - r)!) \bmod r!$
      - ii.  $= ((|n|! \bmod (r!(|n| - r)!)r!(|n| - r)!) \operatorname{div}(|n| - r)!) \bmod r!$
      - iii.  $= ((|n|! \operatorname{div}(r!(|n| - r)!))r!) \bmod r!$
      - iv.  $= 0$ .
    - (c) Also show that (3) iterated  $|n|! \operatorname{div}(r!(|n| - r)!) \operatorname{div}(r!)$  times.
    - (d) **Therefore using procedure I:35, show that  $|b|$** 
      - i.  $= |n|! \operatorname{div}(r!(|n| - r)!)$
      - ii.  $= (|n|! \operatorname{div}(|n| - r)!) \operatorname{div}(r!)$
      - iii.  $= n^x \operatorname{div}(r!)$
      - iv.  $= \binom{n}{r}$ .

### Procedure I:86(1.62)

#### Objective

Choose two positive integers  $a, b$ . The objective of the following instructions is to show that  $\binom{a}{b} = \binom{a-1}{b-1} + \binom{a-1}{b}$ .



## Implementation

$$\text{vii.} = \sum_r^{[0:a+1]} \binom{a}{r} x^r.$$

1. Using **procedure I:32** and **procedure I:33**, show that  $\binom{a-1}{b-1} + \binom{a-1}{b}$

$$(a) = (a-1)^{b-1} \text{div}(b-1)! + (a-1)^b \text{div } b!$$

$$(b) = ((a-1)^{b-1}b) \text{div } b! + (a-1)^b \text{div } b!$$

$$(c) = ((a-1)^{b-1}b + (a-1)^b) \text{div } b!$$

$$(d) = ((a-1)^{b-1}b + (a-1)^{b-1}(a-b)) \text{div } b!$$

$$(e) = ((a-1)^{b-1}a) \text{div } b!$$

$$(f) = a^b \text{div } b!$$

$$(g) = \binom{a}{b}.$$

## Procedure I:87(1.63)

### Objective

Choose an integer  $x$  and a non-negative integer  $a$ . The objective of the following instructions is to show that the  $(1+x)^a = \sum_r^{[0:a+1]} \binom{a}{r} x^r$ .

## Implementation

1. If  $a = 0$ , then do the following:

$$(a) \text{ **Show that** } (1+x)^a = (1+x)^0 = 1 = \sum_r^{[0:1]} \binom{0}{r} x^r = \sum_r^{[0:a+1]} \binom{a}{r} x^r.$$

2. Otherwise, do the following:

$$(a) \text{ Show that } a > 0.$$

$$(b) \text{ Therefore show that } a-1 \geq 0.$$

$$(c) \text{ Use **procedure I:87** on } \langle x, a-1 \rangle \text{ to show that } (1+x)^{a-1} = \sum_r^{[0:a]} \binom{a-1}{r} x^r.$$

$$(d) \text{ Therefore using **procedure I:86**, show that } (1+x)^a$$

$$\text{i.} = (1+x)(1+x)^{a-1}$$

$$\text{ii.} = (1+x) \sum_r^{[0:a]} \binom{a-1}{r} x^r$$

$$\text{iii.} = \sum_r^{[0:a]} \binom{a-1}{r} x^r + \sum_r^{[0:a]} \binom{a-1}{r} x^{r+1}$$

$$\text{iv.} = \sum_r^{[0:a+1]} \binom{a-1}{r} x^r + \sum_r^{[1:a+1]} \binom{a-1}{r-1} x^r$$

$$\text{v.} = 1 + \sum_r^{[1:a+1]} \left( \binom{a-1}{r} + \binom{a-1}{r-1} \right) x^r$$

$$\text{vi.} = 1 + \sum_r^{[1:a+1]} \binom{a}{r} x^r$$

## Part II

# Rational Arithmetic

## Chapter 5

# Rational Arithmetic

### Declaration II:0

The phrase "rational number" will be used as a shorthand for an ordered pair comprising an integer followed by a non-zero natural number.

### Declaration II:1

The phrase "the numerator of  $a$ " and the notation  $\text{nu}(a)$ , where  $a$  is a rational number, will be used as a shorthand for the first entry of  $a$ .

### Declaration II:2

The phrase "the denominator of  $a$ " and the notation  $\text{de}(a)$ , where  $a$  is a rational number, will be used as a shorthand for the second entry of  $a$ .

### Declaration II:3

The phrase " $a = b$ ", where  $a, b$  are rational numbers, will be used as a shorthand for " $\text{nu}(a) \text{de}(b) = \text{de}(a) \text{nu}(b)$ ".

### Procedure II:0(2.27)

#### Objective

Choose a rational number  $a$ . The objective of the following instructions is to show that  $a = a$ .

### Implementation

1. Verify that  $\text{nu}(a) \text{de}(a) = \text{de}(a) \text{nu}(a)$ .
2. **Hence verify that  $a = a$ .**

### Procedure II:1(2.28)

#### Objective

Choose two rational numbers  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $b = a$ .

### Implementation

1. Verify that  $\text{nu}(a) \text{de}(b) = \text{de}(a) \text{nu}(b)$ .
2. Hence verify that  $\text{nu}(b) \text{de}(a) = \text{de}(b) \text{nu}(a)$ .
3. **Hence verify that  $b = a$ .**

### Procedure II:2(2.29)

#### Objective

Choose three rational numbers  $a, b, c$  such that  $a = b$  and  $b = c$ . The objective of the following instructions is to show that  $a = c$ .

## Implementation

1. Using **declaration II:3**, verify that  $\text{nu}(a) \text{de}(b) = \text{de}(a) \text{nu}(b)$ .
2. Using **declaration II:3**, verify that  $\text{nu}(b) \text{de}(c) = \text{de}(b) \text{nu}(c)$ .
3. If  $\text{nu}(b) \neq 0$ , then do the following:
  - (a) Hence verify that  $\text{nu}(a) \text{de}(b) \text{nu}(b) \text{de}(c) = \text{de}(a) \text{nu}(b) \text{de}(b) \text{nu}(c)$ .
  - (b) Hence verify that  $\text{nu}(a) \text{de}(c) = \text{de}(a) \text{nu}(c)$ .
4. Otherwise do the following:
  - (a) Using **declaration II:0**, verify that  $\text{de}(b) \neq 0$ .
  - (b) Verify that  $\text{nu}(a) \text{de}(b) = \text{de}(a) \text{nu}(b) = 0$ .
  - (c) Hence verify that  $\text{nu}(a) = 0$ .
  - (d) Verify that  $0 = \text{nu}(b) \text{de}(c) = \text{de}(b) \text{nu}(c)$ .
  - (e) Hence verify that  $\text{nu}(c) = 0$ .
  - (f) Hence verify that  $\text{nu}(a) \text{de}(c) = 0 = \text{de}(a) \text{nu}(c)$ .
5. **Hence verify that  $a = c$ .**

## Declaration II:4

The notation  $a + b$ , where  $a, b$  are rational numbers, will be used as a shorthand for the pair  $\langle \text{nu}(a) \text{de}(b) + \text{de}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle$ .

## Procedure II:3(2.30)

### Objective

Choose two rational numbers  $a, b, c, d$  such that  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $a + b = c + d$ .

## Implementation

1. Using **declaration II:3**, verify that  $\text{nu}(a) \text{de}(c) = \text{de}(a) \text{nu}(c)$ .
2. Using **declaration II:3**, verify that  $\text{nu}(b) \text{de}(d) = \text{de}(b) \text{nu}(d)$ .
3. Hence verify that  $a + b$ 
  - (a)  $= \langle \text{nu}(a), \text{de}(a) \rangle + \langle \text{nu}(b), \text{de}(b) \rangle$

- (b)  $= \langle \text{nu}(a) \text{de}(b) + \text{de}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle$
- (c)  $= \langle \text{de}(c) \text{de}(d) (\text{nu}(a) \text{de}(b) + \text{de}(a) \text{nu}(b)), \text{de}(c) \text{de}(d) (\text{de}(a) \text{de}(b)) \rangle$
- (d)  $= \langle \text{nu}(a) \text{de}(c) \text{de}(b) \text{de}(d) + \text{de}(a) \text{de}(c) \text{nu}(b) \text{de}(d), \text{de}(c) \text{de}(d) \text{de}(a) \text{de}(b) \rangle$
- (e)  $= \langle \text{de}(a) \text{nu}(c) \text{de}(b) \text{de}(d) + \text{de}(a) \text{de}(c) \text{de}(b) \text{nu}(d), \text{de}(c) \text{de}(d) \text{de}(a) \text{de}(b) \rangle$
- (f)  $= \langle \text{de}(a) \text{de}(b) (\text{nu}(c) \text{de}(d) + \text{de}(c) \text{nu}(d)), \text{de}(a) \text{de}(b) (\text{de}(c) \text{de}(d)) \rangle$
- (g)  $= \langle \text{nu}(c) \text{de}(d) + \text{de}(c) \text{nu}(d), \text{de}(c) \text{de}(d) \rangle$
- (h)  $= \langle \text{nu}(c), \text{de}(c) \rangle + \langle \text{nu}(d), \text{de}(d) \rangle$
- (i)  $= c + d$ .

## Procedure II:4(2.31)

### Objective

Choose three rational numbers  $a, b, c$ . The objective of the following instructions is to show that  $(a + b) + c = a + (b + c)$ .

## Implementation

1. Verify that  $(a + b) + c$ 
  - (a)  $= \langle \text{nu}(a) \text{de}(b) + \text{de}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle + \langle \text{nu}(c), \text{de}(c) \rangle$
  - (b)  $= \langle (\text{nu}(a) \text{de}(b) + \text{de}(a) \text{nu}(b)) \text{de}(c) + (\text{de}(a) \text{de}(b)) \text{nu}(c), (\text{de}(a) \text{de}(b)) \text{de}(c) \rangle$
  - (c)  $= \langle \text{nu}(a) (\text{de}(b) \text{de}(c)) + \text{de}(a) (\text{nu}(b) \text{de}(c) + \text{de}(b) \text{nu}(c)), \text{de}(a) (\text{de}(b) \text{de}(c)) \rangle$
  - (d)  $= \langle \text{nu}(a), \text{de}(a) \rangle + \langle \text{nu}(b) \text{de}(c) + \text{de}(b) \text{nu}(c), \text{de}(b) \text{de}(c) \rangle$
  - (e)  $= a + (b + c)$ .

## Procedure II:5(2.32)

### Objective

Choose two rational numbers  $a, b$ . The objective of the following instructions is to show that  $a + b = b + a$ .

### Implementation

1.  $a + b$ 

$$(a) = \langle \text{nu}(a) \text{de}(b) + \text{de}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle$$

$$(b) = \langle \text{nu}(b) \text{de}(a) + \text{de}(b) \text{nu}(a), \text{de}(b) \text{nu}(a) \rangle$$

$$(c) = b + a.$$

### Declaration II:5

The notation  $\textcolor{red}{a}$ , where  $a$  is an integer, will contextually be used as a shorthand for the pair  $\langle a, 1 \rangle$ .

### Procedure II:6(2.33)

#### Objective

Choose a rational number  $a$ . The objective of the following instructions is to show that  $0 + a = a$ .

### Implementation

1. Verify that  $0 + a$ 

$$(a) = \langle 0, 1 \rangle + \langle \text{nu}(a), \text{de}(a) \rangle$$

$$(b) = \langle 0 \text{de}(a) + 1 \text{nu}(a), 1 \text{de}(a) \rangle$$

$$(c) = \langle \text{nu}(a), \text{de}(a) \rangle$$

$$(d) = a.$$

### Declaration II:6

The notation  $\textcolor{red}{-a}$ , where  $a$  is a rational number, will be used as a shorthand for the pair  $\langle -\text{nu}(a), \text{de}(a) \rangle$ .

### Procedure II:7(2.34)

#### Objective

Choose two rational numbers  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $-a = -b$ .

### Implementation

1. Using **declaration II:3**, verify that  $\text{nu}(a) \text{de}(b) = \text{de}(a) \text{nu}(b)$ .
2. Hence verify that  $-a$ 

$$(a) = \langle -\text{nu}(a), \text{de}(a) \rangle$$

$$(b) = \langle -\text{nu}(a) \text{de}(b), \text{de}(a) \text{de}(b) \rangle$$

$$(c) = \langle -\text{de}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle$$

$$(d) = \langle -\text{nu}(b), \text{de}(b) \rangle$$

$$(e) = -b.$$

### Procedure II:8(2.35)

#### Objective

Choose a rational number  $a$ . The objective of the following instructions is to show that  $-a + a = 0$ .

### Implementation

1. Verify that  $-a + a$ 

$$(a) = (-a) + a$$

$$(b) = \langle -\text{nu}(a), \text{de}(a) \rangle + \langle \text{nu}(a), \text{de}(a) \rangle$$

$$(c) = \langle -\text{nu}(a) \text{de}(a) + \text{de}(a) \text{nu}(a), \text{de}(a)^2 \rangle$$

$$(d) = \langle 0, \text{de}(a)^2 \rangle$$

$$(e) = \langle 0, 1 \rangle$$

$$(f) = 0.$$

### Declaration II:7

The notation  $\textcolor{red}{ab}$ , where  $a, b$  are rational numbers, will be used as a shorthand for the pair  $\langle \text{nu}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle$ .

### Procedure II:9(2.36)

#### Objective

Choose two rational numbers  $a, b, c, d$  such that  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $ab = cd$ .

### Implementation

1. Using **declaration II:3**, verify that  $\text{nu}(a) \text{de}(c) = \text{de}(a) \text{nu}(c)$ .
2. Using **declaration II:3**, verify that  $\text{nu}(b) \text{de}(d) = \text{de}(b) \text{nu}(d)$ .
3. Hence verify that  $ab$

$$(a) = \langle \text{nu}(a), \text{de}(a) \rangle \langle \text{nu}(b), \text{de}(b) \rangle$$

$$(b) = \langle \text{nu}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle$$

$$(c) = \langle (\text{de}(c) \text{de}(d)) \text{nu}(a) \text{nu}(b), (\text{de}(c) \text{de}(d)) \text{de}(a) \text{de}(b) \rangle$$

$$(d) = \langle (\text{nu}(a) \text{de}(c)) (\text{nu}(b) \text{de}(d)), \text{de}(c) \text{de}(d) \text{de}(a) \text{de}(b) \rangle$$

$$(e) = \langle (\text{de}(a) \text{nu}(c)) (\text{de}(b) \text{nu}(d)), \text{de}(c) \text{de}(d) \text{de}(a) \text{de}(b) \rangle$$

$$(f) = \langle (\text{de}(a) \text{de}(b)) \text{nu}(c) \text{nu}(d), (\text{de}(a) \text{de}(b)) \text{de}(c) \text{de}(d) \rangle$$

$$(g) = \langle \text{nu}(c) \text{nu}(d), \text{de}(c) \text{de}(d) \rangle$$

$$(h) = \langle \text{nu}(c), \text{de}(c) \rangle \langle \text{nu}(d), \text{de}(d) \rangle$$

$$(i) = cd.$$

### Procedure II:10(2.37)

#### Objective

Choose three rational numbers  $a, b, c$ . The objective of the following instructions is to show that  $(ab)c = a(bc)$ .

### Implementation

1. Verify that  $(ab)c$ 

$$(a) = \langle \text{nu}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle \langle \text{nu}(c), \text{de}(c) \rangle$$

$$(b) = \langle \text{nu}(a) \text{nu}(b) \text{nu}(c), \text{de}(a) \text{de}(b) \text{de}(c) \rangle$$

$$(c) = \langle \text{nu}(a), \text{de}(a) \rangle \langle \text{nu}(b) \text{nu}(c), \text{de}(b) \text{de}(c) \rangle$$

$$(d) = a(bc).$$

### Procedure II:11(2.38)

#### Objective

Choose two rational numbers  $a, b$ . The objective of the following instructions is to show that  $ab = ba$ .

### Implementation

1.  $ab$ 

$$(a) = \langle \text{nu}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle$$

$$(b) = \langle \text{nu}(b) \text{nu}(a), \text{de}(b) \text{de}(a) \rangle$$

$$(c) = ba.$$

### Procedure II:12(2.39)

#### Objective

Choose a rational number  $a$ . The objective of the following instructions is to show that  $1a = a$ .

### Implementation

1. Verify that  $1a$ 

$$(a) = \langle 1, 1 \rangle \langle \text{nu}(a), \text{de}(a) \rangle$$

$$(b) = \langle 1 \text{nu}(a), 1 \text{de}(a) \rangle$$

$$(c) = \langle \text{nu}(a), \text{de}(a) \rangle$$

$$(d) = a.$$

### Declaration II:8

The notation  $\frac{1}{a}$ , where  $a$  is a rational number such that  $\text{nu}(a) > 0$ , will be used as a shorthand for the pair  $\langle \text{de}(a), \text{nu}(a) \rangle$ .

### Declaration II:9

The notation  $\frac{1}{a}$ , where  $a$  is a rational number such that  $\text{nu}(a) < 0$ , will be used as a shorthand for the pair  $\langle -\text{de}(a), -\text{nu}(a) \rangle$ .

### Procedure II:13(2.40)

#### Objective

Choose two rational numbers  $a, b$  such that  $a = b$  and  $a \neq 0$ . The objective of the following instructions is to show that  $\frac{1}{a} = \frac{1}{b}$ .

### Implementation

1. Using **declaration II:3**, verify that  $\text{nu}(a) \text{de}(b) = \text{de}(a) \text{nu}(b)$ .
2. Using **declaration II:3** and **declaration II:5**, verify that  $\text{nu}(a) = \text{nu}(a) \text{de}(0) \neq \text{de}(a) \text{nu}(0) = 0$ .
3. Hence using **declaration II:0**, verify that  $\text{de}(a) \text{nu}(b) = \text{nu}(a) \text{de}(b) \neq 0$ .
4. Hence verify that  $\text{nu}(b) \neq 0$ .
5. If  $\text{nu}(a) \text{nu}(b) > 0$ , then do the following:
  - (a) Verify that  $\frac{1}{a}$ 
    - i.  $= \langle \text{de}(a) \text{nu}(b), \text{nu}(a) \text{nu}(b) \rangle$
    - ii.  $= \langle \text{nu}(a) \text{de}(b), \text{nu}(a) \text{nu}(b) \rangle$
    - iii.  $= \frac{1}{b}$ .
6. Otherwise do the following:
  - (a) Verify that  $\text{nu}(a) \text{nu}(b) < 0$ .
  - (b) Hence verify that  $\frac{1}{a}$ 
    - i.  $= \langle -\text{de}(a) \text{nu}(b), -\text{nu}(a) \text{nu}(b) \rangle$
    - ii.  $= \langle -\text{nu}(a) \text{de}(b), -\text{nu}(a) \text{nu}(b) \rangle$
    - iii.  $= \frac{1}{b}$ .

### Procedure II:14(2.41)

#### Objective

Choose a rational number  $a$  such that  $a \neq 0$ . The objective of the following instructions is to show that  $\frac{1}{a}a = 1$ .

#### Implementation

1. Using **declaration II:3** and **declaration II:5**, verify that  $\text{nu}(a) = \text{nu}(a) \text{de}(0) \neq \text{de}(a) \text{nu}(0) = 0$ .
2. If  $\text{nu}(a) > 0$ , then do the following:
  - (a) Verify that  $\frac{1}{a}a$ 
    - i.  $= \langle \text{de}(a), \text{nu}(a) \rangle \langle \text{nu}(a), \text{de}(a) \rangle$
    - ii.  $= \langle \text{de}(a) \text{nu}(a), \text{nu}(a) \text{de}(a) \rangle$
    - iii.  $= \langle 1, 1 \rangle$

iv.  $= 1$ .

3. Otherwise do the following:

- (a) Verify that  $\text{nu}(a) < 0$ .
- (b) Hence verify that  $\frac{1}{a}a$ 
  - i.  $= \langle -\text{de}(a), -\text{nu}(a) \rangle \langle \text{nu}(a), \text{de}(a) \rangle$
  - ii.  $= \langle -\text{de}(a) \text{nu}(a), -\text{nu}(a) \text{de}(a) \rangle$
  - iii.  $= \langle 1, 1 \rangle$
  - iv.  $= 1$ .

### Procedure II:15(2.42)

#### Objective

Choose three rational numbers  $a, b, c$ . The objective of the following instructions is to show that  $a(b + c) = ab + ac$ .

#### Implementation

1.  $a(b + c)$ 
  - (a)  $= \langle \text{nu}(a), \text{de}(a) \rangle \langle \text{nu}(b) \text{de}(c) + \text{de}(b) \text{nu}(c), \text{de}(b) \text{de}(c) \rangle$
  - (b)  $= \langle \text{nu}(a)(\text{nu}(b) \text{de}(c) + \text{de}(b) \text{nu}(c)), \text{de}(a)(\text{de}(b) \text{de}(c)) \rangle$
  - (c)  $= \langle \text{nu}(a) \text{nu}(b) \text{de}(c) + \text{nu}(a) \text{de}(b) \text{nu}(c), \text{de}(a) \text{de}(b) \text{de}(c) \rangle$
  - (d)  $= \langle \text{de}(a)(\text{nu}(a) \text{nu}(b) \text{de}(c) + \text{nu}(a) \text{de}(b) \text{nu}(c)), \text{de}(a)(\text{de}(a) \text{de}(b) \text{de}(c)) \rangle$
  - (e)  $= \langle (\text{nu}(a) \text{nu}(b))(\text{de}(a) \text{de}(c)) + (\text{de}(a) \text{de}(b))(\text{nu}(a) \text{nu}(c)), (\text{de}(a) \text{de}(b))(\text{de}(a) \text{de}(c)) \rangle$
  - (f)  $= \langle \text{nu}(a) \text{nu}(b), \text{de}(a) \text{de}(b) \rangle + \langle \text{nu}(a) \text{nu}(c), \text{de}(a) \text{de}(c) \rangle$
  - (g)  $= ab + ac$ .

### Procedure II:16(2.09)

#### Objective

Choose an integer  $a$ . The objective of the following instructions is to show that  $(-1)^{2a} = 1$  and  $(-1)^{2a+1} = -1$ .

### Implementation

Implementation is analogous to that of **procedure I:14**.

### Declaration II:10

The phrase " $a < b$ ", where  $a, b$  are rational numbers, will be used as a shorthand for " $\text{nu}(a) \text{de}(b) < \text{de}(a) \text{nu}(b)$ ".

### Procedure II:17(2.43)

#### Objective

Choose four rational numbers  $a, b, c, d$  such that  $a < b$ ,  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $c < d$ .

### Implementation

1. Using **declaration II:3**, verify that  $\text{nu}(a) \text{de}(c) = \text{de}(a) \text{nu}(c)$ .
2. Using **declaration II:3**, verify that  $\text{nu}(b) \text{de}(d) = \text{de}(b) \text{nu}(d)$ .
3. Using **declaration II:10**, verify that  $\text{nu}(a) \text{de}(b) < \text{de}(a) \text{nu}(b)$ .
4. Hence verify that  $\text{nu}(c) \text{de}(d) \text{de}(a) \text{de}(b)$ 
  - (a)  $= \text{nu}(a) \text{de}(c) \text{de}(d) \text{de}(b)$
  - (b)  $< \text{de}(a) \text{nu}(b) \text{de}(c) \text{de}(d)$
  - (c)  $= \text{de}(b) \text{nu}(d) \text{de}(a) \text{de}(c)$ .
5. Hence verify that  $\text{nu}(c) \text{de}(d) < \text{de}(c) \text{nu}(d)$ .
6. **Hence verify that  $c < d$ .**

### Procedure II:18(2.44)

#### Objective

Choose three rational numbers  $a, b, c$  such that  $a < b$ . The objective of the following instructions is to show that  $a + c < b + c$ .

### Implementation

1. Using **declaration II:10**, verify that  $\text{nu}(a) \text{de}(b) < \text{de}(a) \text{nu}(b)$ .
2. Using **declaration II:0**, verify that  $0 < \text{de}(c)$ .
3. Hence verify that  $\text{nu}(a + c) \text{de}(b + c)$ 
  - (a)  $= (\text{nu}(a) \text{de}(c) + \text{de}(a) \text{nu}(c)) \text{de}(b) \text{de}(c)$
  - (b)  $= \text{nu}(a) \text{de}(c) \text{de}(b) \text{de}(c) + \text{de}(a) \text{nu}(c) \text{de}(b) \text{de}(c)$
  - (c)  $< \text{de}(a) \text{de}(c) \text{nu}(b) \text{de}(c) + \text{de}(a) \text{nu}(c) \text{de}(b) \text{de}(c)$
  - (d)  $= (\text{nu}(b) \text{de}(c) + \text{nu}(c) \text{de}(b)) \text{de}(a) \text{de}(c)$
  - (e)  $= \text{nu}(b + c) \text{de}(a + c)$ .
4. **Hence verify that  $a + c < b + c$ .**

### Procedure II:19(2.45)

#### Objective

Choose two rational numbers  $a, b$  such that  $a < b$ . The objective of the following instructions is to show that  $a \neq b$  and  $b \not\prec a$ .

### Implementation

1. Verify that  $\text{nu}(a) \text{de}(b) < \text{de}(a) \text{nu}(b)$ .
2. Hence verify that  $\text{nu}(a) \text{de}(b) \neq \text{de}(a) \text{nu}(b)$ .
3. **Hence verify that  $a \neq b$ .**
4. Also verify that  $\text{nu}(b) \text{de}(a) \not\prec \text{de}(b) \text{nu}(a)$ .
5. **Hence verify that  $b \not\prec a$ .**

### Procedure II:20(2.46)

#### Objective

Choose two rational numbers  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $a \not\prec b$  and  $b \not\prec a$ .

### Implementation

Implementation is analogous to that of **procedure II:19**.



### Procedure II:21(2.47)

#### Objective

Choose two rational numbers  $a, b$  such that  $a \neq b$ . The objective of the following instructions is to show that  $a < b$  or  $b < a$ .

#### Implementation

1. Verify that  $\text{nu}(a) \text{de}(b) \neq \text{de}(a) \text{nu}(b)$ .
2. If  $\text{nu}(a) \text{de}(b) < \text{de}(a) \text{nu}(b)$ , then do the following:
  - (a) **Verify that**  $a < b$ .
3. Otherwise do the following:
  - (a) Verify that  $\text{nu}(b) \text{de}(a) < \text{de}(b) \text{nu}(a)$ .
  - (b) **Hence verify that**  $b < a$ .

### Procedure II:22(2.48)

#### Objective

Choose two rational numbers  $a, b$  such that  $a \not\leq b$ . The objective of the following instructions is to show that  $a = b$  or  $b < a$ .

#### Implementation

Implementation is analogous to that of [procedure II:21](#).

### Procedure II:23(2.49)

#### Objective

Choose two rational numbers  $a, b$  such that  $0 < a$  and  $0 < b$ . The objective of the following instructions is to show that  $0 < a + b$ .

#### Implementation

1. Using [declaration II:10](#), verify that  $0 = \text{nu}(0) \text{de}(a) < \text{de}(0) \text{nu}(a) = \text{nu}(a)$ .
2. Using [declaration II:0](#), verify that  $0 < \text{de}(a)$ .

3. Using [declaration II:10](#), verify that  $0 = \text{nu}(0) \text{de}(b) < \text{de}(0) \text{nu}(b) = \text{nu}(b)$ .
4. Using [declaration II:0](#), verify that  $0 < \text{de}(b)$ .
5. Hence verify that  $\text{nu}(0) \text{de}(a + b) = 0 < \text{nu}(a) \text{de}(b) + \text{de}(a) \text{nu}(b) = \text{de}(0) \text{nu}(a + b)$ .
6. **Hence verify that**  $0 < a + b$ .

### Procedure II:24(2.50)

#### Objective

Choose two rational numbers  $a, b$  such that  $0 < a$  and  $0 < b$ . The objective of the following instructions is to show that  $0 < ab$ .

#### Implementation

1. Using [declaration II:10](#), verify that  $0 = \text{nu}(0) \text{de}(a) < \text{de}(0) \text{nu}(a) = \text{nu}(a)$ .
2. Using [declaration II:10](#), verify that  $0 = \text{nu}(0) \text{de}(b) < \text{de}(0) \text{nu}(b) = \text{nu}(b)$ .
3. Hence verify that  $\text{nu}(0) \text{de}(ab) = 0 < \text{nu}(a) \text{nu}(b) = \text{de}(0) \text{nu}(ab)$ .
4. **Hence verify that**  $0 < ab$ .

### Procedure II:25(2.81)

#### Objective

Choose two rational numbers  $a, b$ . The objective of the following instructions is to show that  $\|ab\| = \|a\| \|b\|$ .

#### Implementation

Implementation is analogous to that of [procedure I:23](#).

### Procedure II:26(2.82)

#### Objective

Choose two rational numbers  $a, b$ . The objective of the following instructions is to show that  $\|a + b\| \leq \|a\| + \|b\|$ .

### Implementation

Implementation is analogous to that of [procedure I:24](#).

### Procedure II:27(2.83)

#### Objective

Choose two rational numbers  $a, b$ . The objective of the following instructions is to show that  $\|a\| - \|b\| \leq \|a - b\|$ .

#### Implementation

Implementation is analogous to that of [procedure I:25](#).

### Procedure II:28(2.84)

#### Objective

Choose a rational number  $a$ . The objective of the following instructions is to show that  $a = \text{sgn}(a)\|a\|$ .

#### Implementation

Implementation is analogous to that of [procedure I:26](#).

### Declaration II:11

The notation  $\lfloor a \rfloor$ , where  $a$  is a rational number, will be used as a shorthand for  $\text{nu}(a) \text{div de}(a)$ .

### Declaration II:12

The notation  $\lceil a \rceil$ , where  $a$  is a rational number, will be used as a shorthand for  $(\text{nu}(a) \text{div de}(a)) + 1$ .

### Procedure II:29(2.04)

#### Objective

Choose a rational number  $r \neq 1$  and an integer  $n \geq 0$ . The objective of the following instructions is to show that  $\sum_t^{[0:n]} r^t = \frac{1-r^{n+1}}{1-r}$ .

#### Implementation

1. Verify that  $r \sum_t^{[0:n]} r^t = \sum_t^{[0:n]} r^{t+1} = \sum_t^{[1:n+1]} r^t$ .
2. Therefore verify that  $(1 - r) \sum_t^{[0:n]} r^t = \sum_t^{[0:n]} r^t - \sum_t^{[1:n+1]} r^t = 1 - r^{n+1}$ .
3. Therefore verify that  $\sum_t^{[0:n]} r^t = \frac{1-r^{n+1}}{1-r}$ .

### Procedure II:30(2.05)

#### Objective

Choose a rational  $0 < r < 1$  and an integer  $n \geq 0$ . The objective of the following instructions is to show that  $\sum_t^{[0:n]} r^t < \frac{1}{1-r}$ .

#### Implementation

1. Using [procedure II:29](#), verify that  $\sum_t^{[0:n]} r^t = \frac{1-r^{n+1}}{1-r} < \frac{1}{1-r}$ .

### Procedure II:31(2.06)

#### Objective

Choose a non-negative integer  $a$  and a rational number  $x$ . The objective of the following instructions is to show that  $(1+x)^a = \sum_r^{[0:a+1]} \binom{a}{r} x^r$ .

#### Implementation

Instructions are analogous to those of [procedure I:87](#).

## Procedure II:32(2.07)

### Objective

Choose an integer  $r \geq 0$  and a rational number  $x \geq -1$ . The objective of the following instructions is to show that  $(1+x)^r \geq 1+rx$ .

### Implementation

1. If  $-1 \leq x < 0$ , then do the following:
  - (a) Using **procedure II:29**, verify that  $(1+x)^r$ 
    - i.  $= 1 + (1+x)^r - 1$
    - ii.  $= 1 + x \frac{(1+x)^r - 1}{(1+x) - 1}$
    - iii.  $= 1 + x \sum_k^{[0:r]} (1+x)^k$
    - iv.  $\geq 1 + x \sum_k^{[0:r]} 1$
    - v.  $= 1 + rx$ .
2. Otherwise, do the following:
  - (a) Verify that  $x \geq 0$ .
  - (b) Using **procedure II:31**, verify that  $(1+x)^r$ 
    - i.  $= \sum_k^{[0:r+1]} \binom{r}{k} x^k$
    - ii.  $\geq \binom{r}{0} x^0 + \binom{r}{1} x^1$
    - iii.  $= 1 + rx$

5. Hence verify that  $(1 - \frac{x}{1+x})^r \geq 1 - \frac{rx}{1+x} > 0$ .

6. Hence verify that  $(1+x)^r$

(a)  $= (\frac{1}{1+x})^{-r}$

(b)  $= (1 - \frac{x}{1+x})^{-r}$

(c)  $\leq (1 - \frac{rx}{1+x})^{-1}$

(d)  $= \frac{1+x}{1-(r-1)x}$ .

## Procedure II:33(wed2407191348)

### Objective

Choose a non-negative integer  $r$  and a rational number  $x > -1$  such that  $(r-1)x < 1$ . The objective of the following instructions is to show that  $(1+x)^r \leq \frac{1+x}{1-(r-1)x}$ .

### Implementation

1. Verify that  $1 - \frac{x}{1+x} = \frac{1}{1+x} > 0$ .
2. Hence using **procedure II:32**, verify that  $(1 - \frac{x}{1+x})^r \geq 1 - \frac{rx}{1+x}$ .
3. Verify that  $0 < 1 + x - rx$ .
4. Hence verify that  $0 < 1 - \frac{rx}{1+x}$ .

## Chapter 6

# Polynomial Arithmetic

### Declaration II:13

The notation  $\text{min}(c)$ , where  $c$  is a list, will be used as a shorthand for  $\infty$  if  $c$  is empty, otherwise it will stand for the minimum entry of  $c$ .

### Declaration II:14

The notation  $\text{min}_r^R c(r)$ , where  $R$  is a list and  $c[r]$  is a function of  $r$ , will be used as a shorthand for  $\text{min}(c(R))$ .

### Declaration II:15

The notation  $\text{max}(c)$ , where  $c$  is a list, will be used as a shorthand for  $-\infty$  if  $c$  is empty, otherwise it will stand for the maximum entry of  $c$ .

### Declaration II:16

The notation  $\text{max}_r^R c(r)$ , where  $R$  is a list and  $c[r]$  is a function of  $r$ , will be used as a shorthand for  $\text{max}(c(R))$ .

### Declaration II:17

The phrase "polynomial" will be used as a shorthand for a list of rational numbers.

### Declaration II:18

The notation  $a_i$ , where  $a$  is a polynomial and  $i$  is a natural number such that  $i \geq |a|$ , will be used as a shorthand for 0.

### Declaration II:19

The phrase " $a = b$ ", where  $a, b$  are polynomials, will be used as a shorthand for " $a_i = b_i$  for each  $i \in [0 : \max(|a|, |b|)]$ ".

### Declaration II:20

The notation  $\Lambda(a, b)$  will be used as a shorthand for  $\sum_r^{[0:|a|]} a_r b^r$ .

## Procedure II:34(2.51)

### Objective

Choose two polynomials  $a, b$  and a rational number  $c$  such that  $a = b$ . The objective of the following instructions is to show that  $\Lambda(a, c) = \Lambda(b, c)$ .

### Implementation

1. Verify that  $\Lambda(a, c)$

$$(a) = \sum_r^{[0:|a|]} a_r c^r$$

$$(b) = \sum_r^{[0:\max(|a|, |b|)]} a_r c^r$$

$$(c) = \sum_r^{[0:\max(|a|, |b|)]} b_r c^r$$

$$(d) = \sum_r^{[0:|b|]} b_r c^r$$

$$(e) = \Lambda(b, c).$$

### Procedure II:35(2.52)

#### Objective

Choose a natural number  $c$  and two polynomials  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $a_c = b_c$ .

#### Implementation

1. If  $c < \max(|a|, |b|)$ , then do the following:
  - (a) **Verify that**  $a_c = b_c$ .
2. Otherwise do the following:
  - (a) Verify that  $c \geq \max(|a|, |b|)$ .
  - (b) **Hence verify that**  $a_c = 0 = b_c$ .

### Procedure II:36(2.53)

#### Objective

Choose a polynomial  $a$ . The objective of the following instructions is to show that  $a = a$ .

#### Implementation

1. Verify that  $a_i = a_i$  for each  $i \in [0 : \max(|a|, |a|)]$ .
2. **Hence verify that**  $a = a$ .

### Procedure II:37(2.54)

#### Objective

Choose two polynomials  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $b = a$ .

#### Implementation

1. Verify that  $a_i = b_i$  for each  $i \in [0 : \max(|a|, |b|)]$ .
2. Hence verify that  $b_i = a_i$  for each  $i \in [0 : \max(|b|, |a|)]$ .
3. **Hence verify that**  $b = a$ .

### Procedure II:38(2.55)

#### Objective

Choose three polynomials  $a, b, c$  such that  $a = b$  and  $b = c$ . The objective of the following instructions is to show that  $a = c$ .

#### Implementation

1. Using **declaration II:19**, verify that  $a_i = b_i$  for each  $i \in [0 : \max(|a|, |b|, |c|)]$ .
2. Using **declaration II:19**, verify that  $b_i = c_i$  for each  $i \in [0 : \max(|a|, |b|, |c|)]$ .
3. Hence verify that  $a_i = c_i$  for each  $i \in [0 : \max(|a|, |b|, |c|)]$ .
4. **Hence verify that**  $a = c$ .

#### Declaration II:21

The notation  $\langle f(j) \text{ for } j \in R \rangle$ , where  $f[j]$  is a function of  $j$  and  $R$  is a list, will be used as a shorthand for  $\langle f(R) \rangle$ .

#### Declaration II:22

The notation  $a + b$ , where  $a, b$  are polynomials, will be used as a shorthand for the list  $\langle a_i + b_i \text{ for } i \in [0 : \max(|a|, |b|)] \rangle$ .

### Procedure II:39(2.56)

#### Objective

Choose two polynomials  $a, b$  and a rational number  $c$ . The objective of the following instructions is to show that  $\Lambda(a + b, c) = \Lambda(a, c) + \Lambda(b, c)$ .

### Implementation

1. Verify that  $\Lambda(a + b, c)$ 
  - (a)  $= \Lambda(\langle a_r + b_r \text{ for } r \in [0 : \max(|a|, |b|)] \rangle, c)$
  - (b)  $= \sum_r^{[0 : \max(|a|, |b|)]} (a_r + b_r) c^r$
  - (c)  $= \sum_r^{[0 : \max(|a|, |b|)]} a_r c^r + \sum_r^{[0 : \max(|a|, |b|)]} b_r c^r$
  - (d)  $= \sum_r^{[0 : |a|]} a_r c^r + \sum_r^{[0 : |b|]} b_r c^r$
  - (e)  $= \Lambda(a, c) + \Lambda(b, c).$

### Procedure II:40(2.57)

#### Objective

Choose a natural number  $c$  and two polynomials  $a, b$ . The objective of the following instructions is to show that  $(a + b)_c = a_c + b_c$ .

#### Implementation

1. If  $c < \max(|a|, |b|)$ , then do the following:
  - (a) **Verify that**  $(a + b)_c = a_c + b_c$ .
2. Otherwise do the following:
  - (a) Verify that  $c \geq \max(|a|, |b|)$ .
  - (b) Hence verify that  $a_c = 0$ .
  - (c) Also verify that  $b_c = 0$ .
  - (d) Also verify that  $(a + b)_c = 0$ .
  - (e) **Hence verify that**  $(a + b)_c = a_c + b_c$ .

### Procedure II:41(2.58)

#### Objective

Choose four polynomials  $a, b, c, d$  such that  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $a + b = c + d$ .

#### Implementation

1. Verify that  $a_i = c_i$  for each  $i \in [0 : \max(|a|, |b|, |c|, |d|)]$ .

2. Verify that  $b_i = d_i$  for each  $i \in [0 : \max(|a|, |b|, |c|, |d|)]$ .

3. Hence verify that  $a + b$

- (a)  $= \langle a_i + b_i \text{ for } i \in [0 : \max(|a|, |b|, |c|, |d|)] \rangle$
- (b)  $= \langle c_i + d_i \text{ for } i \in [0 : \max(|a|, |b|, |c|, |d|)] \rangle$
- (c)  $= c + d.$

### Procedure II:42(2.59)

#### Objective

Choose three polynomials  $a, b, c$ . The objective of the following instructions is to show that  $(a + b) + c = a + (b + c)$ .

#### Implementation

1. Verify that  $(a + b) + c$ 
  - (a)  $\langle (a + b)_i + c_i \text{ for } i \in [0 : \max(|a + b|, |c|)] \rangle$
  - (b)  $\langle (a_i + b_i) + c_i \text{ for } i \in [0 : \max(|a|, |b|, |c|)] \rangle$
  - (c)  $\langle a_i + (b_i + c_i) \text{ for } i \in [0 : \max(|a|, |b + c|)] \rangle$
  - (d)  $\langle a_i + (b + c)_i \text{ for } i \in [0 : \max(|a|, |b + c|)] \rangle$
  - (e)  $= a + (b + c).$

### Procedure II:43(2.60)

#### Objective

Choose two polynomials  $a, b$ . The objective of the following instructions is to show that  $a + b = b + a$ .

#### Implementation

1. Verify that  $a + b$ 
  - (a)  $= \langle a_i + b_i \text{ for } i \in [0 : \max(|a|, |b|)] \rangle$
  - (b)  $= \langle b_i + a_i \text{ for } i \in [0 : \max(|b|, |a|)] \rangle$
  - (c)  $= b + a.$

### Declaration II:23

The notation  $\textcolor{red}{a}$ , where  $a$  is a rational number, will contextually be used as a shorthand for the list  $\langle a \rangle$ .

## Procedure II:44(2.61)

### Objective

Choose a polynomial  $a$ . The objective of the following instructions is to show that  $0 + a = a$ .

### Implementation

1. Verify that  $0 + a$ 
  - (a)  $= \langle 0_i + a_i \text{ for } i \in [0 : |a|] \rangle$
  - (b)  $= \langle 0 + a_i \text{ for } i \in [0 : |a|] \rangle$
  - (c)  $= a$ .

### Declaration II:24

The notation  $-a$ , where  $a$  is a polynomial, will be used as a shorthand for the list  $\langle -a_i \text{ for } i \in [0 : |a|] \rangle$ .

## Procedure II:45(2.00)

### Objective

Choose a polynomial  $a$  and a rational number  $b$ . The objective of the following instructions is to show that  $\Lambda(-a, b) = -\Lambda(a, b)$ .

### Implementation

1. Verify that  $\Lambda(-a, b)$ 
  - (a)  $= \Lambda(\langle -a_i \text{ for } i \in [0 : |a|] \rangle, b)$
  - (b)  $= \sum_j^{[0:|a|]} (-a_j) b^j$
  - (c)  $= - \sum_j^{[0:|a|]} a_j b^j$
  - (d)  $= -\Lambda(a, b)$ .

## Procedure II:46(2.62)

### Objective

Choose two polynomials  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $-a = -b$ .

### Implementation

1. Verify that  $a_i = b_i$  for  $i \in [0 : \max(|a|, |b|)]$ .
2. Hence verify that  $-a$ 
  - (a)  $= \langle -a_i \text{ for } i \in [0 : \max(|a|, |b|)] \rangle$
  - (b)  $= \langle -b_i \text{ for } i \in [0 : \max(|a|, |b|)] \rangle$
  - (c)  $= -b$ .

## Procedure II:47(2.63)

### Objective

Choose a polynomial  $a$ . The objective of the following instructions is to show that  $-a + a = 0$ .

### Implementation

1. Verify that  $-a + a$ 
  - (a)  $= (-a) + a$
  - (b)  $= \langle -a_i \text{ for } i \in [0 : |a|] \rangle + \langle a_i \text{ for } i \in [0 : |a|] \rangle$
  - (c)  $= \langle -a_i + a_i \text{ for } i \in [0 : |a|] \rangle$
  - (d)  $= \langle 0 \text{ for } i \in [0 : |a|] \rangle$
  - (e)  $= 0$ .

### Declaration II:25

The notation  $ab$ , where  $a, b$  are integers, will be used as a shorthand for the list  $\langle \sum_r^{[0:i+1]} a_r b_{i-r} \text{ for } i \in [0 : |a| + |b| - 1] \rangle$ .

## Procedure II:48(2.64)

### Objective

Choose two polynomials  $a, b$  and a rational number  $c$ . The objective of the following instructions is to show that  $\Lambda(ab, c) = \Lambda(a, c)\Lambda(b, c)$ .

## Implementation

1. Verify that  $\Lambda(ab, c)$

- (a)  $= \Lambda(\langle \sum_r^{[0:j+1]} a_r b_{j-r} \text{ for } j \in [0 : |a| + |b| - 1] \rangle, c)$
- (b)  $= \sum_j^{[0:|a|+|b|-1]} (\sum_r^{[0:j+1]} a_r b_{j-r}) c^j$
- (c)  $= \sum_j^{[0:|a|+|b|-1]} \sum_r^{[0:j+1]} a_r c^r b_{j-r} c^{j-r}$
- (d)  $= \sum_r^{[0:|a|+|b|-1]} \sum_j^{[r:|a|+|b|-1]} a_r c^r b_{j-r} c^{j-r}$
- (e)  $= \sum_r^{[0:|a|+|b|-1]} a_r c^r \sum_j^{[r:|a|+|b|-1]} b_{j-r} c^{j-r}$
- (f)  $= \sum_r^{[0:|a|+|b|-1]} a_r c^r \sum_j^{[0:|a|+|b|-1-r]} b_j c^j$
- (g)  $= \sum_r^{[0:|a|]} a_r c^r \sum_j^{[0:|a|+|b|-1-r]} b_j c^j$
- (h)  $= \sum_r^{[0:|a|]} a_r c^r \sum_j^{[0:|b|]} b_j c^j$
- (i)  $= (\sum_j^{[0:|a|]} a_j c^j) (\sum_j^{[0:|b|]} b_j c^j)$
- (j)  $= \Lambda(a, c) \Lambda(b, c).$

## Procedure II:49(2.65)

### Objective

Choose a natural number  $c$  and two polynomials  $a, b$ . The objective of the following instructions is to show that  $(ab)_c = \sum_r^{[0:c+1]} a_r b_{c-r}$ .

### Implementation

1. If  $c < |a| + |b| - 1$ , then do the following:

- (a) **Verify that**  $(ab)_c = \sum_r^{[0:c+1]} a_r b_{c-r}$ .

2. Otherwise do the following:

- (a) Verify that  $c \geq |a| + |b| - 1$ .
- (b) Verify that  $(ab)_c$ 
  - i.  $= 0$
  - ii.  $= \sum_r^{[0:|a|]} 0a_r + \sum_r^{[|a|:c+1]} 0b_{c-r}$
  - iii.  $= \sum_r^{[0:|a|]} a_r b_{c-r} + \sum_r^{[|a|:c+1]} a_r b_{c-r}$
  - iv.  $= \sum_r^{[0:c+1]} a_r b_{c-r}.$

## Procedure II:50(2.66)

### Objective

Choose four polynomials  $a, b, c, d$  such that  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $ab = cd$ .

### Implementation

- 1. Using **declaration II:19**, verify that  $a_i = c_i$  for  $i \in [0 : \max(|a|, |c|) + \max(|b|, |d|) - 1]$ .
- 2. Using **declaration II:19**, verify that  $b_i = d_i$  for  $i \in [0 : \max(|a|, |c|) + \max(|b|, |d|) - 1]$ .
- 3. Hence verify that  $ab$ 
  - (a)  $= \langle \sum_r^{[0:i+1]} a_r b_{i-r} \text{ for } i \in [0 : \max(|a|, |c|) + \max(|b|, |d|) - 1] \rangle$
  - (b)  $= \langle \sum_r^{[0:i+1]} c_r d_{i-r} \text{ for } i \in [0 : \max(|a|, |c|) + \max(|b|, |d|) - 1] \rangle$
  - (c)  $= cd.$

## Procedure II:51(2.67)

### Objective

Choose three polynomials  $a, b, c$ . The objective of the following instructions is to show that  $(ab)c = a(bc)$ .

### Implementation

1. Verify that  $(ab)c$

- (a)  $= \langle \sum_t^{[0:j+1]} (ab)_t c_{j-t} \text{ for } j \in [0 : |ab| + |c| - 1] \rangle$
- (b)  $= \langle \sum_t^{[0:j+1]} \langle \sum_r^{[0:i+1]} a_r b_{i-r} \text{ for } i \in [0 : |a| + |b| - 1] \rangle_t c_{j-t} \text{ for } j \in [0 : |a| + |b| + |c| - 2] \rangle$
- (c)  $= \langle \sum_t^{[0:j+1]} \sum_r^{[0:t+1]} a_r b_{t-r} c_{j-t} \text{ for } j \in [0 : |a| + |b| + |c| - 2] \rangle$
- (d)  $= \langle \sum_r^{[0:j+1]} \sum_t^{[r:j+1]} a_r b_{t-r} c_{j-t} \text{ for } j \in [0 : |a| + |b| + |c| - 2] \rangle$
- (e)  $= \langle \sum_r^{[0:j+1]} a_r \sum_t^{[r:j+1]} b_{t-r} c_{j-t} \text{ for } j \in [0 : |a| + |b| + |c| - 2] \rangle$



- (f) =  $\langle \sum_r^{[0:j+1]} a_r \sum_t^{[0:j-r+1]} b_t c_{j-r-t} \text{ for } j \in [0 : |a| + |b| + |c| - 2] \rangle$
- (g) =  $\langle \sum_r^{[0:j+1]} a_r \langle \sum_t^{[0:i+1]} b_t c_{i-t} \text{ for } i \in [0 : |b| + |c| - 1] \rangle_{j-r} \text{ for } j \in [0 : |a| + |b| + |c| - 2] \rangle$
- (h) =  $\langle \sum_r^{[0:j+1]} a_r (bc)_{j-r} \text{ for } j \in [0 : |a| + |bc| - 1] \rangle$
- (i) =  $a(bc)$ .

## Procedure II:52(2.68)

### Objective

Choose two polynomials  $a, b$ . The objective of the following instructions is to show that  $ab = ba$ .

### Implementation

1.  $ab$
- (a) =  $\langle \sum_r^{[0:i+1]} a_r b_{i-r} \text{ for } i \in [0 : |a| + |b| - 1] \rangle$
- (b) =  $\langle \sum_r^{[0:i+1]} b_r a_{i-r} \text{ for } i \in [0 : |a| + |b| - 1] \rangle$
- (c) =  $ba$ .

## Procedure II:53(2.69)

### Objective

Choose a polynomial  $a$ . The objective of the following instructions is to show that  $1a = a$ .

### Implementation

1. Verify that  $1a$
- (a) =  $\langle \sum_r^{[0:i+1]} 1_r a_{i-r} \text{ for } i \in [0 : |1| + |a| - 1] \rangle$
- (b) =  $\langle 1_0 a_{i-0} \text{ for } i \in [0 : |a|] \rangle$
- (c) =  $\langle a_i \text{ for } i \in [0 : |a|] \rangle$
- (d) =  $a$ .

## Procedure II:54(2.70)

### Objective

Choose three polynomials  $a, b, c$ . The objective of the following instructions is to show that  $a(b + c) = ab + ac$ .

### Implementation

1.  $a(b + c)$
- (a) =  $\langle \sum_r^{[0:i+1]} a_r (b + c)_{i-r} \text{ for } i \in [0 : |a| + |b + c| - 1] \rangle$
- (b) =  $\langle \sum_r^{[0:i+1]} a_r (b_{i-r} + c_{i-r}) \text{ for } i \in [0 : |a| + |b + c| - 1] \rangle$
- (c) =  $\langle \sum_r^{[0:i+1]} (a_r b_{i-r} + a_r c_{i-r}) \text{ for } i \in [0 : |a| + |b + c| - 1] \rangle$
- (d) =  $\langle \sum_r^{[0:i+1]} a_r b_{i-r} + \sum_r^{[0:i+1]} a_r c_{i-r} \text{ for } i \in [0 : |a| + |b + c| - 1] \rangle$
- (e) =  $\langle \sum_r^{[0:i+1]} a_r b_{i-r} \text{ for } i \in [0 : |a| + |b| - 1] \rangle + \langle \sum_r^{[0:i+1]} a_r c_{i-r} \text{ for } i \in [0 : |a| + |c| - 1] \rangle$
- (f) =  $ab + ac$ .

### Declaration II:26

The notation  $\lambda$  will be used as a shorthand for the list  $\langle 0, 1 \rangle$ .

## Procedure II:55(2.71)

### Objective

Choose a polynomial  $a$ . The objective of the following instructions is to show that  $\lambda a = \langle 0 \rangle \cap a$ .

### Implementation

1. Verify that  $|\lambda a| = |\lambda| + |a| - 1 = |a| + 1$ .
2. For  $j \in [1 : |a| + 1]$ , do the following:
  - (a) Verify that  $(\lambda a)_j$ 
    - i. =  $\sum_r^{[0:j+1]} \lambda_r a_{j-r}$
    - ii. =  $\sum_r^{[0:j+1]} [r = 1] a_{j-r}$

$$\text{iii. } = a_{j-1}$$

$$3. \text{ Verify that } (\lambda a)_0 = \sum_r^{[0:1]} \lambda_r a_{0-r} = \lambda_0 a_0 = 0.$$

$$4. \text{ Hence verify that } \lambda a = \langle 0 \rangle \frown a.$$

## Procedure II:56(2.72)

### Objective

Choose a natural number  $n$ . The objective of the following instructions is to show that  $\lambda^n = \langle [j = n] \text{ for } j \in [0 : n + 1] \rangle$ .

### Implementation

1. If  $n = 0$ , then do the following:
  - (a) Verify that  $\lambda^n$ 
    - i.  $= \lambda^0$
    - ii.  $= \langle 1 \rangle$
    - iii.  $= \langle [j = 0] \text{ for } j \in [0 : 1] \rangle$
    - iv.  $= \langle [j = n] \text{ for } j \in [0 : n + 1] \rangle$ .
2. Otherwise do the following:
  - (a) Execute **procedure II:63** on  $\langle n - 1 \rangle$ .
  - (b) Hence verify that  $\lambda^{n-1} = \langle [j = n - 1] \text{ for } j \in [0 : n] \rangle$ .
  - (c) Hence verify that  $\lambda^n$ 
    - i.  $= \lambda \lambda^{n-1}$
    - ii.  $= \lambda \langle [j = n - 1] \text{ for } j \in [0 : n] \rangle$
    - iii.  $= \langle 0 \rangle \frown \langle [j = n - 1] \text{ for } j \in [0 : n] \rangle$
    - iv.  $= \langle [j = n] \text{ for } j \in [0 : n + 1] \rangle$ .

### Declaration II:27

The notation **deg**( $a$ ), where  $a$  is a polynomial such that  $a \neq 0$ , will be used as a shorthand for the largest natural number  $j < |a|$  such that  $a_j \neq 0$ .

## Procedure II:57(2.73)

### Objective

Choose two polynomials  $a, b$  such that  $a \neq b$  and  $a \neq 0$ . The objective of the following instructions is to show that  $\deg(a) = \deg(b)$ .

### Implementation

1. For  $j \in [\max(|a|, |b|) : 0]$ , do the following:
  - (a) If  $a_j = 0$ , then do the following:
    - i. Verify that  $0 = a_j = b_j$ .
  - (b) Otherwise do the following:
    - i. Verify that  $0 \neq a_j = b_j$ .
    - ii. Hence verify that  $j < \min(|a|, |b|)$ .
- iii. **Hence verify that**  $\deg(a) = j = \deg(b)$ .

## Procedure II:58(2.74)

### Objective

Let  $\deg(0) = -1$ . Choose two polynomials  $a, b$  such that  $\deg(a) < \deg(b)$ . The objective of the following instructions is to show that  $\deg(a + b) = \deg(b)$ .

### Implementation

1. For  $j \in [\max(|a|, |b|) : \deg(b) + 1]$ , do the following:
  - (a) Verify that  $j > \deg(b) > \deg(a)$ .
  - (b) Hence verify that  $a_j = b_j = 0$ .
  - (c) Hence verify that  $(a + b)_j = a_j + b_j = 0$ .
2. Verify that  $\deg(b) > \deg(a)$ .
3. Hence verify that  $(a + b)_{\deg(b)} = a_{\deg(b)} + b_{\deg(b)} = 0 + b_{\deg(b)} = b_{\deg(b)} \neq 0$ .
4. **Hence verify that**  $\deg(a + b) = \deg(b)$ .

## Procedure II:59(2.75)

### Objective

Let  $\deg(0) = -1$ . Choose two polynomials  $a, b$ . The objective of the following instructions is to show that  $\deg(a + b) \leq \max(\deg(a), \deg(b))$ .

### Implementation

1. For  $j \in [\max(|a|, |b|) : \max(\deg(a), \deg(b)) + 1]$ , do the following:
  - (a) Verify that  $j > \deg(a)$ .
  - (b) Verify that  $j > \deg(b)$ .
  - (c) Hence verify that  $a_j = b_j = 0$ .
  - (d) Hence verify that  $(a + b)_j = a_j + b_j = 0$ .
2. **Hence verify that  $\deg(a + b) \leq \max(\deg(a), \deg(b))$ .**

## Procedure II:60(2.76)

### Objective

Let  $\deg(0) = -1$ . Choose a polynomial  $a$ . The objective of the following instructions is to show that  $\deg(-a) = \deg(a)$ .

### Implementation

1. For  $j \in [|a| : \deg(a) + 1]$ , do the following:
  - (a) Verify that  $j > \deg(a)$ .
  - (b) Hence verify that  $a_j = 0$ .
  - (c) Hence verify that  $(-a)_j = -(a_j) = -0 = 0$ .
2. Verify that  $a_{\deg(a)} \neq 0$ .
3. Hence verify that  $(-a)_{\deg(a)} = -(a_{\deg(a)}) \neq 0$ .
4. **Hence verify that  $\deg(-a) = \deg(a)$ .**

## Procedure II:61(2.77)

### Objective

Choose two polynomials  $a, b$  such that  $a \neq 0$  and  $b \neq 0$ . The objective of the following instructions is to show that  $(ab)_{\deg(a)+\deg(b)} = a_{\deg(a)}b_{\deg(b)} \neq 0$ .

### Implementation

1. Verify that  $a_{\deg(a)} \neq 0$ .
2. Verify that  $b_{\deg(b)} \neq 0$ .
3. Hence verify that  $(ab)_{\deg(a)+\deg(b)}$ 
  - (a)  $= \sum_r^{[0:\deg(a)+\deg(b)+1]} a_r b_{\deg(a)+\deg(b)-r}$
  - (b)  $= \sum_r^{[0:\deg(a)]} a_r b_{\deg(a)+\deg(b)-r} + a_{\deg(a)} b_{\deg(a)+\deg(b)-\deg(a)} + \sum_r^{[\deg(a)+1:\deg(a)+\deg(b)+1]} a_r b_{\deg(a)+\deg(b)-r}$
  - (c)  $= \sum_r^{[0:\deg(a)]} 0a_r + a_{\deg(a)} b_{\deg(b)} + \sum_r^{[\deg(a)+1:\deg(a)+\deg(b)+1]} 0b_{\deg(a)+\deg(b)-r}$
  - (d)  $= a_{\deg(a)} b_{\deg(b)}$
  - (e)  $\neq 0$ .

## Procedure II:62(2.78)

### Objective

Choose two polynomials  $a, b$  such that  $a \neq 0$  and  $b \neq 0$ . The objective of the following instructions is to show that  $\deg(ab) = \deg(a) + \deg(b)$ .

### Implementation

1. For  $j \in [\deg(a) + \deg(b) + 1 : |a| + |b| - 1]$ , do the following:
  - (a) Verify that  $(ab)_j$ 
    - i.  $= \sum_r^{[0:j+1]} a_r b_{j-r}$
    - ii.  $= \sum_r^{[0:\deg(a)+1]} a_r b_{j-r} + \sum_r^{[\deg(a)+1:j+1]} a_r b_{j-r}$
    - iii.  $= \sum_r^{[0:\deg(a)+1]} 0a_r + \sum_r^{[\deg(a)+1:j+1]} 0b_{j-r}$
    - iv.  $= 0$ .
2. Now using **procedure II:61**, verify that  $(ab)_{\deg(a)+\deg(b)} = a_{\deg(a)}b_{\deg(b)} \neq 0$ .

3. Hence verify that  $\deg(ab) = \deg(a) + \deg(b)$ .

#### Declaration II:28

The phrase "monic polynomial" will be used to refer to polynomials  $p$  such that  $p \neq 0$  and  $p_{\deg(p)} = 1$ .

#### Declaration II:29

The notation  $\text{mon}(p)$ , where  $p$  is a polynomial such that  $p \neq 0$ , will be used as a shorthand for  $\frac{p}{p_{\deg(p)}}$ .

#### Procedure II:63(2.25)

##### Objective

Choose two polynomials,  $a, b$  such that  $b \neq 0$ . The objective of the following instructions is to construct two polynomials  $u, w$  such that  $a = ub + w$  and  $\deg(w) < \deg(b)$ .

##### Implementation

1. If  $\deg(a) \geq \deg(b)$ , then do the following:
  - (a) Let  $y = \frac{a_{\deg(a)}}{b_{\deg(b)}} \lambda^{\deg(a) - \deg(b)}$
  - (b) Let  $e = a - yb$ .
  - (c) Verify that  $\deg(e) < \deg(a)$ .
  - (d) Execute **procedure II:63** on the tuple  $\langle e, b \rangle$  and let  $\langle c, d \rangle$  receive.
  - (e) Verify that  $cb + d = e$ .
  - (f) **Verify that**  $\deg(d) < \deg(b)$ .
  - (g) Therefore verify that  $cb + d = a - yb$
  - (h) **Therefore verify that**  $(y + c)b + d = a$ .
  - (i) **Now yield the tuple**  $\langle y + c, d \rangle$ .
2. Otherwise do the following:
  - (a) **Verify that**  $0b + a = a$ .
  - (b) **Verify that**  $\deg(a) < \deg(b)$ .
  - (c) **Yield the tuple**  $\langle 0, a \rangle$ .

#### Declaration II:30

The notation  $a \text{ div } b$ , where  $a, b$  are polynomials, will be used to refer to the first part of the pair yielded by executing **procedure II:63** on  $\langle a, b \rangle$ .

#### Declaration II:31

The notation  $a \text{ mod } b$ , where  $a, b$  are polynomials, will be used to refer to the second part of the pair yielded by executing **procedure II:63** on  $\langle a, b \rangle$ .

#### Procedure II:64(2.79)

##### Objective

Choose a polynomial  $a$  and a rational number  $b$ . The objective of the following instructions is to show that  $a \text{ mod } (\lambda - b) = \Lambda(a, b)$ .

##### Implementation

1. Let  $d = \lambda - b$ .
2. Verify that  $d \neq 0$ .
3. Let  $c = a \text{ div } d$ .
4. Verify that  $a = cd + (a \text{ mod } d)$ .
5. Also verify that  $\deg(a \text{ mod } d) < \deg(d) = 1$ .
6. Hence verify that  $\deg(a \text{ mod } d) = 0$ .
7. Now verify that  $\Lambda(a, b)$ 
  - (a)  $= \Lambda(cd + (a \text{ mod } d), b)$
  - (b)  $= \Lambda(cd, b) + \Lambda(a \text{ mod } d, b)$
  - (c)  $= \Lambda(c, b)\Lambda(d, b) + \Lambda(a \text{ mod } d, b)$
  - (d)  $= \Lambda(c, b)(-b + b) + \Lambda(a \text{ mod } d, b)$
  - (e)  $= 0\Lambda(c, b) + \Lambda(a \text{ mod } d, b)$
  - (f)  $= \Lambda(a \text{ mod } d, b)$
  - (g)  $= a \text{ mod } d$ .

# Chapter 7

## Polynomial Sign Changes

### Procedure II:65(2.80)

#### Objective

Choose a polynomial  $p \neq 0$  and rational numbers  $a_0 < a_1 < \dots < a_{\deg(p)-2} < a_{\deg(p)-1}$  in such a way that  $\Lambda(p, a_i) = 0$  for  $i \in [0 : \deg(p)]$ . The objective of the following instructions is to either show that  $p = q_0 \prod_j^{[0:n]} (\lambda - a_j)$  or  $0 \neq 0$ .

#### Implementation

1. Let  $n = \deg(p)$ .
2. Let  $q = p$ .
3. For  $i$  in  $[0 : n]$ , do the following:
  - (a) Verify that  $p = q \prod_k^{[0:i]} (\lambda - a_k)$ .
  - (b) If  $\Lambda(q, a_i) \neq 0$ , do the following:
    - i. Verify that  $\Lambda(p, a_i) = \Lambda(q \prod_k^{[0:i]} (\lambda - a_k), a_i) = \Lambda(q, a_i) \prod_k^{[0:i]} \Lambda(\lambda - a_k, a_i) = \Lambda(q, a_i) \prod_k^{[0:i]} (a_i - a_k) \neq 0$ .
    - ii. Therefore using the precondition and (i), verify that  $0 \neq 0$ .
    - iii. **Abort procedure.**
  - (c) Otherwise do the following:
    - i. Let  $b = q$ .
    - ii. Let  $q = b \operatorname{div}(\lambda - a_i)$ .
    - iii. Verify that  $\Lambda(b, a_i) = 0$ .
    - iv. Execute **procedure II:64** on  $\langle b, a_i \rangle$ .

v. Hence verify that  $b = (\lambda - a_i)q + b \bmod (\lambda - a_i) = (\lambda - a_i)q$ .

vi. Hence verify that  $p = q \prod_j^{[0:i+1]} (\lambda - a_j)$ .

4. Now verify that  $0 \neq p = q \prod_j^{[0:n]} (\lambda - a_j)$ .

5. Hence verify that  $q \neq 0$ .

6. Hence verify that  $n = \deg(p) = \deg(q) + \sum_j^{[0:n]} \deg(\lambda - a_j) = \deg(q) + n$ .

7. Hence verify that  $\deg(q) = 0$ .

8. Hence verify that  $q = q_0 \neq 0$ .

9. **Hence verify that**  $p = q_0 \prod_j^{[0:n]} (\lambda - a_j)$ .

### Procedure II:66(2.16)

#### Objective

Choose a polynomial  $p \neq 0$  and rational numbers  $a_0 < a_1 < \dots < a_{\deg(p)-1} < a_{\deg(p)}$  in such a way that  $\Lambda(p, a_i) = 0$  for  $i \in [0 : \deg(p) + 1]$ . The objective of the following instructions is to show that  $0 \neq 0$ .

#### Implementation

1. Let  $n = \deg(p)$ .
2. Execute **procedure II:65** on  $\langle p, a \rangle$ .
3. Hence verify that  $p = q_0 \prod_j^{[0:n]} (\lambda - a_j)$ .

4. Hence verify that  $\Lambda(p, a_n) = \Lambda(q_0 \prod_j^{[0:n]} (\lambda - a_j), a_n) = \Lambda(q_0, a_n) \prod_j^{[0:n]} \Lambda(\lambda - a_j, a_n) = q_0 \prod_j^{[0:n]} (a_n - a_j) \neq 0$ .
5. Therefore using the precondition and (4), verify that  $0 = \Lambda(p, a_n) \neq 0$ .
6. **Abort procedure.**

## Procedure II:67(2.15)

### Objective

Choose a polynomial  $p$  and a rational number  $X$ . The objective of the following instructions is to construct a rational number  $a$  and a procedure  $q(y, z)$  to show that  $|\Lambda(p, z) - \Lambda(p, y)| \leq a|z - y|$  when two rational numbers  $y, z$  such that  $|y| \leq X$  and  $|z| \leq X$  are chosen.

### Implementation

1. Let  $a = \sum_r^{[1:|p|]} r|p_r|X^{r-1}$ .
2. Let  $q(y, z)$  be the following procedure:
  - (a) Verify that  $|\Lambda(p, z) - \Lambda(p, y)|$ 
    - i.  $= |(\sum_r^{[0:|p|]} p_r z^r) - (\sum_r^{[0:|p|]} p_r y^r)|$
    - ii.  $= |\sum_r^{[1:|p|]} p_r(z^r - y^r)|$
    - iii.  $= |\sum_r^{[1:|p|]} p_r(z - y) \sum_t^{[0:r]} z^t y^{r-1-t}|$
    - iv.  $= |(z - y) \sum_r^{[1:|p|]} p_r \sum_t^{[0:r]} z^t y^{r-1-t}|$
    - v.  $= |z - y| |\sum_r^{[1:|p|]} p_r \sum_t^{[0:r]} z^t y^{r-1-t}|$
    - vi.  $\leq |z - y| \sum_r^{[1:|p|]} |p_r| \sum_t^{[0:r]} z^t y^{r-1-t}|$
    - vii.  $= |z - y| \sum_r^{[1:|p|]} |p_r| |\sum_t^{[0:r]} z^t y^{r-1-t}|$
    - viii.  $\leq |z - y| \sum_r^{[1:|p|]} |p_r| \sum_t^{[0:r]} |z^t y^{r-1-t}|$
    - ix.  $= |z - y| \sum_r^{[1:|p|]} |p_r| \sum_t^{[0:r]} |z|^t |y|^{r-1-t}$
    - x.  $\leq |z - y| \sum_r^{[1:|p|]} |p_r| \sum_t^{[0:r]} X^t X^{r-1-t}$
    - xi.  $= |z - y| \sum_r^{[1:|p|]} |p_r| \sum_t^{[0:r]} X^{r-1}$
    - xii.  $= |z - y| \sum_r^{[1:|p|]} r|p_r|X^{r-1}$
    - xiii.  $= a|z - y|$
3. **Yield the tuple  $\langle a, q \rangle$ .**

## Procedure II:68(2.17)

### Objective

Choose a polynomial  $f$ . Choose rational numbers  $a < b$  such that  $\text{sgn}(\Lambda(f, a)) = -\text{sgn}(\Lambda(f, b))$ . Choose a rational number target  $B > 0$ . The objective of the following instructions is to construct a rational number  $d$  such that  $a \leq d \leq b$  and  $|f(d)| < B$ .

### Implementation

1. Execute **procedure II:67** on  $\langle f, \max(|a|, |b|) \rangle$  and let  $\langle G, q \rangle$  receive the result.
2. Let  $c = a$  and  $d = b$ .
3. Until  $G|d - c| < B$ 
  - (a) Let  $e = \frac{c+d}{2}$ .
  - (b) **Verify that  $a \leq c < e < d \leq b$ .**
  - (c) If  $\text{sgn}(\Lambda(f, c)) = -\text{sgn}(\Lambda(f, e))$ , then do the following:
    - i. Let  $d = e$ .
  - (d) Otherwise if  $\text{sgn}(\Lambda(f, e)) = -\text{sgn}(\Lambda(f, d))$ , then do the following:
    - i. Let  $c = e$ .
  - (e) Otherwise if  $\Lambda(f, e) = 0$ , then do the following:
    - i. **Verify that  $|\Lambda(f, e)| = 0 < B$ .**
    - ii. **Yield the tuple  $\langle e \rangle$ .**
4. Execute procedure  $q$  on  $\langle c, d \rangle$ .
5. **Hence verify that  $|\Lambda(f, c)| < |\Lambda(f, d) - \Lambda(f, c)| \leq G|d - c| < B$ .**
6. **Yield the tuple  $\langle c \rangle$ .**

## Procedure II:69(2.18)

### Objective

Choose a polynomial  $f \neq 0$  and pairs of rational numbers  $(a_{\deg(f)}, b_{\deg(f)}), (a_{\deg(f)-1}, b_{\deg(f)-1}), \dots, (a_0, b_0)$  in such a way that:

1.  $a_{\deg(f)} < b_{\deg(f)} \leq a_{\deg(f)-1} < b_{\deg(f)-1} \leq \dots \leq a_1 < b_1 \leq a_0 < b_0$ .

2.  $\text{sgn}(\Lambda(f, a_i)) = -\text{sgn}(\Lambda(f, b_i))$  for  $i \in [0 : \deg(f) + 1]$ .

The objective of the following instructions is to show that  $1 = -1$ .

### Implementation

1. If  $\deg(f) > 0$ :
  - (a) Let  $B = \min_k^{[0:\deg(f)-1]} \min(|\Lambda(f, a_k)|, |\Lambda(f, b_k)|)$ .
  - (b) For  $k \in [0 : \deg(f)]$ , verify that  $|\Lambda(f, a_k)| \geq B$ .
  - (c) Execute **procedure II:68** on the formal polynomial  $f$ , interval  $(a_{\deg(f)}, b_{\deg(f)})$ , and target of  $B$ . Let the tuple  $\langle d \rangle$  receive the result.
  - (d) Verify that  $|\Lambda(f, d)| < B$ .
  - (e) Let  $h = f \text{ div } (\lambda - d)$ .
  - (f) Execute **procedure II:64** on  $\langle f, d \rangle$ .
  - (g) Hence verify that  $f = (\lambda - d)h + f \bmod (\lambda - d) = (\lambda - d)h + \Lambda(f, d)$ .
  - (h) Hence verify that  $0 \neq f - \Lambda(f, d) = (\lambda - d)h$ .
  - (i) Hence verify that  $h \neq 0$ .
  - (j) Hence verify that  $\deg(f) = \deg(f - \Lambda(f, d)) = \deg((\lambda - d)h) = \deg(\lambda - d) + \deg(h) = 1 + \deg(h)$ .
  - (k) Hence verify that  $\deg(h) = \deg(f) - 1$ .
  - (l) For  $k \in [0 : \deg(h) + 1]$ , do the following:
    - i. If  $\Lambda(f, a_k) \geq B$ , in-order verify that:
      - A.  $\Lambda(f, a_k) \geq B > |\Lambda(f, d)| \geq \Lambda(f, d)$ .
      - B.  $\Lambda(f, a_k) - \Lambda(f, d) > 0$ .
      - C.  $(a_k - d)\Lambda(h, a_k) > 0$ .
      - D.  $\Lambda(h, a_k) > 0$ .
      - E.  $\Lambda(f, b_k) \leq -B < -|\Lambda(f, d)| \leq \Lambda(f, d)$ .
      - F.  $\Lambda(f, b_k) - \Lambda(f, d) < 0$ .
      - G.  $(b_k - d)\Lambda(h, b_k) < 0$ .
      - H.  $\Lambda(h, b_k) < 0$ .
    - ii. Otherwise, if  $\Lambda(f, a_k) \leq -B$ , do the following:

A. Using steps analogous to (ji), verify that  $\Lambda(h, a_k) < 0$ .

B. Using steps analogous to (ji), verify that  $\Lambda(h, b_k) > 0$ .

- (m) Execute **procedure II:69** on  $h$  and  $a_{\deg(h)} < b_{\deg(h)} \leq a_{\deg(h)-1} < b_{\deg(h)-1} \leq \dots \leq a_1 < b_1 \leq a_0 < b_0$ .
2. Otherwise, do the following:
  - (a) Verify that  $\deg(f) = 0$ .
  - (b) Therefore verify that  $f = f_0 \neq 0$ .
  - (c) Therefore verify that  $\text{sgn}(f_0) = \text{sgn}(\Lambda(f, a_0)) = -\text{sgn}(\Lambda(f, b_0)) = -\text{sgn}(f_0)$ .
  - (d) **Therefore verify that  $1 = -1$ .**
  - (e) **Abort procedure.**

### Procedure II:70(2.19)

#### Objective

Choose two lists of polynomials  $s, q$  in such a way that:

1.  $|s| > 1$ .
2. For  $i$  in  $[0 : |s|]$ ,  $\deg(s_i) = i$ .
3. For  $i$  in  $[0 : |s|]$ ,  $\text{sgn}((s_i)_i) = \text{sgn}((s_m)_m)$ .
4. For  $i$  in  $[1 : |s| - 1]$ ,  $s_{i-1} + s_{i+1} = q_i s_i$ .

The objective of the following instructions is to construct lists of polynomials  $g, h$  such that  $g_i s_{i+1} + h_i s_i = 1$  for  $i$  in  $[0 : |s| - 1]$ .

#### Implementation

1. Let  $m = |s| - 1$ .
2. Let  $g = h = \langle \rangle$ .
3. If  $m > 1$ , do the following:
  - (a) Verify that  $q_{m-1}s_{m-1} - s_m = s_{m-2}$ .
  - (b) Execute **procedure II:70** on  $s_{[0:m]}$  and  $q_{[1:m-1]}$  and let the tuple  $\langle, , g, h \rangle$  receive.
  - (c) Verify that  $g_{m-2}s_{m-1} + h_{m-2}s_{m-2} = 1$ .
  - (d) Let  $g_{m-1} = -h_{m-2}$ .
  - (e) Let  $h_{m-1} = g_{m-2} + h_{m-2}q_{m-1}$ .

- (f) Therefore verify that  $g_{m-1}s_m + h_{m-1}s_{m-1}$
- i.  $= g_{m-2}s_{m-1} + h_{m-2}(q_{m-1}s_{m-1} - s_m)$
  - ii.  $= g_{m-2}s_{m-1} + h_{m-2}s_{m-2}$
  - iii.  $= 1$ .
4. Otherwise, if  $m = 1$  do the following:
- (a) Let  $g_0 = 0$ .
  - (b) Let  $h_0 = \frac{1}{s_0}$ .
  - (c) **Therefore verify that**  $g_0s_1 + h_0s_0 = 1$ .
5. **Yield the tuple**  $\langle s, q, g, h \rangle$ .

### Declaration II:32

The notation  $J_s(x)$ , where  $s$  is a list of polynomials and  $x$  is a rational number, will be used as a shorthand for the number of changes observed when the list  $\text{sgn}(\Lambda(s, x))$  is iterated through linearly.

### Procedure II:71(2.20)

#### Objective

Execute **procedure II:70** and let  $\langle s, q, g, h \rangle$  receive. Choose a rational number  $X$ . The objective of the following instructions is to construct a rational number  $l$  and a procedure  $u(c, d)$  to show that either  $0 < 0$  or  $|J_s(d) - J_s(c)| = [\text{sgn}(\Lambda(s_{|s|-1}, c)) \neq \text{sgn}(\Lambda(s_{|s|-1}, d))]$ , when rational numbers  $c, d$  such that  $|c| \leq X$ ,  $|d| \leq X$ ,  $|d - c| \leq l$ ,  $0 \notin \Lambda(s, c)$ , and  $0 \notin \Lambda(s, d)$  are chosen.

#### Implementation

1. Execute **procedure II:67** on  $\langle s, X \rangle$  and let  $\langle G, t \rangle$  receive the result.
2. Let  $B = \max_i^{[0:|s|]} G_i$ .
3. Let  $C = \max_i^{[0:|s|-1]} \max(|\Lambda(\|g_i\|, X)|, |\Lambda(\|h_i\|, X)|)$ .
4. Let  $D = \max_i^{[1:|s|-1]} \max(|\Lambda(\|g_i\|, X)|, 2)$ .
5. Let  $l = \frac{1}{BCD}$ .
6. Let  $u(c, d)$  be the following procedure:
  - (a) Let  $i = 0$ .

- (b) If  $i + 1 < |s|$ , do the following:
  - i. Verify that  $\text{sgn}(\Lambda(s_i, c)) = \text{sgn}(\Lambda(s_i, d))$ .
  - ii. Verify that  $J_{s_{[0:i+1]}}(c) = J_{s_{[0:i+1]}}(d)$ .
  - iii. If  $\text{sgn}(\Lambda(s_{i+1}, c)) = \text{sgn}(\Lambda(s_{i+1}, d))$ , do the following:
    - A. Verify that  $J_{s_{[0:i+2]}}(c) = J_{s_{[0:i+2]}}(d)$ .
    - B. Set  $i$  to  $i + 1$  and go to (b).
  - iv. Otherwise if  $\text{sgn}(\Lambda(s_{i+1}, c)) \neq \text{sgn}(\Lambda(s_{i+1}, d))$  or  $i + 2 \geq |s|$ , do the following:
    - A. Verify that  $\text{sgn}(\Lambda(s_{i+1}, c)) \neq \text{sgn}(\Lambda(s_{i+1}, d))$ .
    - B. Verify that  $|J_{s_{[0:i+2]}}(c) - J_{s_{[0:i+2]}}(d)| = 1$ .
    - C. Verify that  $i + 2 = |s|$ .
    - D. Go to (c).
  - v. Execute **subprocedure II:72:0** on  $i$ .
  - vi. If  $\text{sgn}(\Lambda(s_{i+2}, c)) \neq \text{sgn}(\Lambda(s_{i+2}, d))$ , do the following:
    - A. Execute procedure  $t_{i+2}$  on  $\langle c, d \rangle$ .
    - B. Hence verify that  $|\Lambda(s_{i+2}, c)| < |\Lambda(s_{i+2}, d) - \Lambda(s_{i+2}, c)| = |(d - c)G_{i+2}| \leq \frac{1}{BCD} \cdot B = \frac{1}{CD} = \frac{1}{C} \cdot \frac{1}{D} \leq \frac{1}{C}(1 - \frac{1}{D})$ .
    - C. Using (A) and (i), verify that  $\frac{1}{C}(1 - \frac{1}{D}) < |s_{i+2}(c)| < \frac{1}{C}(1 - \frac{1}{D})$ .
    - D. **Abort procedure.**
  - vii. Otherwise if  $\text{sgn}(\Lambda(s_i, c)) = \text{sgn}(\Lambda(s_{i+2}, c))$ , do the following:
    - A. Verify that  $2\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_i, c)| + |\Lambda(s_{i+2}, c)| = |\Lambda(s_i, c) + \Lambda(s_{i+2}, c)| = |q_{i+1}(c)\Lambda(s_{i+1}, c)| < D\frac{1}{CD}$ .
    - B. Verify that  $2(1 - \frac{1}{D}) < 1$ .
    - C. Using (B) and the construction of  $D$ , verify that  $2 \leq D < 2$ .
    - D. **Abort procedure.**
  - viii. Otherwise, do the following:
    - A. Verify that  $\text{sgn}(\Lambda(s_i, d)) = \text{sgn}(\Lambda(s_i, c)) \neq \text{sgn}(\Lambda(s_{i+2}, c)) = \text{sgn}(\Lambda(s_{i+2}, d))$ .
    - B. Therefore verify that  $1 = J_{s_{[0:i+3]}}(c) - J_{s_{[0:i+1]}}(c) = J_{s_{[0:i+3]}}(d) - J_{s_{[0:i+1]}}(d)$ .



C. Therefore verify that  $J_{s_{[0:i+1]}}(c) + 1 = J_{s_{[0:i+3]}}(c) = J_{s_{[0:i+3]}}(d) = J_{s_{[0:i+1]}}(d) + 1$ .

D. Set  $i$  to  $i + 2$  and go to (b).

(c) If  $\text{sgn}(\Lambda(s_{[i-1]}, c)) = \text{sgn}(\Lambda(s_{[i-1]}, d))$ , then do the following:

i. **Verify that**  $J_s(c) = J_s(d)$ .

(d) Otherwise do the following:

i. **Verify that**  $|J_s(d) - J_s(c)| = 1$ .

7. **Yield the tuple**  $\langle l, u \rangle$ .

### Subprocedure II:72:0

**Objective** Choose a non-negative integer  $i < m$  and natural numbers  $c, d$  such that  $\text{sgn}(\Lambda(s_{i+1}, c)) \neq \text{sgn}(\Lambda(s_{i+1}, d))$  and  $i + 2 \leq m$ . The objective of the following instructions is to show that  $|\Lambda(s_{i+1}, c)| < \frac{1}{CD}$ ,  $|\Lambda(s_{i+1}, d)| < \frac{1}{CD}$ ,  $\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_i, c)|$ ,  $\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_i, d)|$ ,  $\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_{i+2}, c)|$ , and  $\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_{i+2}, d)|$ .

### Implementation

1. Verify the following in order:

(a) Execute procedure  $t_{i+1}$  on  $\langle c, d \rangle$ .

(b)  $|\Lambda(s_{i+1}, c)| < |\Lambda(s_{i+1}, c) - \Lambda(s_{i+1}, d)| = |c - d| |G_{i+1}| \leq |c - d| B \leq \frac{1}{BCD} \cdot B = \frac{1}{CD}$

(c)  $|\Lambda(s_{i+1}, d)| < |\Lambda(s_{i+1}, c) - \Lambda(s_{i+1}, d)| \leq \frac{1}{CD}$

(d)  $1 = \Lambda(g_i, c)\Lambda(s_{i+1}, c) + \Lambda(h_i, c)\Lambda(s_i, c) = |\Lambda(g_i, c)\Lambda(s_{i+1}, c) + \Lambda(h_i, c)\Lambda(s_i, c)| \leq |\Lambda(g_i, c)| |\Lambda(s_{i+1}, c)| + |\Lambda(h_i, c)| |\Lambda(s_i, c)| < C(\frac{1}{CD} + |\Lambda(s_i, c)|)$

(e)  $\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_i, c)|$

(f)  $1 < C(\frac{1}{CD} + |\Lambda(s_i, d)|)$

(g)  $\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_i, d)|$

(h)  $1 = \Lambda(g_{i+1}, c)\Lambda(s_{i+2}, c) + \Lambda(h_{i+1}, c)\Lambda(s_{i+1}, c) = |\Lambda(g_{i+1}, c)\Lambda(s_{i+2}, c) + \Lambda(h_{i+1}, c)\Lambda(s_{i+1}, c)| \leq |\Lambda(g_{i+1}, c)| |\Lambda(s_{i+2}, c)| + |\Lambda(h_{i+1}, c)| |\Lambda(s_{i+1}, c)| < C(|\Lambda(s_{i+2}, c)| + \frac{1}{CD})$

(i)  $\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_{i+2}, c)|$

(j)  $1 < C(|\Lambda(s_{i+2}, d)| + \frac{1}{CD})$

(k)  $\frac{1}{C}(1 - \frac{1}{D}) < |\Lambda(s_{i+2}, d)|$

### Procedure II:72(2.21)

#### Objective

Choose a polynomial  $p \neq 0$ . Choose a rational number  $k > 1 + \max_i^{[0:\deg(p)]} |\frac{p_i}{p_{\deg(p)}}|$ . The objective of the following instructions is to show that  $\text{sgn}(\Lambda(p, k)) = \text{sgn}(p_{\deg(p)})$ .

#### Implementation

1. Let  $n = \deg(p)$ .

2. In reverse order verify the following:

(a)  $\text{sgn}(\Lambda(p, k)) = \text{sgn}(p_{\deg(p)})$

(b)  $\text{sgn}(p_n k^n + p_{n-1} k^{n-1} + \dots + p_0 k^0) = \text{sgn}(p_n)$

(c)  $\text{sgn}(k^n + \frac{p_{n-1}}{p_n} k^{n-1} + \dots + \frac{p_0}{p_n} k^0) = 1$

(d)  $k^n + \frac{p_{n-1}}{p_n} k^{n-1} + \dots + \frac{p_0}{p_n} k^0 > 0$

(e)  $k^n > -(\frac{p_{n-1}}{p_n} k^{n-1} + \dots + \frac{p_0}{p_n} k^0)$

(f)  $k^n > |\frac{p_{n-1}}{p_n} k^{n-1} + \dots + \frac{p_0}{p_n} k^0|$

(g)  $k^n > |\max_i^{[0:n]} |\frac{p_i}{p_n}| (k^{n-1} + \dots + k^0)|$

(h)  $k^n > \max_i^{[0:n]} |\frac{p_i}{p_n}| \frac{k^n - 1}{k - 1}$

(i)  $k^{n+1} - k^n > \max_i^{[0:n]} |\frac{p_i}{p_n}| (k^n - 1)$

(j)  $k^{n+1} - (1 + \max_i^{[0:n]} |\frac{p_i}{p_n}|) k^n + \max_i^{[0:n]} |\frac{p_i}{p_n}| > 0$

(k)  $k > 1 + \max_i^{[0:n]} |\frac{p_i}{p_n}|$

### Procedure II:73(2.22)

#### Objective

Choose a polynomial  $p \neq 0$ . Choose a rational number  $k < -(1 + \max_i^{[0:\deg(p)]} |\frac{p_i}{p_{\deg(p)}}|)$ . The objective of the following instructions is to show that  $\text{sgn}(\Lambda(p, k)) = (-1)^{\deg(p)} \text{sgn}(p_{\deg(p)})$ .

## Implementation

1. Let  $t = \deg(p)$ .
2. Let  $q = \langle (-1)^{t-i} p_i \text{ for } i \in [0 : t + 1] \rangle$ .
3. Verify that  $k < -(1 + \max_i^{[1:t+1]} |\frac{q_i}{q_{\deg(q)}}|)$ .
4. Therefore verify that  $-k > 1 + \max_i^{[0:t]} |\frac{q_i}{q_{\deg(q)}}|$ .
5. Execute **procedure II:72** on  $\langle q, -k \rangle$ .
6. Hence verify that  $(-1)^t \text{sgn}(\Lambda(p, k))$ 
  - (a)  $= \text{sgn}((-1)^t \Lambda(p, k))$
  - (b)  $= \text{sgn}((-1)^t \sum_i^{[0:t+1]} p_i k^i)$
  - (c)  $= \text{sgn}(\sum_i^{[0:t+1]} (-1)^i (-1)^{t-i} p_i k^i)$
  - (d)  $= \text{sgn}(\sum_i^{[0:t+1]} q_i (-k)^i)$
  - (e)  $= \text{sgn}(\Lambda(q, -k))$
  - (f)  $= \text{sgn}(q_t)$
  - (g)  $= \text{sgn}(p_t)$ .
7. **Therefore verify that**  $\text{sgn}(\Lambda(p, k)) = (-1)^t (-1)^t \text{sgn}(\Lambda(p, k)) = (-1)^t \text{sgn}(p_t)$ .

## Procedure II:74(2.23)

### Objective

Choose a list of polynomials,  $s$ , and rational numbers  $a, l, c$  such that  $a < c$  and  $l > 0$ . The objective of the following instructions is to either show that  $0 < 0$  or to construct a list of rational numbers,  $b$ , such that  $a = b_0 < b_1 < \dots < b_{|b|-1} = c$ ,  $b_i - b_{i-1} \leq l$  for  $i$  in  $[1 : |b|]$ , and  $0 \notin \Lambda(s, b_i)$  for  $i$  in  $[1 : |b| - 1]$ .

### Implementation

1. Let  $e = \langle \langle \rangle, \langle \rangle, \dots, \langle \rangle \rangle$ .
2. Let  $f = \sum_r^{[0:|s|]} \deg(s_r)$ .
3. Let  $b = \langle a \rangle$ .
4. Let  $d = b_1$ .
5. While  $d + l < c$ , do the following:
  - (a) Let  $m = l$ .

- (b) While  $0 \in \Lambda(s, d + m)$  and  $\sum |e| \leq f$ , do the following:
  - i. Let  $0 \leq i < |s|$  be an integer such that  $\Lambda(s_i, d + m) = 0$ .
  - ii. Append  $d + m$  onto  $e_i$ .
  - iii. Set  $m = \frac{m}{2}$ .
- (c) If  $\sum |e| > f$ , then do the following:
  - i. If  $|e_i| \leq \deg(s_i)$  for  $0 \leq i < |s|$ , then do the following:
    - A. Verify that  $\sum |e| \leq f$ .
    - B. Therefore using (c), verify that  $\sum |e| \leq f < \sum |e|$ .
    - C. **Abort procedure.**
  - ii. Otherwise, do the following:
    - A. Let  $0 \leq i < |s|$  be an integer such that  $|e_i| > \deg(s_i)$ .
    - B. Execute **procedure II:66** on  $s_i$  and a sorted  $e_i$ .
    - C. **Abort procedure.**
- (d) Otherwise, do the following:
  - i. **Verify that**  $0 \notin \Lambda(s, d + m)$ .
  - ii. Append  $d + m$  onto  $b$ .
  - iii. **Verify that**  $0 < b_{|b|-1} - b_{|b|-2} = m \leq l$ .
  - iv. Set  $d$  to  $d + m$ .
  - v. Using (5), verify that  $d < c$ .
6. Verify that  $d < c$ .
7. Verify that  $d + l \geq c$ .
8. **Therefore verify that**  $0 < c - d \leq l$ .
9. Append  $c$  onto  $b$ .
10. **Yield**  $\langle b \rangle$ .

## Procedure II:75(2.24)

### Objective

Execute **procedure II:70** and let  $\langle s, q, g, h \rangle$  receive. Let  $m = |s| - 1$ . The objective of the following instructions is to either show that  $0 < 0$  or to construct two lists of rational numbers  $c, d$  such

that  $c_0 < d_0 \leq c_1 < d_1 \leq \dots \leq c_{m-1} < d_{m-1}$  and  $0 \neq \text{sgn}(\Lambda(s_m, c_i)) = -\text{sgn}(\Lambda(s_m, d_i))$  for  $i$  in  $[0 : m]$ .

### Implementation

1. Let  $U = 1 + \max_i^{[0:s]} \left( 1 + \max_j^{[1:i+1]} \left| \frac{(s_i)_{i-j}}{(s_i)_i} \right| \right)$
2. Using **procedure II:72**, verify that  $J(U) = 0$ .
3. Using **procedure II:73**, verify that  $J(-U) = m$ .
4. Execute **procedure II:71** on the tuple  $\langle s, q, U \rangle$  and let  $\langle l, u \rangle$  receive.
5. Execute **procedure II:74** on  $s$  with endpoints  $-U, U$  and a step size of  $l$  and let  $\langle e \rangle$  receive the result.
6. Let  $c = d = \langle \rangle$ .
7. For  $i = 1$  to  $i = |e| - 1$ :
  - (a) Execute procedure  $u$  on the tuple  $\langle e_{i-1}, e_i \rangle$ .
  - (b) If  $J_m(e_{i-1}) \neq J_m(e_i)$ , then do the following:
    - i. Append  $e_{i-1}$  to  $c$ .
    - ii. Append  $e_i$  to  $d$ .
    - iii. Verify that  $0 \neq |J_s(d_{|d|-1}) - J_s(c_{|c|-1})| = [\text{sgn}(\Lambda(s_{|s|-1}, c_{|c|-1})) \neq \text{sgn}(\Lambda(s_{|s|-1}, d_{|d|-1}))]$ .
    - iv. Therefore verify that  $\text{sgn}(s_m(c_{|c|-1})) \neq \text{sgn}(s_m(d_{|d|-1}))$ .
    - v. Therefore verify that  $|J_m(d_{|d|-1}) - J_m(c_{|c|-1})| = 1$ .
    - vi. Also verify that  $0 \notin \Lambda(s, c_{|c|-1})$ .
    - vii. Hence verify that  $\Lambda(s_m, c_{|c|-1}) \neq 0$ .
    - viii. Also verify that  $0 \notin \Lambda(s, d_{|d|-1})$ .
    - ix. Hence verify that  $\Lambda(s_m, d_{|d|-1}) \neq 0$ .
    - x. **Therefore verify that**  $0 \neq \text{sgn}(s_m(c_{|c|-1})) = -\text{sgn}(s_m(d_{|d|-1}))$ .
    - xi. **Also verify that**  $d_{|d|-2} \leq c_{|c|-1} < d_{|d|-1}$ .
8. If  $|c| = |d| < m$ , then do the following:
  - (a) Verify that each change of  $J_m(x)$  over the course of (7) was by 1.

- (b) Verify that  $J_m(x)$  changed less than  $m$  times over the course of (12).
  - (c) Therefore verify that  $|J_m(U) - J_m(-U)| < m$ .
  - (d) Therefore using (2) and (3), verify that  $m = |J_m(U) - J_m(-U)| < m$ .
  - (e) **Abort procedure.**
9. Otherwise, do the following:
- (a) **Verify that**  $m \leq |c| = |d|$ .
  - (b) **Yield the tuple**  $\langle c, d \rangle$ .

### Procedure II:76(2.26)

#### Objective

Choose two lists of polynomials  $s, q$  and a non-negative integer  $k$  in such a way that, letting  $m = |s| - 1$ ,

1.  $k < m$ .
2. For  $k \leq i \leq m$ ,  $\deg(s_i) = i$ .
3. For  $k < i < m$ ,  $s_{i-1} + s_{i+1} = q_i s_i$ .

Let  $\deg(0) = -1$ . The objective of the following instructions is to construct polynomials  $g, h$  such that  $s_k = g s_{m-1} + h s_m$ ,  $\deg(g) = m - 1 - k$ , and  $\deg(h) = m - 2 - k$ .

#### Implementation

1. If  $k < m - 2$ , do the following:
  - (a) Verify that  $s_k + s_{k+2} = q_{k+1} s_{k+1}$ .
  - (b) Therefore verify that  $s_k = q_{k+1} s_{k+1} - s_{k+2}$ .
  - (c) Execute **procedure II:76** on  $s, q, k + 1$  and let the tuple  $\langle g_1, h_1 \rangle$  receive.
  - (d) Verify that  $s_{k+1} = g_1 s_{m-1} + h_1 s_m$ .
  - (e) Verify that  $\deg(g_1) = m - 1 - (k + 1) = m - k - 2$ .
  - (f) Verify that  $\deg(h_1) = m - 2 - (k + 1) = m - k - 3$ .
  - (g) Execute **procedure II:76** on  $s, q, k + 2$  and let the tuple  $\langle g_2, h_2 \rangle$  receive.
  - (h) Verify that  $s_{k+2} = g_2 s_{m-1} + h_2 s_m$ .

- (i) Verify that  $\deg(g_2) = m - 1 - (k + 2) = m - k - 3$ .
  - (j) Verify that  $\deg(h_2) = m - 2 - (k + 2) = m - k - 4$ .
  - (k) Let  $g = q_{k+1}g_1 - g_2$ .
  - (l) **Verify that**  $\deg(g) = \max(1 + (m - k - 2), m - k - 3) = m - 1 - k$ .
  - (m) Let  $h = q_{k+1}h_1 - h_2$ .
  - (n) **Verify that**  $\deg(h) = \max(1 + (m - k - 3), m - k - 4) = m - 2 - k$ .
  - (o) **Verify that**  $s_k = q_{k+1}(g_1s_{m-1} + h_1s_m) - (g_2s_{m-1} + h_2s_m) = (q_{k+1}g_1 - g_2)s_{m-1} + (q_{k+1}h_1 - h_2)s_m = gs_{m-1} + hs_m$ .
2. Otherwise, if  $k = m - 2$  do the following:
- (a) Verify that  $s_{m-2} + s_m = q_{m-1}s_{m-1}$ .
  - (b) Let  $g = q_{m-1}$ .
  - (c) **Verify that**  $\deg(g) = 1 = m - 1 - k$ .
  - (d) Let  $h = -1$ .
  - (e) **Verify that**  $\deg(h) = 0 = m - 2 - k$ .
  - (f) **Therefore verify that**  $s_k = s_{m-2} = q_{m-1}s_{m-1} - s_m = gs_{m-1} + hs_m$ .
3. Otherwise, if  $k = m - 1$  do the following:
- (a) Let  $g = 1$ .
  - (b) **Verify that**  $\deg(g) = 0 = m - 1 - k$ .
  - (c) Let  $h = 0$ .
  - (d) **Verify that**  $\deg(h) = -1 = m - 2 - k$ .
  - (e) **Verify that**  $s_k = s_{m-1} = gs_{m-1} + hs_m$ .
4. **Yield the tuple**  $\langle g, h \rangle$ .

## Part III

# Complex Arithmetic

## Chapter 8

# Complex Arithmetic

### Declaration III:0

The phrase "complex number" will be used as a shorthand for an ordered pair of rational numbers.

### Declaration III:1

The phrase "the real part of  $a$ " and the notation  $\text{re}(a)$ , where  $a$  is a complex number, will be used as a shorthand for the first entry of  $a$ .

### Declaration III:2

The phrase "the imaginary part of  $a$ " and the notation  $\text{im}(a)$ , where  $a$  is a complex number, will be used as a shorthand for the second entry of  $a$ .

### Declaration III:3

The phrase " $a = b$ ", where  $a, b$  are complex numbers, will be used as a shorthand for " $\text{re}(a) = \text{re}(b)$  and  $\text{im}(a) = \text{im}(b)$ ".

### Procedure III:0(3.68)

#### Objective

Choose a complex number  $a$ . The objective of the following instructions is to show that  $a = a$ .

### Implementation

1. Show that  $\text{re}(a) = \text{re}(a)$ .
2. Show that  $\text{im}(a) = \text{im}(a)$ .
3. **Hence show that  $a = a$ .**

### Procedure III:1(3.69)

#### Objective

Choose two complex numbers  $a, b$  such that  $a = b$ . The objective of the following instructions is to show that  $b = a$ .

### Implementation

1. Show that  $\text{re}(b) = \text{re}(a)$  given that  $\text{re}(a) = \text{re}(b)$ .
2. Show that  $\text{im}(b) = \text{im}(a)$  given that  $\text{im}(a) = \text{im}(b)$ .
3. **Hence show that  $b = a$ .**

### Procedure III:2(3.70)

#### Objective

Choose three complex numbers  $a, b, c$  such that  $a = b$  and  $b = c$ . The objective of the following instructions is to show that  $a = c$ .

### Implementation

1. Show that  $\text{re}(a) = \text{re}(c)$ 
  - (a) given that  $\text{re}(a) = \text{re}(b)$
  - (b) and  $\text{re}(b) = \text{re}(c)$ .
2. Show that  $\text{im}(a) = \text{im}(c)$ 
  - (a) given that  $\text{im}(a) = \text{im}(b)$
  - (b) and  $\text{im}(b) = \text{im}(c)$ .
3. **Hence verify that  $a = c$ .**

### Declaration III:4

The notation  $a + b$ , where  $a, b$  are complex numbers, will be used as a shorthand for the pair  $\langle \text{re}(a) + \text{re}(b), \text{im}(a) + \text{im}(b) \rangle$ .

### Procedure III:3(3.71)

#### Objective

Choose two complex numbers  $a, b, c, d$  such that  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $a + b = c + d$ .

### Implementation

1. Using **declaration III:3**, show that
  - (a)  $\text{re}(a) = \text{re}(c)$
  - (b)  $\text{im}(a) = \text{im}(c)$
  - (c)  $\text{re}(b) = \text{re}(d)$
  - (d)  $\text{im}(b) = \text{im}(d)$ .
2. Hence show that  $a + b$ 
  - (a)  $= \langle \text{re}(a), \text{im}(a) \rangle + \langle \text{re}(b), \text{im}(b) \rangle$
  - (b)  $= \langle \text{re}(a) + \text{re}(b), \text{im}(a) + \text{im}(b) \rangle$
  - (c)  $= \langle \text{re}(c) + \text{re}(d), \text{im}(c) + \text{im}(d) \rangle$
  - (d)  $= \langle \text{re}(c), \text{im}(c) \rangle + \langle \text{re}(d), \text{im}(d) \rangle$
  - (e)  $= c + d$ .

### Procedure III:4(3.72)

#### Objective

Choose three complex numbers  $a, b, c$ . The objective of the following instructions is to show that  $(a + b) + c = a + (b + c)$ .

### Implementation

1. Show that  $(a + b) + c$ 
  - (a)  $= \langle \text{re}(a) + \text{re}(b), \text{im}(a) + \text{im}(b) \rangle + \langle \text{re}(c), \text{im}(c) \rangle$
  - (b)  $= \langle (\text{re}(a) + \text{re}(b)) + \text{re}(c), (\text{im}(a) + \text{im}(b)) + \text{im}(c) \rangle$
  - (c)  $= \langle \text{re}(a) + (\text{re}(b) + \text{re}(c)), \text{im}(a) + (\text{im}(b) + \text{im}(c)) \rangle$
  - (d)  $= \langle \text{re}(a), \text{im}(a) \rangle + \langle \text{re}(b) + \text{re}(c), \text{im}(b) + \text{im}(c) \rangle$
  - (e)  $= a + (b + c)$ .

### Procedure III:5(3.73)

#### Objective

Choose two complex numbers  $a, b$ . The objective of the following instructions is to show that  $a + b = b + a$ .

### Implementation

1. Show that  $a + b$ 
  - (a)  $= \langle \text{re}(a) + \text{re}(b), \text{im}(a) + \text{im}(b) \rangle$
  - (b)  $= \langle \text{re}(b) + \text{re}(a), \text{im}(b) + \text{im}(a) \rangle$
  - (c)  $= b + a$ .

### Declaration III:5

The notation  $a$ , where  $a$  is a rational number, will contextually be used as a shorthand for the pair  $\langle a, 0 \rangle$ .

### Procedure III:6(3.74)

#### Objective

Choose a complex number  $a$ . The objective of the following instructions is to show that  $0 + a = a$ .

#### Implementation

1. Show that  $0 + a$ 
  - (a)  $= \langle 0, 0 \rangle + \langle \text{re}(a), \text{im}(a) \rangle$
  - (b)  $= \langle 0 + \text{re}(a), 0 + \text{im}(a) \rangle$
  - (c)  $= \langle \text{re}(a), \text{im}(a) \rangle$
  - (d)  $= a$ .

#### Declaration III:6

The notation  $-a$ , where  $a$  is a complex number, will be used as a shorthand for the pair  $\langle -\text{re}(a), -\text{im}(a) \rangle$ .

### Procedure III:7(3.75)

#### Objective

Choose a complex number  $a$ . The objective of the following instructions is to show that  $-a + a = 0$ .

#### Implementation

1. Show that  $-a + a$ 
  - (a)  $= (-a) + a$
  - (b)  $= \langle -\text{re}(a), -\text{im}(a) \rangle + \langle \text{re}(a), \text{im}(a) \rangle$
  - (c)  $= \langle -\text{re}(a) + \text{re}(a), -\text{im}(a) + \text{im}(a) \rangle$
  - (d)  $= \langle 0, 0 \rangle$
  - (e)  $= 0$ .

#### Declaration III:7

The notation  $ab$ , where  $a, b$  are complex numbers, will be used as a shorthand for the pair  $\langle \text{re}(a)\text{re}(b) - \text{im}(a)\text{im}(b), \text{re}(a)\text{im}(b) + \text{im}(a)\text{re}(b) \rangle$ .

### Procedure III:8(3.76)

#### Objective

Choose four complex numbers  $a, b, c, d$  such that  $a = c$  and  $b = d$ . The objective of the following instructions is to show that  $ab = cd$ .

#### Implementation

1. Using **declaration III:3**, show that
  - (a)  $\text{re}(a) = \text{re}(c)$
  - (b)  $\text{im}(a) = \text{im}(c)$
  - (c)  $\text{re}(b) = \text{re}(d)$
  - (d)  $\text{im}(b) = \text{im}(d)$ .
2. Hence show that  $ab$ 
  - (a)  $= \langle \text{re}(a), \text{im}(a) \rangle \langle \text{re}(b), \text{im}(b) \rangle$
  - (b)  $= \langle \text{re}(a)\text{re}(b) - \text{im}(a)\text{im}(b), \text{re}(a)\text{im}(b) + \text{im}(a)\text{re}(b) \rangle$
  - (c)  $= \langle \text{re}(c)\text{re}(d) - \text{im}(c)\text{im}(d), \text{re}(c)\text{im}(d) + \text{im}(c)\text{re}(d) \rangle$
  - (d)  $= \langle \text{re}(c), \text{im}(c) \rangle \langle \text{re}(d), \text{im}(d) \rangle$
  - (e)  $= cd$ .

### Procedure III:9(3.77)

#### Objective

Choose three complex numbers  $a, b, c$ . The objective of the following instructions is to show that  $(ab)c = a(bc)$ .

#### Implementation

1. Show that  $(ab)c$ 
  - (a)  $= \langle \text{re}(a)\text{re}(b) - \text{im}(a)\text{im}(b), \text{re}(a)\text{im}(b) + \text{im}(a)\text{re}(b) \rangle \langle \text{re}(c), \text{im}(c) \rangle$
  - (b)  $= \langle (\text{re}(a)\text{re}(b) - \text{im}(a)\text{im}(b))\text{re}(c) - (\text{re}(a)\text{im}(b) + \text{im}(a)\text{re}(b))\text{im}(c), (\text{re}(a)\text{re}(b) - \text{im}(a)\text{im}(b))\text{im}(c) + (\text{re}(a)\text{im}(b) + \text{im}(a)\text{re}(b))\text{re}(c) \rangle$



$$(c) = \langle \text{re}(a)(\text{re}(b)\text{re}(c) - \text{im}(b)\text{im}(c)) - \text{im}(a)(\text{re}(b)\text{im}(c) + \text{im}(b)\text{re}(c)), \text{re}(a)(\text{re}(b)\text{im}(c) + \text{im}(b)\text{re}(c)) + \text{im}(a)(\text{re}(b)\text{re}(c) - \text{im}(b)\text{im}(c)) \rangle$$

$$(d) = \langle \text{re}(a), \text{im}(a) \rangle \langle \text{re}(b)\text{re}(c) - \text{im}(b)\text{im}(c), \text{re}(b)\text{im}(c) + \text{im}(b)\text{re}(c) \rangle$$

$$(e) = a(bc).$$

### Procedure III:10(3.78)

#### Objective

Choose two complex numbers  $a, b$ . The objective of the following instructions is to show that  $ab = ba$ .

#### Implementation

1. Show that  $ab$

$$(a) = \langle \text{re}(a)\text{re}(b) - \text{im}(a)\text{im}(b), \text{re}(a)\text{im}(b) + \text{im}(a)\text{re}(b) \rangle$$

$$(b) = \langle \text{re}(b)\text{re}(a) - \text{im}(b)\text{im}(a), \text{re}(b)\text{im}(a) + \text{im}(b)\text{re}(a) \rangle$$

$$(c) = ba.$$

### Procedure III:11(3.79)

#### Objective

Choose a complex number  $a$ . The objective of the following instructions is to show that  $1a = a$ .

#### Implementation

1. Show that  $1a$

$$(a) = \langle 1, 0 \rangle \langle \text{re}(a), \text{im}(a) \rangle$$

$$(b) = \langle 1\text{re}(a) - 0\text{im}(a), 1\text{im}(a) + 0\text{re}(a) \rangle$$

$$(c) = \langle \text{re}(a), \text{im}(a) \rangle$$

$$(d) = a.$$

### Procedure III:12(sun2107190636)

#### Objective

Choose a non-negative integer  $a$  and a complex number  $x$ . The objective of the following instructions is to show that  $(1+x)^a = \sum_r^{[0:a+1]} \binom{a}{r} x^r$ .

#### Implementation

Instructions are analogous to those of [procedure I:87](#).

### Declaration III:8

The notation  $\overline{a}$ , where  $a$  is a complex number, will be used as a shorthand for  $\langle \text{re}(a), -\text{im}(a) \rangle$ .

### Procedure III:13(3.00)

#### Objective

Choose two complex numbers  $a, b$ . The objective of the following instructions is to show that  $\overline{a+b} = \overline{a} + \overline{b}$ .

#### Implementation

1. Show that  $\overline{a+b}$

$$(a) = \langle \text{re}(a+b), -\text{im}(a+b) \rangle$$

$$(b) = \langle \text{re}(a) + \text{re}(b), -\text{im}(a) - \text{im}(b) \rangle$$

$$(c) = \overline{a} + \overline{b}.$$

### Procedure III:14(3.01)

#### Objective

Choose two complex numbers  $a, b$ . The objective of the following instructions is to show that  $\overline{ab} = \overline{a}\overline{b}$ .

#### Implementation

1. Show that  $\overline{ab}$

$$(a) = \langle \text{re}(ab), -\text{im}(ab) \rangle$$

$$\begin{aligned}
(b) &= \langle \text{re}(a) \text{re}(b) - \text{im}(a) \text{im}(b), -\text{re}(a) \text{im}(b) - \text{im}(a) \text{re}(b) \rangle \\
(c) &= \langle \text{re}(a), -\text{im}(a) \rangle \langle \text{re}(b), -\text{im}(b) \rangle \\
(d) &= \bar{a}\bar{b}.
\end{aligned}$$

### Declaration III:9

The notation  $\|a\|^2$ , where  $a$  is a complex number, will be used as a shorthand for  $\text{re}(a)^2 + \text{im}(a)^2$ .

### Procedure III:15(3.02)

#### Objective

Choose a complex number  $a$ . The objective of the following instructions is to show that  $a\bar{a} = \|a\|^2$ .

#### Implementation

1. Show that  $a\bar{a} = \|a\|^2$ .

### Procedure III:16(3.04)

#### Objective

Choose a list of complex numbers  $a$ . The objective of the following instructions is to show that  $\|\sum_r^{[0:|a|]} a_r\|^2 \leq |a| \sum_r^{[0:|a|]} \|a_r\|^2$ .

#### Implementation

1. Show that  $\|\sum_r^{[0:|a|]} a_r\|^2$ 
  - (a)  $= \sum_r^{[0:|a|]} \sum_k^{[0:|a|]} a_r \bar{a}_k$
  - (b)  $= \sum_r^{[0:|a|]} \|a_r\|^2 + 2 \sum_r^{[0:|a|]} \sum_k^{[r+1:|a|]} (\text{re}(a_r) \text{re}(a_k) + \text{im}(a_r) \text{im}(a_k))$
  - (c)  $= \sum_r^{[0:|a|]} \|a_r\|^2 + 2 \sum_r^{[0:|a|]} \sum_k^{[r+1:|a|]} (\text{re}(a_r)^2 - \text{re}(a_r) \text{re}(a_k) + \text{re}(a_k)^2 + \text{im}(a_r)^2 - \text{im}(a_r) \text{im}(a_k) + \text{im}(a_k)^2)$
  - (d)  $\leq \sum_r^{[0:|a|]} \|a_r\|^2 + \sum_r^{[0:|a|]} \sum_k^{[r+1:|a|]} (\text{re}(a_r)^2 + \text{re}(a_k)^2 + \text{im}(a_r)^2 + \text{im}(a_k)^2)$
  - (e)  $= \sum_r^{[0:|a|]} \|a_r\|^2 + \sum_r^{[0:|a|]} \sum_k^{[r+1:|a|]} (\|a_r\|^2 + \|a_k\|^2)$

$$\begin{aligned}
(f) &= \sum_r^{[0:|a|]} \|a_r\|^2 + \frac{1}{2} \sum_r^{[0:|a|]} \sum_k^{[0:r] \cap [r+1:|a|]} (\|a_r\|^2 + \|a_k\|^2) \\
(g) &= \sum_r^{[0:|a|]} \|a_r\|^2 + \frac{1}{2} (\sum_r^{[0:|a|]} (|a| - 1) \|a_r\|^2 + \sum_k^{[0:|a|]} (|a| - 1) \|a_k\|^2) \\
(h) &= \sum_r^{[0:|a|]} \|a_r\|^2 + \sum_r^{[0:|a|]} (|a| - 1) \|a_r\|^2 \\
(i) &= |a| \sum_r^{[0:|a|]} \|a_r\|^2
\end{aligned}$$

### Procedure III:17(3.05)

#### Objective

Choose a list of complex numbers  $a$ . The objective of the following instructions is to show that  $\frac{\|a_0\|^2}{|a|} - \sum_r^{[1:|a|]} \|a_r\|^2 \leq \|a_0 - \sum_r^{[1:|a|]} a_r\|^2$ .

#### Implementation

1. Using [procedure III:16](#), show that  $\|a_0\|^2$ 
  - (a)  $= \|\sum_r^{[1:|a|]} a_r + (a_0 - \sum_r^{[1:|a|]} a_r)\|^2$
  - (b)  $\leq |a| \sum_r^{[1:|a|]} \|a_r\|^2 + |a| \|a_0 - \sum_r^{[1:|a|]} a_r\|^2$
2. Therefore show that  $\frac{\|a_0\|^2}{|a|} - \sum_r^{[1:|a|]} \|a_r\|^2 \leq \|a_0 - \sum_r^{[1:|a|]} a_r\|^2$ .

### Procedure III:18(3.04aa)

#### Objective

Choose a list of complex numbers  $a$  and a list of rational numbers  $b$  such that  $|a| = |b|$  and  $\|a_i\|^2 \leq b_i^2$  for each  $i \in [0 : |a|]$ . The objective of the following instructions is to show that  $\|\sum_r^{[0:|a|]} a_r\|^2 \leq (\sum_r^{[0:|b|]} b_r)^2$ .

#### Implementation

1. If  $|a| = 0$ , then do the following:
  - (a) Show that  $\|\sum_i^{[0:|a|]} a_i\|^2 = \|0\|^2 = (\sum_i^{[0:|b|]} b_i)^2$ .
2. Otherwise do the following:
  - (a) Show that  $|a| > 0$ .

(b) Show that  $\|\sum_i^{[1:|a|]} a_i\|^2 \leq (\sum_i^{[1:|b|]} b_i)^2$  using **procedure III:18** on  $a_{[1:|a|]}$  and  $b_{[1:|b|]}$ .

(c) Show that  $\operatorname{re}(\overline{a_0} \sum_i^{[1:|a|]} a_i)^2$

$$\text{i.} \leq \|\overline{a_0} \sum_i^{[1:|a|]} a_i\|^2$$

$$\text{ii.} = \|\overline{a_0}\|^2 \|\sum_i^{[1:|a|]} a_i\|^2$$

$$\text{iii.} \leq b_0^2 (\sum_i^{[1:|a|]} b_i)^2.$$

(d) Hence show that  $\|\sum_i^{[0:|a|]} a_i\|^2$

$$\text{i.} = (a_0 + \sum_i^{[1:|a|]} a_i) \overline{(a_0 + \sum_i^{[1:|a|]} a_i)}$$

$$\text{ii.} = \|a_0\|^2 + a_0 \overline{\sum_i^{[1:|a|]} a_i} + \overline{a_0} \sum_i^{[1:|a|]} a_i + \|\sum_i^{[1:|a|]} a_i\|^2$$

$$\text{iii.} \leq b_0^2 + \overline{a_0} \sum_i^{[1:|a|]} a_i + \overline{a_0} \sum_i^{[1:|a|]} a_i + (\sum_i^{[1:|a|]} b_i)^2$$

$$\text{iv.} = b_0^2 + 2 \operatorname{re}(\overline{a_0} \sum_i^{[1:|a|]} a_i) + (\sum_i^{[1:|a|]} b_i)^2$$

$$\text{v.} \leq b_0^2 + 2b_0 \sum_i^{[1:|a|]} b_i + (\sum_i^{[1:|a|]} b_i)^2$$

$$\text{vi.} = (b_0 + \sum_i^{[1:|a|]} b_i)^2$$

$$\text{vii.} = (\sum_i^{[0:|a|]} b_i)^2.$$

### Procedure III:19(sat1708191238)

#### Objective

Choose two complex numbers  $a, d$  and two rational numbers  $b, c$  such that  $\|a\|^2 \leq b^2 < c^2 \leq \|d\|^2$ . The objective of the following instructions is to show that  $\|d - a\|^2 \geq (c - b)^2$ .

#### Implementation

1. Show that  $\operatorname{re}(\frac{a}{d})^2$

$$\text{(a)} = \operatorname{re}(\frac{a\overline{d}}{\|d\|^2})^2 = \frac{\operatorname{re}(a\overline{d})^2}{\|d\|^4} \leq \frac{\|a\overline{d}\|^2}{\|d\|^4}$$

$$\text{(b)} = \frac{\|a\|^2 \|d\|^2}{\|d\|^4} = \frac{\|a\|^2}{\|d\|^2} \leq \frac{b^2}{c^2} = (\frac{b}{c})^2.$$

2. Hence show that  $\operatorname{re}(\frac{a}{d}) \leq \frac{b}{c} < 1$ .

3. Hence show that  $\|d - a\|^2$

$$\text{(a)} = \|\frac{d-a}{d}\|^2 \|d\|^2$$

$$\text{(b)} = (\operatorname{re}(1 - \frac{a}{d})^2 + \operatorname{im}(1 - \frac{a}{d})^2) \|d\|^2$$

$$\text{(c)} \geq \operatorname{re}(1 - \frac{a}{d})^2 \|d\|^2$$

$$\text{(d)} = (1 - \operatorname{re}(\frac{a}{d}))^2 \|d\|^2$$

$$\text{(e)} \geq (1 - \frac{b}{c})^2 c^2$$

$$\text{(f)} = (c - b)^2.$$

### Declaration III:10

The notation  $\frac{1}{a}$ , where  $a$  is a complex number, will be used as a shorthand for the pair  $\frac{1}{\|a\|^2} \overline{a}$ .

### Procedure III:20(3.81)

#### Objective

Choose a complex number  $a$  such that  $a \neq 0$ . The objective of the following instructions is to show that  $\frac{1}{a}a = 1$ .

#### Implementation

1. Show that  $\operatorname{re}(a) \neq \operatorname{re}(0) = 0$  or  $\operatorname{im}(a) \neq \operatorname{im}(0) = 0$  using **declaration III:3**.

2. Hence show that  $\|a\|^2 = \operatorname{re}(a)^2 + \operatorname{im}(a)^2 > 0$ .

3. Hence show that  $\frac{1}{a}a$

$$\text{(a)} = (\frac{1}{\|a\|^2} \overline{a})a$$

$$\text{(b)} = \frac{1}{\|a\|^2} (\overline{a}a)$$

$$\text{(c)} = \frac{1}{\|a\|^2} \|a\|^2$$

$$\text{(d)} = 1.$$

### Procedure III:21(3.82)

#### Objective

Choose three complex numbers  $a, b, c$ . The objective of the following instructions is to show that  $a(b + c) = ab + ac$ .

### Implementation

1.  $a(b + c)$ 
  - (a)  $= \langle \text{re}(a), \text{im}(a) \rangle \langle \text{re}(b) + \text{re}(c), \text{im}(b) + \text{im}(c) \rangle$
  - (b)  $= \langle \text{re}(a)(\text{re}(b) + \text{re}(c)) - \text{im}(a)(\text{im}(b) + \text{im}(c)), \text{re}(a)(\text{im}(b) + \text{im}(c)) + \text{im}(a)(\text{re}(b) + \text{re}(c)) \rangle$
  - (c)  $= \langle (\text{re}(a)\text{re}(b) - \text{im}(a)\text{im}(b)) + (\text{re}(a)\text{re}(c) - \text{im}(a)\text{im}(c)), (\text{re}(a)\text{im}(b) + \text{im}(a)\text{re}(b)) + (\text{re}(a)\text{im}(c) + \text{im}(a)\text{re}(c)) \rangle$
  - (d)  $= \langle \text{re}(a)\text{re}(b) - \text{im}(a)\text{im}(b), \text{re}(a)\text{im}(b) + \text{im}(a)\text{re}(b) \rangle + \langle \text{re}(a)\text{re}(c) - \text{im}(a)\text{im}(c), \text{re}(a)\text{im}(c) + \text{im}(a)\text{re}(c) \rangle$
  - (e)  $= ab + ac.$

### Declaration III:11

The notation  $i$  will be used as a shorthand for  $\langle 0, 1 \rangle$ .

### Procedure III:22(3.03)

#### Objective

Choose an integer  $a$ . The objective of the following instructions is to show that  $i^{4a} = 1$ ,  $i^{4a+1} = i$ ,  $i^{4a+2} = -1$ , and  $i^{4a+3} = -i$ .

#### Implementation

1. Show that  $i^2 = -1$ .
2. Hence show that  $i^4 = (-1)^2 = 1$ .
3. **Hence show that**
  - (a)  $i^{4a} = (i^4)^a = 1^a = 1$
  - (b)  $i^{4a+1} = i^{4a}i = 1i = i$
  - (c)  $i^{4a+2} = i^{4a+1}i = i^2 = -1$
  - (d)  $i^{4a+3} = i^{4a+2}i = (-1)i = -i$ .

### Declaration III:12

The notation  $a \equiv b$  (err  $c_1$ ) (err  $c_2$ )  $\cdots$  (err  $c_n$ ) will be used as a shorthand for  $\|b - a\|^2 \leq \|c_1\|^2 \leq \|c_2\|^2 \leq \cdots \leq \|c_n\|^2$ .

### Procedure III:23(mon1908191916)

#### Objective

Choose four complex numbers  $a, b, c, d$  in such a way that  $a \equiv b$  (err  $c$ ) and  $\|c\|^2 \leq \|d\|^2$ . The objective of the following instructions is to show that  $a \equiv b$  (err  $c$ ) (err  $d$ ).

#### Implementation

1. Show that  $\|b - a\| \leq \|c\|^2 \leq \|d\|^2$ .
2. **Hence show that  $a \equiv b$  (err  $c$ ) (err  $d$ ) using declaration III:12.**

### Procedure III:24(mon1908191825)

#### Objective

Choose three complex numbers  $a, b, c$  and two non-negative rational numbers  $d, e$  in such a way that  $a \equiv b$  (err  $d$ ) and  $b \equiv c$  (err  $e$ ). The objective of the following instructions is to show that  $a \equiv c$  (err  $d + e$ ).

#### Implementation

1. Show that  $\|b - a\|^2 \leq \|d\|^2 = d^2$ .
2. Show that  $\|c - b\|^2 \leq \|e\|^2 = e^2$ .
3. Hence show that  $\|c - a\|^2 = \|(c - b) + (b - a)\|^2 \leq (e + d)^2$  using **procedure III:18**.
4. **Hence show that  $a \equiv c$  (err  $e + d$ ) using declaration III:12.**

### Procedure III:25(mon1908191839)

#### Objective

Choose four complex numbers  $a, b, c, d$  and two non-negative rational numbers  $e, f$  in such a way that  $a \equiv b$  (err  $e$ ) and  $c \equiv d$  (err  $f$ ). The objective of the following instructions is to show that  $a + c \equiv b + d$  (err  $e + f$ ).

### Implementation

1. Show that  $\|b - a\|^2 \leq \|e\|^2 = e^2$ .
2. Show that  $\|d - c\|^2 \leq \|f\|^2 = f^2$ .
3. Hence show that  $\|(b + d) - (a + c)\|^2 = \|(b - a) + (d - c)\|^2 \leq (e + f)^2$  using **procedure III:18**.
4. **Hence show that**  $a + c \equiv b + d$  (err  $e + f$ ) using **declaration III:12**.

### Procedure III:26(mon1908191849)

#### Objective

Choose three complex numbers  $a, b, c$  in such a way that  $a \equiv b$  (err  $c$ ). The objective of the following instructions is to show that  $-a \equiv -b$  (err  $c$ ).

#### Implementation

1. Show that  $\|(-b) - (-a)\|^2 = \|b - a\|^2 \leq \|c\|^2$ .
2. **Hence show that**  $-a \equiv -b$  (err  $c$ ) using **declaration III:12**.

### Procedure III:27(mon1908191857)

#### Objective

Choose four complex numbers  $a, b, c, d$  in such a way that  $a \equiv b$  (err  $c$ ). The objective of the following instructions is to show that  $ad \equiv bd$  (err  $cd$ ).

#### Implementation

1. Show that  $\|b - a\|^2 \leq \|c\|^2$ .
2. Hence show that  $\|bd - ad\|^2 \leq \|cd\|^2$ .
3. **Hence show that**  $ad \equiv bd$  (err  $cd$ ) using **declaration III:12**.

### Procedure III:28(mon1908191905)

#### Objective

Choose two complex numbers  $a, b, c$  in such a way that  $a \neq 0$ ,  $b \neq 0$ , and  $a \equiv b$  (err  $c$ ). The ob-

jective of the following instructions is to show that  $\frac{1}{a} \equiv \frac{1}{b}$  (err  $\frac{c}{ab}$ ).

#### Implementation

1. Show that  $\|b - a\| \leq \|c\|^2$ .
2. Hence show that  $\|\frac{1}{b} - \frac{1}{a}\|^2 = \|\frac{a-b}{ba}\|^2 \leq \|\frac{c}{ba}\|^2$ .
3. **Hence show that**  $\frac{1}{a} \equiv \frac{1}{b}$  (err  $\frac{c}{ab}$ ) using **declaration III:12**.

## Chapter 9

# Exponential and Trigonometric Functions

### Declaration III:13

The notation  $\exp_n(a)$ , where  $a$  is a complex number, will be used as a shorthand for  $(1 + \frac{a}{n})^n$ .

### Procedure III:29(3.08)

#### Objective

Choose a rational number  $a$  and a positive integer  $n$  such that  $-n < a < 1$ . The objective of the following instructions is to show that  $\exp_n(a) \leq \frac{1}{1-a}$ .

#### Implementation

1. Using **procedure II:32**, show that  $\exp_n(a)$

$$\begin{aligned} \text{(a)} &= \left(\frac{n+a}{n}\right)^n \\ \text{(b)} &= \left(\frac{n}{n+a}\right)^{-n} \\ \text{(c)} &= \frac{1}{\left(1+\frac{-a}{n+a}\right)^n} \\ \text{(d)} &\leq \frac{1}{1+\frac{-an}{n+a}} \\ \text{(e)} &\leq \frac{1}{1-a}. \end{aligned}$$

### Procedure III:30(3.09)

#### Objective

Choose a rational number  $a$  and a positive integer  $n$  such that  $a > -n$ . The objective of the following instructions is to show that  $\frac{\exp_{n+1}(a)}{\exp_n(a)} \geq 1$ .

#### Implementation

1. Using **procedure II:32**, show that  $\frac{\exp_{n+1}(a)}{\exp_n(a)}$ 
  - (a)  $= \frac{(\frac{n+1+a}{n+1})^n}{(\frac{n+a}{n})^n} \left(1 + \frac{a}{n+1}\right)$
  - (b)  $= \left(\frac{(n+1+a)n}{(n+1)(n+a)}\right)^n \left(1 + \frac{a}{n+1}\right)$
  - (c)  $= \left(\frac{n^2+n+na}{n^2+an+n+a}\right)^n \left(1 + \frac{a}{n+1}\right)$
  - (d)  $= \left(1 - \frac{a}{(n+1)(n+a)}\right)^n \left(1 + \frac{a}{n+1}\right)$
  - (e)  $\geq \left(1 - \frac{an}{(n+1)(n+a)}\right) \left(1 + \frac{a}{n+1}\right)$
  - (f)  $= 1 + \frac{a(n+a)}{(n+1)(n+a)} - \frac{an}{(n+1)(n+a)} - \frac{a^2n}{(n+1)^2(n+a)}$
  - (g)  $= 1 + \frac{a^2}{(n+1)(n+a)} - \frac{a^2n}{(n+1)^2(n+a)}$
  - (h)  $= 1 + \frac{a^2}{(n+1)^2(n+a)}$
  - (i)  $\geq 1$

### Procedure III:31(3.10)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct positive rational numbers  $a, b$  such that  $a > 1$ , and a procedure,  $p(x, r, n)$ , to show that  $(1 + \frac{x}{n})^r \leq a^2$  when given a rational number  $x$  and a non-negative integers  $r, n$  such that  $r \leq n$ ,  $n \geq b$  and  $x^2 \leq X^2$  are chosen.

#### Implementation

1. Let  $a = 2^{\lceil X \rceil}$ .
2. Let  $b = X$ .
3. Let  $p(x, r, n)$  be the following procedure:
  - (a) Show that  $0 \leq 1 + \frac{x}{n} \leq 2$ 
    - i. given that  $-1 \leq \frac{x}{n} \leq 1$
    - ii. given that  $-X \leq x \leq X$
    - iii. given that  $x^2 \leq X^2$ .
  - (b) Hence using **procedure III:29** and **procedure III:30**, show that  $(1 + \frac{x}{n})^r$ 
    - i.  $\leq (1 + \frac{X}{n})^r$
    - ii.  $\leq \exp_n(X)$
    - iii.  $\leq (1 + \frac{X}{2^{\lceil X \rceil} n})^{2^{\lceil X \rceil} n}$
    - iv.  $= ((1 + \frac{X}{2^{\lceil X \rceil} n})^n)^{2^{\lceil X \rceil}}$
    - v.  $= \exp_n(\frac{X}{2^{\lceil X \rceil}})^{2^{\lceil X \rceil}}$
    - vi.  $\leq (\frac{1}{1 - \frac{X}{2^{\lceil X \rceil}}})^{2^{\lceil X \rceil}}$
    - vii.  $\leq 2^{2^{\lceil X \rceil}}$
    - viii.  $= a^2$ .
4. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:32(3.11)

#### Objective

Choose a rational number  $X \leq 0$ . The objective of the following instructions is to construct two rational numbers  $a > 0, b$ , and a procedure  $p(x, r, n)$  to

show that  $(1 + \frac{x}{n})^r \geq a^2$  when a rational number  $x$  and non-negative integers  $r, n$  such that  $X \leq x \leq 0$ ,  $r \leq n$ , and  $n > b$  are chosen.

#### Implementation

1. Execute **procedure III:31** on  $\langle -2X \rangle$  and let  $\langle c, d, q \rangle$  receive.
2. Let  $a = c^{-1}$ .
3. Let  $b = \max(-2X, d)$ .
4. Let  $p(x, r, n)$  be the following procedure:
  - (a) Show that  $0 \leq -2x \leq -2X$ 
    - i. given that  $2X \leq 2x \leq 0$
    - ii. given that  $X \leq x \leq 0$ .
  - (b) Show that  $(1 + \frac{-2x}{n})^r \leq c^2$  using procedure  $q$ .
  - (c) Show that  $\frac{n}{2} \leq n + x < n$ 
    - i. given that  $-\frac{n}{2} \leq x \leq 0$
    - ii. given that  $n > b \geq -2X \geq -2x \geq 0$ .
  - (d) Hence show that  $(1 + \frac{x}{n})^r$ 
    - i.  $= (\frac{n+x}{n})^r$
    - ii.  $= (\frac{n}{n+x})^{-r}$
    - iii.  $= (1 - \frac{x}{n+x})^{-r}$
    - iv.  $\geq (1 - \frac{x}{\frac{1}{2}n})^{-r}$
    - v.  $= (1 - \frac{2x}{n})^{-r}$
    - vi.  $= ((1 + \frac{-2x}{n})^r)^{-1}$
    - vii.  $\geq (c^2)^{-1}$
    - viii.  $= a^2$ .
5. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:33(3.12)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a > 0, b$ , and a procedure  $p(x, r, n)$  to show that  $(1 + \frac{x}{n})^r \geq a^2$  when a rational number  $x$

and non-negative integers  $r, n$  such that  $x^2 \leq X^2$ ,  $r \leq n$ , and  $n > b$  are chosen.

### Implementation

1. Execute **procedure III:32** on  $\langle -X \rangle$  and let  $\langle c, b, q \rangle$  receive.
2. Let  $a = \min(1, c)$ .
3. Let  $p(x, r, n)$  be the following procedure:
  - (a) If  $x < 0$ , then do the following:
    - i. Show that  $-X \leq x \leq 0$  given that  $x^2 \leq X^2$ .
    - ii. **Hence show that**  $(1 + \frac{x}{n})^r \geq c^2 \geq a^2$  **using procedure  $q$ .**
  - (b) Otherwise do the following:
    - i. Verify that  $x \geq 0$ .
    - ii. **Show that**  $(1 + \frac{x}{n})^r \geq 1 + \frac{rx}{n} \geq 1 \geq a^2$  **using procedure II:32.**
4. **Yield the tuple**  $\langle a, b, p \rangle$ .

### Procedure III:34(3.13)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct positive rational numbers  $a, b$  such that  $a > 1$ , and a procedure,  $p(x, r, n)$ , to show that  $\|(1 + \frac{x}{n})^r\|^2 \leq a^2$  when a complex number  $x$  and non-negative integers  $r, n$  such that  $\|x\|^2 \leq X^2$ ,  $r \leq n$ , and  $n > b$  are chosen.

### Implementation

1. Let  $c = 2X + X^2$ .
2. Execute **procedure III:31** on  $\langle c \rangle$  and let  $\langle a, b, q \rangle$  receive.
3. Let  $p(x, r, n)$  be the following procedure:
  - (a) Let  $y = 2|\operatorname{re}(x)| + \|x\|^2$ .
  - (b) Show that  $|y| = y \leq 2X + X^2 = c$ 
    - i. given that  $|\operatorname{re}(x)| \leq X$
    - ii. given that  $|\operatorname{re}(x)|^2 \leq \|x\|^2 \leq X^2$ .

(c) Hence show that  $(1 + \frac{y}{n})^r \leq a^2$  using procedure  $q$ .

(d) Now using **procedure III:15** show that  $\|(1 + \frac{x}{n})^r\|^2$

$$\text{i.} = (1 + \frac{x}{n})^r \overline{(1 + \frac{x}{n})^r}$$

$$\text{ii.} = (1 + \frac{x}{n})^r (1 + \frac{\bar{x}}{n})^r$$

$$\text{iii.} = (1 + \frac{2\operatorname{re}(x)}{n} + \frac{\|x\|^2}{n^2})^r$$

$$\text{iv.} \leq (1 + \frac{2|\operatorname{re}(x)|}{n} + \frac{\|x\|^2}{n^2})^r$$

$$\text{v.} \leq (1 + \frac{2|\operatorname{re}(x)| + \|x\|^2}{n})^r$$

$$\text{vi.} = (1 + \frac{y}{n})^r$$

$$\text{vii.} \leq a^2.$$

4. **Yield the tuple**  $\langle a, b, p \rangle$ .

### Procedure III:35(3.14)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, b$  and a procedure,  $p(x, r, n)$ , to show that  $\|(1 + \frac{x}{n})^r\|^2 \geq a^2$  when a rational number  $x$  and non-negative integers  $r, n$  such that  $\|x\|^2 \leq X^2$ ,  $r \leq n$  and  $n > b$  are chosen.

### Implementation

1. Let  $c = 2X + X^2$ .
2. Execute **procedure III:33** on  $\langle c \rangle$  and let  $\langle a, d, q \rangle$  receive.
3. Let  $b = \max(c, d)$ .
4. Let  $p(x, r, n)$  be the following procedure:
  - (a) Let  $y = 2|\operatorname{re}(x)| + \|x\|^2$ .
  - (b) Verify that  $|-y| = y \leq 2X + X^2 = c$ .
  - (c) Hence show that  $(1 + \frac{-y}{n})^r \geq a^2$  using procedure  $q$ .
  - (d) Also, show that  $n > b \geq c \geq y$ .
  - (e) Hence show that  $\|(1 + \frac{x}{n})^r\|^2$ 
    - i.  $= (1 + \frac{x}{n})^r \overline{(1 + \frac{x}{n})^r}$
    - ii.  $= (1 + \frac{x}{n})^r (1 + \frac{\bar{x}}{n})^r$



- iii.  $= (1 + \frac{2 \operatorname{re}(x)}{n} + \frac{\|x\|^2}{n^2})^r$
- iv.  $\geq (1 - \frac{2|\operatorname{re}(x)|}{n} - \frac{\|x\|^2}{n^2})^r$
- v.  $\geq (1 - \frac{2|\operatorname{re}(x)| + \|x\|^2}{n})^r$
- vi.  $= (1 + \frac{-y}{n})^r$
- vii.  $\geq a^2$ .

5. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:36(3.15)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct rational numbers  $a, b$  such that  $a > 0$ , and a procedure,  $p$ , to show that  $\exp_n(x + y) \equiv \exp_n(x) \exp_n(y)$  (err  $\frac{axy}{n}$ ) (err  $\frac{aX^2}{n}$ ) when two complex numbers  $x, y$  and a positive integer  $n > b$  such that  $\|x\|^2 \leq X^2$ ,  $\|y\|^2 \leq X^2$  are chosen.

#### Implementation

1. Execute **procedure III:34** on  $\langle 2X \rangle$  and let  $\langle c, q \rangle$  receive.
2. Let  $a = c^3$ .
3. Let  $p(x, y, n)$  be the following procedure:
  - (a) Using procedure  $q$ , show that  $\exp_n(x + y) \equiv \exp_n(x) \exp_n(y)$ 
    - i. (err  $\exp_n(x) \exp_n(y) - \exp_n(x + y)$ )
    - ii. (err  $(1 + \frac{x}{n})^n (1 + \frac{y}{n})^n - (1 + \frac{x+y}{n})^n$ )
    - iii. (err  $(1 + \frac{x+y}{n} + \frac{xy}{n^2})^n - (1 + \frac{x+y}{n})^n$ )
    - iv. (err  $\frac{xy}{n^2} \sum_r^{[0:n]} (1 + \frac{x+y}{n} + \frac{xy}{n^2})^r (1 + \frac{x+y}{n})^{n-1-r}$ )
    - v. (err  $\frac{xy}{n^2} \sum_r^{[0:n]} (1 + \frac{x}{n})^r (1 + \frac{y}{n})^r (1 + \frac{x+y}{n})^{n-1-r}$ )
    - vi. (err  $\frac{xy}{n^2} \sum_r^{[0:n]} c^3$ )
    - vii. (err  $\frac{axy}{n}$ )
    - viii. (err  $\frac{aX^2}{n}$ ).
4. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:37(thu2507191359)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct rational numbers  $a, b$  such that  $a > 0$  and a procedure  $p(x, y, n)$  to show that  $\exp_n(x - y) \equiv \frac{\exp_n(x)}{\exp_n(y)}$  (err  $\frac{a}{n}$ ) when two complex numbers  $x, y$  and a positive integer  $n$  such that  $\|x\|^2 \leq X$ ,  $\|y\|^2 \leq X$ , and  $n > b$  are chosen.

#### Implementation

1. Execute **procedure III:36** on  $\langle X \rangle$  and let  $\langle c, d \rangle$  receive.
2. Execute **procedure III:35** on  $\langle X \rangle$  and let  $\langle e, f \rangle$  receive.
3. Execute **procedure III:34** on  $\langle X \rangle$  and let  $\langle g, h \rangle$  receive.
4. Let  $b = \max(d, f, h)$ .
5. Let  $a = c(1 + \frac{g}{e})X^2$ .
6. Let  $p(x, y, n)$  be the following procedure:
  - (a) Using procedures  $q, r, t$ , show that  $\exp_n(x - y)$ 
    - i.  $\equiv \exp_n(x) \exp_n(-y)$  (err  $\frac{cX^2}{n}$ )
    - ii.  $= \exp_n(x) \frac{\exp_n(y) \exp_n(-y)}{\exp_n(y)}$
    - iii.  $\equiv \exp_n(x) \frac{\exp_n(0)}{\exp_n(y)}$  (err  $\frac{g}{e} \cdot \frac{cX^2}{n}$ )
    - iv.  $= \frac{\exp_n(x)}{\exp_n(y)}$ .
  - (b) **Hence show that**  $\exp_n(x - y) \equiv \frac{\exp_n(x)}{\exp_n(y)}$  (err  $\frac{cX^2}{n} + \frac{gcX^2}{en}$ ) (err  $\frac{a}{n}$ ).
7. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:38(3.16)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct positive rational numbers  $a, b$  and a procedure,  $p(x, k, n)$ , to show that  $\exp_n(kx) \equiv \exp_n(x)^k$  (err  $\frac{ak}{n}$ ) when a

complex number  $x$ , and non-negative integers  $k, n$  such that  $n > b$  and  $\|kx\|^2 \leq X^2$  are chosen.

### Implementation

1. Execute **procedure III:34** on  $\langle X \rangle$  and let  $\langle c, d, q \rangle$  receive.
2. Execute **procedure III:36** on  $\langle X \rangle$  and let  $\langle e, f, t \rangle$  receive.
3. Let  $a = ecX^2$
4. Let  $b = \max(d, f)$ .
5. Let  $p(x, k, n)$  be the following procedure:
  - (a) If  $k > 0$ , then for  $r \in [1 : k]$  do the following:
    - i. Show that  $\|xr\|^2 \leq \|kx\|^2 \leq X^2$ .
    - ii. Hence show that  $\|\exp_{nr}(xr)\|^2 \leq c^2$  using procedure  $q$ .
    - iii. Hence show that  $\|\exp_n(x)^r\|^2 = \|(1 + \frac{x}{n})^{nr}\|^2 = \|(1 + \frac{xr}{nr})^{nr}\|^2 = \|\exp_{nr}(xr)\|^2 \leq \frac{c^2}{c^2}$
  - (b) Hence using procedure  $t$ , show that  $\exp_n(kx)$ 
    - i.  $= \exp_n(x)^0 \exp_n(kx)$
    - ii.  $\equiv \exp_n(x)^1 \exp_n((k-1)x) \text{ (err } \frac{ceX^2}{n})$
    - iii.  $\equiv \exp_n(x)^2 \exp_n((k-2)x) \text{ (err } \frac{ceX^2}{n})$
    - iv.  $\vdots$
    - v.  $\equiv \exp_n(x)^k \exp_n((k-k)x) \text{ (err } \frac{ceX^2}{n})$
    - vi.  $= \exp_n(x)^k$ .
  - (c) **Hence show that**  $\exp_n(kx) \equiv \exp_n(x)^k \text{ (err } \frac{kceX^2}{n}) \text{ (err } \frac{ak}{n})$ .
6. **Yield the tuple**  $\langle a, b, p \rangle$ .

### Procedure III:39(thu2507191307)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct positive rational numbers  $a, b$ , and a procedure  $p(x, y, n)$  to show that  $\exp_n(y) \equiv \exp_n(x) \text{ (err } a(x-y))$  when two

complex numbers  $x, y$  and a positive integer  $n > b$  such that  $\|x\|^2 \leq X$  and  $\|y\|^2 \leq X$  are chosen.

### Implementation

1. Execute **procedure III:34** on  $\langle X \rangle$  and let  $\langle c, b, q \rangle$  receive.
2. Let  $a = c^2$ .
3. Let  $p(x, y, n)$  be the following procedure:
  - (a) Using procedure  $q$ , show that  $\exp_n(x) \equiv \exp_n(y)$ 
    - i.  $(\text{err } \exp_n(y) - \exp_n(x))$
    - ii.  $(\text{err } (1 + \frac{y}{n})^n - (1 + \frac{x}{n})^n)$
    - iii.  $(\text{err } (\frac{y}{n} - \frac{x}{n}) \sum_r^{[0:n]} (1 + \frac{y}{n})^r (1 + \frac{x}{n})^{n-1-r})$
    - iv.  $(\text{err } (y - x) (\frac{1}{n} \sum_r^{[0:n]} c^2))$
    - v.  $(\text{err } a(y - x))$ .
4. **Yield the tuple**  $\langle a, b, p \rangle$ .

### Procedure III:40(3.21)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, N$ , and a procedure,  $p(x, n)$ , to show that  $\exp_n(x) \equiv \sum_r^{[0:n+1]} \frac{x^r}{r!} \text{ (err } \frac{a}{n})$  when a complex number  $x$  and an integer  $n > N$  such that  $\|x\|^2 \leq X^2$  are chosen.


### Implementation

1. Let  $N = \lfloor X \rfloor + 1$ .
2. Let  $a = X^2 (\sum_r^{[0:N]} \frac{X^r}{r!} + \frac{X^N}{N!} \cdot \frac{1}{1 - \frac{X}{N}})$ .
3. Let  $p(x, n)$  be the following procedure:
  - (a) Using **procedure II:31**, **procedure III:16**, **procedure II:30**, and **procedure II:32**, show that  $\exp_n(x) \equiv \sum_r^{[0:n+1]} \frac{x^r}{r!}$ 
    - i.  $(\text{err } \sum_r^{[0:n+1]} \frac{x^r}{r!} - \exp_n(x))$
    - ii.  $(\text{err } \sum_r^{[0:n+1]} \frac{x^r}{r!} - \sum_r^{[0:n+1]} \frac{n^r}{r!} \cdot \frac{x^r}{n^r})$
    - iii.  $(\text{err } \sum_r^{[1:n+1]} (1 - \frac{n^r}{n^r}) \frac{x^r}{r!})$

- iv.  $(\text{err } \sum_r^{[1:n+1]} (1 - \frac{n^r}{n^r}) \frac{X^r}{r!})$   
v.  $(\text{err } \sum_r^{[2:n+1]} (1 - \frac{(n-r+1)^r}{n^r}) \frac{X^r}{r!})$   
vi.  $(\text{err } \sum_r^{[2:n+1]} (1 - (1 - \frac{r-1}{n})^r) \frac{X^r}{r!})$   
vii.  $(\text{err } \sum_r^{[2:n+1]} (1 - (1 - \frac{(r-1)r}{n}) \frac{X^r}{r!})$   
viii.  $(\text{err } \sum_r^{[2:n+1]} \frac{(r-1)r}{n} \frac{X^r}{r!})$   
ix.  $(\text{err } \frac{1}{n} \sum_r^{[2:n+1]} \frac{X^r}{(r-2)!})$   
x.  $(\text{err } \frac{X^2}{n} \sum_r^{[0:n-1]} \frac{X^r}{r!})$   
xi.  $(\text{err } \frac{X^2}{n} (\sum_r^{[0:N]} \frac{X^r}{r!} + \sum_r^{[N:n-1]} \frac{X^r}{r!}))$   
xii.  $(\text{err } \frac{X^2}{n} (\sum_r^{[0:N]} \frac{X^r}{r!} + \sum_r^{[N:n-1]} \frac{X^r}{N!N^{r-N}}))$   
xiii.  $(\text{err } \frac{X^2}{n} (\sum_r^{[0:N]} \frac{X^r}{r!} + \frac{X^N}{N!} \sum_r^{[N:n-1]} \frac{X^{r-N}}{N^{r-N}}))$   
xiv.  $(\text{err } \frac{X^2}{n} (\sum_r^{[0:N]} \frac{X^r}{r!} + \frac{X^N}{N!} \sum_r^{[0:n-N-1]} \frac{X^r}{N^r}))$   
xv.  $(\text{err } \frac{X^2}{n} (\sum_r^{[0:N]} \frac{X^r}{r!} + \frac{X^N}{N!} \cdot \frac{1}{1-\frac{X}{N}}))$   
xvi.  $(\text{err } \frac{a}{n}).$

4. Yield the tuple  $\langle a, N, p \rangle$ .

Figure III:0



A plot of the list of complex numbers  $(1 + \frac{4i}{10})^{[0:11]}$ . Notice that each multiplication of a complex number by  $1 + \frac{4i}{10}$  results in an anti-clockwise rotation about the origin and a small radial movement outwards. This can be seen to reflect the computation  $(1 + \frac{4i}{10})a = 1a + \frac{4}{10}(ai)$  after one notes that  $ai$  is perpendicular to  $a$ . Also note that each line segment has a length of roughly  $\frac{4}{10}$  units. Hence the entire path has a length of approximately  $10 * \frac{4}{10} = 4$  units.

#### Declaration III:14

The notation  $\text{cos}_n(z)$ , where  $z$  is a complex number and  $n$  is a positive integer, will be used as a shorthand for  $\frac{\exp_n(iz) + \exp_n(-iz)}{2}$ .

#### Procedure III:41(3.22)

##### Objective

Choose a rational number  $x$  and a positive integer  $n$ . The objective of the following instructions is to show that  $\text{re}(\exp_n(ix)) = \text{cos}_n(x)$ .

##### Implementation

1. Show that  $\text{re}(\exp_n(ix))$

$$(a) = \frac{\exp_n(ix) + \overline{\exp_n(ix)}}{2}$$

$$(b) = \frac{\exp_n(ix) + \exp_n(\overline{ix})}{2}$$

$$(c) = \frac{\exp_n(ix) + \exp_n(-ix)}{2}$$

$$(d) = \text{cos}_n(x).$$

#### Declaration III:15

The notation  $\text{sin}_n(z)$ , where  $z$  is a complex number and  $n$  is a positive integer, will be used as a shorthand for  $\frac{\exp_n(iz) - \exp_n(-iz)}{2i}$ .

### Procedure III:42(3.23)

#### Objective

Choose a rational number  $x$  and a positive integer  $n$ . The objective of the following instructions is to show that  $\text{im}(\exp_n(ix)) = \sin_n(x)$ .

#### Implementation

1. Show that  $\text{im}(\exp_n(ix))$

$$(a) = \frac{\exp_n(ix) - \overline{\exp_n(ix)}}{2i}$$

$$(b) = \frac{\exp_n(ix) - \exp_n(\overline{ix})}{2i}$$

$$(c) = \frac{\exp_n(ix) - \exp_n(-ix)}{2i}$$

$$(d) = \sin_n(x).$$

### Procedure III:43(3.24)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, b$ , and a procedure,  $p(x, y, n)$ , to show that  $\cos_n(x + y) \equiv \cos_n(x) \cos_n(y) - \sin_n(x) \sin_n(y)$  (err  $\frac{axy}{n}$ ) (err  $\frac{aX^2}{n}$ ) when two complex numbers  $x, y$  and a positive integer  $n > b$  such that  $\|x\|^2 \leq X^2$  and  $\|y\|^2 \leq X^2$  are chosen.

#### Implementation

1. Execute [procedure III:36](#) on  $\langle X \rangle$  and let  $\langle a, b, q \rangle$  receive.
2. Let  $p(x, y, n)$  be the following procedure:
  - (a) Using procedure  $q$ , show that  $\cos_n(x + y)$ 
    - i.  $= \frac{1}{2}(\exp_n(i(x + y)) + \exp_n(-i(x + y)))$
    - ii.  $\equiv \frac{1}{2}(\exp_n(ix) \exp_n(iy) + \exp_n(-i(x + y)))$  (err  $\frac{a(ix)(iy)}{2n}$ )
    - iii.  $\equiv \frac{1}{2}(\exp_n(ix) \exp_n(iy) + \exp_n(-ix) \exp_n(-iy))$  (err  $\frac{a(-ix)(-iy)}{2n}$ )
    - iv.  $= \frac{1}{4}(\exp_n(ix) \exp_n(iy) + \exp_n(-ix) \exp_n(-iy)) + \frac{1}{4}(\exp_n(ix) \exp_n(iy) + \exp_n(-ix) \exp_n(-iy))$

$$\begin{aligned} \text{v.} &= \frac{1}{4}(\exp_n(ix)(\exp_n(iy) + \exp_n(-iy)) + \\ &(\exp_n(-ix) - \exp_n(ix)) \exp_n(-iy)) + \\ &\frac{1}{4}((\exp_n(ix) - \exp_n(-ix)) \exp_n(iy) + \\ &\exp_n(-ix)(\exp_n(iy) + \exp_n(-iy))) \end{aligned}$$

$$\text{vi.} = \frac{1}{2} \exp_n(ix) \cos_n(y) + \frac{1}{2i} \sin_n(x) \exp_n(-iy) - \frac{1}{2i} \sin_n(x) \exp_n(iy) + \frac{1}{2} \exp_n(-ix) \cos_n(y)$$

$$\text{vii.} = \cos_n(x) \cos_n(y) - \sin_n(x) \sin_n(y)$$

$$\begin{aligned} \text{(b) Hence show that } \cos_n(x + y) &\equiv \\ \cos_n(x) \cos_n(y) - \sin_n(x) \sin_n(y) &(\text{err } \frac{axy}{n}) (\text{err } \frac{aX^2}{n}). \end{aligned}$$

3. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:44(3.25)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, b$ , and a procedure,  $p(x, y, n)$ , to show that  $\sin_n(x + y) \equiv \sin_n(x) \cos_n(y) - \cos_n(x) \sin_n(y)$  (err  $\frac{axy}{n}$ ) (err  $\frac{aX^2}{n}$ ) when two complex numbers  $x, y$  and a positive integer  $n > b$  such that  $\|x\|^2 \leq X^2$  and  $\|y\|^2 \leq X^2$  are chosen.

#### Implementation

Implementation is analogous to that of [procedure III:43](#).

### Procedure III:45(3.26)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, b$ , and a procedure,  $p(x, n)$ , to show that  $\cos_n(x)^2 + \sin_n(x)^2 \equiv 1$  (err  $\frac{a\|x\|^2}{n}$ ) (err  $\frac{aX^2}{n}$ ) when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$  and  $n > b$  are chosen.

#### Implementation

1. Execute [procedure III:36](#) on  $\langle X \rangle$  and let  $\langle a, b, q \rangle$  receive.
2. Let  $p(x, n)$  be the following procedure:

(a) Using procedure  $q$ , show that  $\cos_n(x)^2 + \sin_n(y)^2$

$$\text{i.} = \frac{1}{4}(\exp_n(ix) + \exp_n(-ix))^2 + \frac{1}{4i^2}(\exp_n(ix) - \exp_n(-ix))^2$$

$$\text{ii.} = \frac{1}{4}(\exp_n(ix)^2 + 2\exp_n(ix)\exp_n(-ix) + \exp_n(-ix)^2) - \exp_n(ix)^2 + 2\exp_n(ix)\exp_n(-ix) - \exp_n(-ix)^2$$

$$\text{iii.} = \exp_n(ix)\exp_n(-ix)$$

$$\text{iv.} \equiv 1 \text{ (err } \frac{a(-ix)(ix)}{n} \text{)}.$$

(b) Hence show that  $\cos_n(x)^2 + \sin_n(y)^2 \equiv 1 \text{ (err } \frac{a\|x\|^2}{n} \text{) (err } \frac{aX^2}{n} \text{)}.$

3. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:46(sat0308190647)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, b$ , and a procedure,  $p(x, y, n)$ , to show that  $\|x \exp_n(iy)\|^2 \equiv \|x\|^2 \text{ (err } \frac{a\|x\|^2\|y\|^2}{n} \text{) (err } \frac{a\|x\|^2X^2}{n} \text{)}$  when a complex number  $x$ , a rational number  $y$ , and a positive integer  $n$  such that  $\|y\|^2 \leq X^2$  and  $n > b$  are chosen.

#### Implementation

1. Execute **procedure III:45** on  $\langle X \rangle$  and let  $\langle a, b, q \rangle$  receive.

2. Let  $p(x, y, n)$  be the following procedure:

(a) Using procedure  $q$ , **procedure III:41**, and **procedure III:42**, show that  $\|x \exp_n(iy)\|^2$

$$\text{i.} = \|x\|^2 \|\exp_n(iy)\|^2$$

$$\text{ii.} = \|x\|^2 \|\cos_n(y) + i \sin_n(y)\|^2$$

$$\text{iii.} = \|x\|^2 (\cos_n(y)^2 + \sin_n(y)^2)$$

$$\text{iv.} \equiv \|x\|^2 \cdot 1 \text{ (err } \|x\|^2 \cdot \frac{a\|y\|^2}{n} \text{)}.$$

(b) Hence show that  $\|x \exp_n(iy)\|^2 \equiv \|x\|^2 \text{ (err } \frac{a\|xy\|^2}{n} \text{) (err } \frac{a\|x\|^2X^2}{n} \text{)}.$

### Procedure III:47(3.29)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, N$ , and a procedure,  $p(x, n)$ , to show that  $\cos_n(x) \equiv \sum_r^{[0: \lceil \frac{n}{2} \rceil]} \frac{(-1)^r x^{2r}}{(2r)!} \text{ (err } \frac{a}{n} \text{)}$  when a complex number  $x$  and an integer  $n > N$  such that  $\|x\|^2 \leq X^2$  is chosen.

#### Implementation

1. Execute **procedure III:40** on  $\langle X \rangle$  and let  $\langle a, N, q \rangle$  receive.

2. Let  $p(x, n)$  be the following procedure:

(a) Using procedure  $q$ , show that  $\cos_n(x)$

$$\text{i.} = \frac{\exp_n(ix)}{2} + \frac{\exp_n(-ix)}{2}$$

$$\text{ii.} \equiv \frac{1}{2} \sum_r^{[0:n+1]} \frac{(ix)^r}{r!} + \frac{\exp_n(-ix)}{2} \text{ (err } \frac{a}{2n} \text{)}$$

$$\text{iii.} \equiv \frac{1}{2} \sum_r^{[0:n+1]} \frac{(ix)^r}{r!} + \frac{1}{2} \sum_r^{[0:n+1]} \frac{(-ix)^r}{r!} \text{ (err } \frac{a}{2n} \text{)}$$

$$\text{iv.} = \sum_r^{[0:n+1]} \frac{(i^r + (-i)^r)x^r}{2(r!)}$$

$$\text{v.} = \sum_r^{[0:n+1]} \frac{[r \bmod 2=0](-1)^{\frac{r}{2}}x^r}{r!}$$

$$\text{vi.} = \sum_r^{[0: \lceil \frac{n}{2} \rceil]} \frac{(-1)^r x^{2r}}{(2r)!}.$$

(b) Hence show that  $\cos_n(x) \equiv \sum_r^{[0: \lceil \frac{n}{2} \rceil]} \frac{(-1)^r x^{2r}}{(2r)!} \text{ (err } \frac{a}{n} \text{)}.$

### Procedure III:48(3.30)

#### Objective

Choose a rational number  $X \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, N$ , and a procedure,  $p(x, n)$ , to show that  $\sin_n(x) \equiv \sum_r^{[0: \lfloor \frac{n+1}{2} \rfloor]} \frac{(-1)^r x^{2r+1}}{(2r+1)!} \text{ (err } \frac{a}{n} \text{)}$  when a complex number  $x$  and an integer  $n > N$  such that  $\|x\|^2 \leq X^2$  is chosen.

#### Implementation

Implementation is analogous to that of **procedure III:47**.

# Chapter 10

## Binomial and Mercator Series

### Declaration III:16

The notation  $(1+x)_n^a$ , where  $x, a$  are complex numbers and  $n$  is a positive integer, will be used as a shorthand for  $\sum_r^{[0:n]} \binom{a}{r} x^r$ .

### Procedure III:49(sun2107190619)

#### Objective

Choose a complex number  $x$  and two non-negative integers  $a, n$  such that  $n > a$ . The objective of the following instructions is to show that  $(1+x)_n^a = (1+x)^a$ .

#### Implementation

1. Using [procedure III:12](#), show that  $(1+x)_n^a =$

$$(a) = \sum_r^{[0:n]} \binom{a}{r} x^r$$

$$(b) = \sum_r^{[0:n]} \frac{a^r}{r!} x^r$$

$$(c) = \sum_r^{[0:a+1]} \frac{a^r}{r!} x^r + \sum_r^{[a+1:n]} \frac{a^r}{r!} x^r$$

$$(d) = \sum_r^{[0:a+1]} \frac{a^r}{r!} x^r + \sum_r^{[a+1:n]} \frac{0}{r!} x^r$$

$$(e) = \sum_r^{[0:a+1]} \binom{a}{r} x^r$$

$$(f) = (1+x)^a.$$

### Procedure III:50(sun2107190640)

#### Objective

Choose two complex numbers  $x, y$  and a positive integer  $N$ . The objective of the following instructions is to show that  $\binom{x+y}{N} = \sum_k^{N+1} \binom{x}{k} \binom{y}{N-k}$ .

#### Implementation

1. If  $N = 0$ , then do the following:

$$(a) \text{ Show that } \binom{x+y}{N} = 1 = \sum_k^{[0:N+1]} \binom{x}{k} \binom{y}{N-k}.$$

2. Otherwise do the following:

(a) Show that  $N > 0$ .

$$(b) \text{ Show that } \binom{x+y-1}{N-1} = \sum_k^{[0:N]} \binom{x-1}{k} \binom{y}{N-1-k} \text{ using } \textcolor{red}{\text{procedure III:50}}.$$

$$(c) \text{ Show that } \binom{x+y-1}{N-1} = \sum_k^{[0:N]} \binom{x}{k} \binom{y-1}{N-1-k} \text{ using } \textcolor{red}{\text{procedure III:50}}.$$

(d) Hence show that  $\binom{x+y}{N}$

$$\text{i.} = \frac{x+y}{N} \binom{x+y-1}{N-1}$$

$$\text{ii.} = \frac{x}{N} \binom{x+y-1}{N-1} + \frac{y}{N} \binom{x+y-1}{N-1}$$

$$\text{iii.} = \frac{x}{N} \sum_k^{[0:N]} \binom{x-1}{k} \binom{y}{N-1-k} + \frac{y}{N} \sum_k^{[0:N]} \binom{x}{k} \binom{y-1}{N-1-k}$$

$$\text{iv.} = \frac{x}{N} \sum_k^{[1:N+1]} \binom{x-1}{k-1} \binom{y}{N-k} + \frac{y}{N} \sum_k^{[0:N]} \binom{x}{k} \binom{y-1}{N-1-k}$$

$$\text{v.} = \sum_k^{[0:N+1]} \frac{k}{N} \binom{x}{k} \binom{y}{N-k} + \sum_k^{[0:N+1]} \frac{N-k}{N} \binom{x}{k} \binom{y}{N-k}$$

$$\text{vi.} = \sum_k^{[0:N+1]} \binom{x}{k} \binom{y}{N-k}.$$

### Procedure III:51(sun2107191133)

#### Objective

Choose complex numbers  $a, b, x$  and a natural number  $n$ . The objective of the following instructions is to show that  $(1+x)_n^a(1+x)_n^b - (1+x)_n^{a+b} = \sum_k^{[1:n]} \sum_r^{[k:n]} \binom{a}{k+n-1-r} \binom{b}{r} x^{k+n-1}$ .

#### Implementation

1. Show that  $(1+x)_n^a(1+x)_n^b - (1+x)_n^{a+b}$

$$(a) = \left( \sum_k^{[0:n]} \binom{a}{k} x^k \right) \left( \sum_r^{[0:n]} \binom{b}{r} x^r \right) - \sum_k^{[0:n]} \binom{a+b}{k} x^k$$

$$(b) = \sum_k^{[0:n]} \sum_r^{[0:n]} \binom{a}{k} \binom{b}{r} x^{k+r} - \sum_k^{[0:n]} \binom{a+b}{k} x^k$$

$$(c) = \sum_k^{[0:n]} \sum_r^{[0:k+1]} \binom{a}{k-r} \binom{b}{r} x^k + \sum_k^{[n:2n-1]} \sum_r^{[k-n+1:n]} \binom{a}{k-r} \binom{b}{r} x^k$$

$$(d) = \sum_k^{[0:n]} \binom{a+b}{k} x^k + \sum_k^{[1:n]} \sum_r^{[k:n]} \binom{a}{k+n-1-r} \binom{b}{r} x^{k+n-1}$$

$$(e) = \sum_k^{[1:n]} \sum_r^{[k:n]} \binom{a}{k+n-1-r} \binom{b}{r} x^{k+n-1}.$$

### Procedure III:52(sun2107191247)

#### Objective

Choose two rational numbers  $A > 0$  and  $0 < X < 1$ . The objective of the following instructions is to construct rational numbers  $Y > 0$ ,  $0 < Z < 1$  and a procedure  $p(a, x, n)$  to show that  $\|(a)_n x^n\|^2 \leq (YZ^n)^2$  when complex numbers  $a, x$  such that  $\|a+1\|^2 < A^2$  and  $\|x\|^2 < X^2$  are chosen.

#### Implementation

1. Let  $e = \frac{AX}{1-X} - 1$ .

2. Let  $d = \lfloor \frac{AX}{1-X} \rfloor$ .

3. Show that  $d > e > -1$ .

4. Let  $Z = (1 + \frac{A}{d+1})X$ .

5. Show that  $0 < Z < (1 + \frac{A}{e+1})X = 1$ .

6. Let  $Y = Z^{-d} \prod_k^{[0:d]} \frac{(A+k+1)X}{k+1} = Z^{-d} \prod_k^{[0:d]} X(1 + \frac{A}{k+1})$ .

7. Let  $p(a, x, n)$  be the following procedure:

(a) Show that  $|\operatorname{re}(a+1)| \leq A$  given that  $\operatorname{re}(a+1)^2 \leq \|a+1\|^2 \leq A^2$ .

(b) Hence show that  $\|(a)_n x^n\|^2$

$$\text{i.} = \|\frac{a^n}{n!} x^n\|^2$$

$$\text{ii.} = \|\prod_k^{[0:n]} (\frac{a+1-(k+1)}{k+1} \cdot x)\|^2$$

$$\text{iii.} = \prod_k^{[0:n]} \frac{\|(a+1)-(k+1)\|^2 \|x\|^2}{(k+1)^2}$$

$$\text{iv.} = \prod_k^{[0:n]} \frac{(\|a+1\|^2 - 2\operatorname{re}(a+1)(k+1) + (k+1)^2) \|x\|^2}{(k+1)^2}$$

$$\text{v.} \leq \prod_k^{[0:n]} \frac{(A^2 + 2A(k+1) + (k+1)^2) X^2}{(k+1)^2}$$

$$\text{vi.} = \left( \prod_k^{[0:n]} \frac{(A+k+1)X}{k+1} \right)^2$$

$$\text{vii.} = \left( \prod_k^{[0:n]} X(1 + \frac{A}{k+1}) \right)^2.$$

(c) If  $n \leq d$ , then do the following:

i. Show that  $\|(a)_n x^n\|^2$

$$A. \leq \left( \prod_k^{[0:n]} X(1 + \frac{A}{k+1}) \right)^2$$

$$B. = \left( \prod_k^{[0:d]} X(1 + \frac{A}{k+1}) \right)^2 \left( \prod_k^{[n:d]} X(1 + \frac{A}{k+1}) \right)^{-2}$$

$$C. \leq \left( \prod_k^{[0:d]} X(1 + \frac{A}{k+1}) \right)^2 (X(1 + \frac{A}{d+1}))^{-2(d-n)}$$

$$D. = Y^2 Z^{2n}.$$

(d) Otherwise do the following:

i. Show that  $\|(a)_n x^n\|^2$

$$A. \leq \left( \prod_k^{[0:n]} X(1 + \frac{A}{k+1}) \right)^2$$

$$B. = \left( \prod_k^{[0:d]} X(1 + \frac{A}{k+1}) \right)^2 \left( \prod_k^{[d:n]} X(1 + \frac{A}{k+1}) \right)^2$$

$$C. \leq \left( \prod_k^{[0:d]} X(1 + \frac{A}{k+1}) \right)^2 (X(1 + \frac{A}{d+1}))^{2(n-d)}$$

$$D. = Y^2 Z^{2n}.$$

8. Yield the tuple  $\langle Y, Z, p \rangle$ .

### Procedure III:53(wed2407191422)

#### Objective

Choose a rational number  $0 < X < 1$  and a positive integer  $k$ . The objective of the following instructions is to construct rational numbers  $Y > 0$ ,  $0 < Z < 1$  and a procedure  $p(x, n)$  to show that  $\|n^k x^n\|^2 \leq (Y Z^n)^2$  when a complex number  $x$  such that  $\|x\|^2 \leq X^2$  is chosen.

#### Implementation

1. Let  $e = \frac{k}{1-X} - 1$ .
2. Let  $d = \lfloor \frac{k}{1-X} \rfloor$ .
3. Show that  $d > e > k - 1$ .
4. Let  $Z = (1 + \frac{1}{d})^k X$ .
5. Show that  $Z < (1 + \frac{1}{e})^k X$ .
6. Now show that  $0 < Z < (1 + \frac{1}{e})^k X \leq \frac{1+\frac{1}{e}}{1-(k-1)\frac{1}{e}} \cdot X = 1$  using [procedure II:33](#).
7. Let  $Y = Z^{-d} X \prod_r^{[1:d]} X(1 + \frac{1}{r})^k$ .
8. Let  $p(x, n)$  be the following procedure:
  - (a) Show that  $\|n^k x^n\|^2$ 
    - i.  $\leq \|x \prod_r^{[1:n]} x \cdot \frac{(r+1)^k}{r^k}\|^2$
    - ii.  $= \|x\|^2 \prod_r^{[1:n]} \|x\|^2 (\frac{(r+1)^k}{r^k})^2$
    - iii.  $\leq X^2 \prod_r^{[1:n]} ((1 + \frac{1}{r})^k X)^2$ .
  - (b) If  $n \leq d$ , then do the following:
    - i. Show that  $\|n^k x^n\|^2$ 
      - A.  $\leq X^2 (\prod_r^{[1:n]} X(1 + \frac{1}{r})^k)^2$
      - B.  $= X^2 (\prod_r^{[1:d]} X(1 + \frac{1}{r})^k)^2 \cdot (\prod_r^{[n:d]} X(1 + \frac{1}{r})^k)^{-2}$
      - C.  $\leq X^2 (\prod_r^{[1:d]} X(1 + \frac{1}{r})^k)^2 (X(1 + \frac{1}{d})^k)^{-2(d-n)}$
      - D.  $= Y^2 Z^{2n}$ .
  - (c) Otherwise do the following:
    - i. Show that  $\|n^k x^n\|^2$ 
      - A.  $\leq X^2 (\prod_r^{[1:n]} X(1 + \frac{1}{r})^k)^2$

$$B. = X^2 (\prod_r^{[1:d]} X(1 + \frac{1}{r})^k)^2 (\prod_r^{[d:n]} X(1 + \frac{1}{r})^k)^2$$

$$C. \leq X^2 (\prod_r^{[1:d]} X(1 + \frac{1}{r})^k)^2 (X(1 + \frac{1}{d})^k)^{2(n-d)}$$

$$D. = Y^2 Z^{2n}.$$

9. Yield the tuple  $\langle Y, Z, p \rangle$ .

### Procedure III:54(wed2407191521)

#### Objective

Choose two rational numbers  $A > 0$ ,  $1 > X > 0$ . The objective of the following instructions is to construct rational numbers  $D > 0$ ,  $0 < G < 1$ , and a procedure  $p(x, a, b, n)$  to show that  $(1+x)_n^{a+b} \equiv (1+x)_n^a (1+x)_n^b$  (err  $DG^n$ ) when  $\|x\|^2 \leq X$ , and  $\|a\|^2, \|b\|^2 < A$ .

#### Implementation

1. Execute [procedure III:52](#) on  $\langle A, X \rangle$  and let  $\langle B, C, q \rangle$  receive.
2. Execute [procedure III:53](#) on  $\langle C, 1 \rangle$  and let  $\langle F, G, t \rangle$  receive.
3. Let  $D = \frac{B^2 F}{1-C}$ .
4. Let  $p(x, a, b, n)$  be the following procedure:
  - (a) For each  $r \in [1 : n]$ , do the following:
    - i. Show that  $\|\binom{a}{r} x^r\|^2 \leq (BC^r)^2$  using procedure  $q$ .
    - ii. Show that  $\|\binom{b}{r} x^r\|^2 \leq (BC^r)^2$  using procedure  $q$ .
  - (b) Show that  $\|n C^n\|^2 \leq (FG^n)^2$  using procedure  $t$ .
  - (c) Hence show that  $(1+x)_n^{a+b} \equiv (1+x)_n^a (1+x)_n^b$ 
    - i. (err  $(1+x)_n^a (1+x)_n^b - (1+x)_n^{a+b}$ )
    - ii. (err  $\sum_k^{[1:n]} \sum_r^{[k:n]} \binom{a}{k+n-1-r} \binom{b}{r} x^{k+n-1}$ )
    - iii. (err  $\sum_k^{[1:n]} \sum_r^{[k:n]} \binom{a}{k+n-1-r} x^{k+n-1-r} \binom{b}{r} x^r$ )
    - iv. (err  $\sum_k^{[1:n]} \sum_r^{[k:n]} BC^{k+n-1-r} BC^r$ )
    - v. (err  $B^2 C^n \sum_k^{[1:n]} \sum_r^{[k:n]} C^{k-1}$ )
    - vi. (err  $B^2 C^n \sum_r^{[1:n]} \sum_k^{[r+1:n]} C^{k-1}$ )



- vii. (err  $B^2 C^n \sum_r^{[1:n]} \frac{1}{1-C}$ )
- viii. (err  $\frac{B^2}{1-C} \cdot n C^n$ )
- ix. (err  $\frac{B^2 F}{1-C} G^n$ )
- x. (err  $D G^n$ ).

5. Yield the tuple  $\langle D, G, p \rangle$ .

### Procedure III:55(wed2407191611)

#### Objective

Choose two rational numbers  $A > 0$ ,  $1 > X > 0$ . The objective of the following instructions is to construct a rational number  $D$  and a procedure  $p(x, n, a, k)$  to show that  $\|(1+x)_n^a\|^2 < D^2$  when complex numbers  $x, a$  and positive integers  $n, k$  such that  $\|x\|^2 < X^2$  and  $\|ka\|^2 < A^2$ .

#### Implementation

1. Execute **procedure III:34** on  $\langle \frac{ABX}{1-C} \rangle$  and let  $\langle E, N, t \rangle$  receive.
2. Execute **procedure III:52** on  $\langle A+1, X \rangle$  and let  $\langle B, C, q \rangle$  receive.
3. Let  $D = \max(E, (1 + \frac{ABX}{1-C})^{\lfloor N \rfloor})$ .
4. Let  $p(x, n, a, k)$  be the following procedure:
  - (a) For each  $r \in [1 : n]$ , do the following:
    - i. Show that  $\|a\|^2 \leq \|ka\|^2 \leq A^2$ .
    - ii. Show that  $\|a-1\|^2 \leq (A+1)^2$ .
    - iii. Hence show that  $\|(\frac{a-1}{r-1})x^{r-1}\|^2 \leq (BC^r)^2$  using procedure  $q$ .
  - (b) Hence show that  $\|k \sum_r^{[1:n]} \binom{a}{r} x^r\|^2$ 
    - i.  $= \|k \sum_r^{[1:n]} \frac{a}{r} \binom{a-1}{r-1} x^r\|^2$
    - ii.  $= \|kax \sum_r^{[1:n]} \frac{1}{r} \binom{a-1}{r-1} x^{r-1}\|^2$
    - iii.  $\leq (AX \sum_r^{[1:n]} BC^{r-1})^2$
    - iv.  $\leq (\frac{ABX}{1-C})^2$ .
  - (c) If  $k > N$ , then do the following:
    - i. Hence using procedure  $t$ , show that  $\|(1+x)_n^a\|^2$ 
      - A.  $= \|(\sum_r^{[0:n]} \binom{a}{r} x^r)^k\|^2$

- B.  $= \|(1 + \sum_r^{[1:n]} \binom{a}{r} x^r)^k\|^2$
- C.  $= \|\exp_k(k \sum_r^{[1:n]} \binom{a}{r} x^r)\|^2$
- D.  $\leq E^2$
- E.  $\leq D^2$ .

(d) Otherwise do the following:

- i. Show that  $\|\sum_r^{[1:n]} \binom{a}{r} x^r\|^2$ 
  - A.  $\leq \|k \sum_r^{[1:n]} \binom{a}{r} x^r\|^2$
  - B.  $\leq (\frac{ABX}{1-C})^2$ .
- ii. Hence show that  $\|(1+x)_n^a\|^2$ 
  - A.  $= (\|(1+x)_n^a\|^2)^k$
  - B.  $= (\|1 + \sum_r^{[1:n]} \binom{a}{r} x^r\|^2)^k$
  - C.  $\leq (1 + \frac{ABX}{1-C})^{2k}$
  - D.  $\leq D^2$ .

5. Yield  $\langle D, p \rangle$ .

### Procedure III:56(tue2008190712)

#### Objective

Choose two rational numbers  $A > 0$ ,  $1 > X > 0$ . The objective of the following instructions is to construct positive rational numbers  $D, N$ , and a procedure  $p(x, a, n)$  to show that  $\|(1+x)_n^a\|^2 \geq D^2$  when complex numbers  $x, a$  and an integer  $n$  such that  $\|x\|^2 \leq X^2$ ,  $\|a\| \leq A^2$ , and  $n > N$  are chosen.

#### Implementation

1. Execute **procedure III:54** on  $\langle A, X \rangle$  and let  $\langle a_1, b_1, p_1 \rangle$  receive.
2. Execute **procedure III:53** on  $\langle b_1, 1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:55** on  $\langle A, X \rangle$  and let  $\langle a_3, p_3 \rangle$  receive.
4. Let  $D = \frac{1}{2a_3}$ .
5. Let  $N = 2a_1a_2$ .
6. Let  $p(x, a, n)$  be the following procedure:
  - (a) Show that  $\|nb_1^n\|^2 \leq (a_2b_2^n)^2 \leq a_2^2$  using procedure  $p_2$ .

- (b) Hence show that  $(a_1 b_1^n)^2 \leq (\frac{a_1 a_2}{n})^2$ .
- (c) Show that  $\|(1+x)_n^{-a}\|^2 \leq a_3^2$  using procedure  $p_3$ .
- (d) Using procedure  $p_1$ , show that  $\|(1+x)_n^{-a}(1+x)_n^a - 1\|^2$
- $\|(1+x)_n^{-a}(1+x)_n^a - (1+x)_n^{-a+a}\|^2$
  - $\leq (a_1 b_1^n)^2$ .
- (e) Hence using **procedure III:17**, show that  $\frac{1}{2} - \|(1+x)_n^{-a}(1+x)_n^a\|^2$
- $\frac{1}{2}\|1\|^2 - \|(1+x)_n^{-a}(1+x)_n^a\|^2$
  - $\leq \|1 - (1+x)_n^{-a}(1+x)_n^a\|^2$
  - $\leq (a_1 b_1^n)^2$
  - $\leq (\frac{a_1 a_2}{n})^2$
  - $\leq \frac{1}{4}$ .
- (f) Hence show that  $(\frac{1}{2})^2$
- $\leq \|(1+x)_n^{-a}(1+x)_n^a\|^2$
  - $\leq a_3^2 \|(1+x)_n^a\|^2$ .
- (g) **Hence show that  $D^2 \leq \|(1+x)_n^a\|^2$ .**

7. **Yield the tuple  $\langle D, N, p \rangle$ .**

### Procedure III:57(tue2008190849)

#### Objective

Choose two rational numbers  $A > 0$  and  $1 > X > 0$ . The objective of the following instructions is to construct positive rational numbers  $B, C, D$ , and a procedure  $p(x, a, b, n)$  to show that  $(1+x)_n^{a-b} \equiv \frac{(1+x)_n^a}{(1+x)_n^b}$  (err  $BC^n$ ) when complex numbers  $x, a, b$  and an integer  $n$  such that  $\|x\|^2 \leq X^2$ ,  $\|a\|^2 \leq A^2$ ,  $\|b\|^2 \leq A^2$ , and  $n > D$  are chosen.

#### Implementation

- Execute **procedure III:54** on  $\langle A, X \rangle$  and let  $\langle a_1, C, p_1 \rangle$  receive.
- Execute **procedure III:56** on  $\langle A, X \rangle$  and let  $\langle a_2, D, p_2 \rangle$  receive.
- Execute **procedure III:55** on  $\langle A, X \rangle$  and let  $\langle a_3, p_3 \rangle$  receive.

- Let  $B = (1 + \frac{a_3}{a_2})a_1$ .
- Let  $p(x, a, b, n)$  be the following procedure:
  - Using procedures  $p_1, p_2, p_3$ , show that  $(1+x)_n^{a-b}$ 
    - $\equiv (1+x)_n^a(1+x)_n^{-b}$  (err  $a_1 C^n$ )
    - $\equiv (1+x)_n^a \frac{(1+x)_n^b (1+x)_n^{-b}}{(1+x)_n^b}$
    - $\equiv ((1+x)_n^a)^1 \frac{(1+x)_n^{b-b}}{(1+x)_n^b}$  (err  $a_3 \frac{a_1 C^n}{a_2}$ )
    - $\equiv \frac{(1+x)_n^a}{(1+x)_n^b}$
- Hence show that  $(1+x)_n^{a-b} \equiv \frac{(1+x)_n^a}{(1+x)_n^b}$  (err  $(1 + \frac{a_3}{a_2})a_1 C^n$ ) (err  $BC^n$ ).**
- Yield the tuple  $\langle B, C, D, p \rangle$ .**

### Procedure III:58(wed2407191627)

#### Objective

Choose two rational numbers  $A > 0$ ,  $1 > X > 0$ . The objective of the following instructions is to construct rational numbers  $G > 0$ ,  $0 < C < 1$ , and a procedure  $p(x, n, a, k)$  to show that  $(1+x)_n^{ka} \equiv ((1+x)_n^a)^k$  (err  $GkC^n$ ) when a non-negative integer  $k$  and complex numbers  $x, a$  such that  $\|x\|^2 \leq X^2$  and  $\|ka\|^2 < A^2$  are chosen.

#### Implementation

- Execute **procedure III:55** on  $\langle A, X \rangle$  and let  $\langle D, t \rangle$  receive.
- Execute **procedure III:54** on  $\langle A, X \rangle$  and let  $\langle B, C, q \rangle$  receive.
- Let  $G = DB$ .
- Let  $p(x, n, a, k)$  be the following procedure:
  - Hence using procedures  $t, q$ , show that  $(1+x)_n^{ka}$ 
    - $\equiv ((1+x)_n^a)^0 (1+x)_n^{ka}$
    - $\equiv ((1+x)_n^a)^1 (1+x)_n^{(k-1)a}$  (err  $DBC^n$ )
    - $\equiv ((1+x)_n^a)^2 (1+x)_n^{(k-2)a}$  (err  $DBC^n$ )
    - $\vdots$
    - $\equiv ((1+x)_n^a)^k (1+x)_n^{(k-k)a}$  (err  $DBC^n$ )

- vi.  $= ((1+x)^a_n)^k$ .
- (b) **Hence show that**  $(1+x)^{ka}_n \equiv ((1+x)^a_n)^k \text{ (err } kDBC^n) \text{ (err } GkC^n)$ .
5. **Yield the tuple**  $\langle G, C, D, p \rangle$ .

### Procedure III:59(sun0812190858)

#### Objective

Choose two non-negative rational numbers  $a, b$  and two non-negative integers  $r, n$  such that  $b < r < n - a - 1$ . The objective of the following instructions is to show that  $\text{sgn}(\binom{b}{r} \binom{a}{n-r}) = \text{sgn}(\binom{b}{r+1} \binom{a}{n-r-1})$ .

#### Implementation

1. Show that  $\text{sgn}(\frac{b-r}{r+1} \cdot \frac{n-r}{a-n+r+1}) = 1$ 
  - (a) given that  $\frac{b-r}{r+1} \cdot \frac{n-r}{a-n+r+1} > 0$
  - (b) given that  $\frac{b-r}{(a+1)-(n-r)} > 0$
  - (c) given that  $r > b$  and  $n - r > a + 1$ .
2. Hence show that  $\text{sgn}(\binom{b}{r+1} \binom{a}{n-r-1})$ 
  - (a)  $= \text{sgn}(\frac{b-r}{r+1} \binom{b}{r} \cdot \frac{n-r}{a-n+r+1} \binom{a}{n-r})$
  - (b)  $= \text{sgn}(\frac{b-r}{r+1} \cdot \frac{n-r}{a-n+r+1}) \text{sgn}(\binom{b}{r} \binom{a}{n-r})$
  - (c)  $= \text{sgn}(\binom{b}{r} \binom{a}{n-r})$ .

### Procedure III:60(sun0812190920)

#### Objective

Choose two non-negative rational numbers  $a, b$  and an integer  $n \geq \lceil a \rceil + \lceil b \rceil$ . The objective of the following instructions is to show that  $\sum_r^{[0:n+1]} \|\binom{b}{r} \binom{a}{n-r}\| = \sum_r^{[0:\lceil b \rceil]} \|\binom{b}{r} \binom{a}{n-r}\| + \|\sum_r^{\lceil b \rceil : [n-a]} \binom{b}{r} \binom{a}{n-r}\| + \sum_r^{\lceil n-a \rceil : n+1} \|\binom{b}{r} \binom{a}{n-r}\|$ .

#### Implementation

1. Verify that  $\lceil b \rceil \leq \lfloor n - a \rfloor$ .
2. For  $r$  in  $[\lceil b \rceil : \lfloor n - a \rfloor]$ , do the following:
  - (a) Show that  $\text{sgn}(\binom{b}{r} \binom{a}{n-r}) = \text{sgn}(\binom{b}{r+1} \binom{a}{n-r-1})$  using **procedure III:59**.

$$3. \text{ Hence show that } \sum_r^{\lceil b \rceil : \lceil n-a \rceil} \|\binom{b}{r} \binom{a}{n-r}\| = \|\sum_r^{\lceil b \rceil : \lceil n-a \rceil} \binom{b}{r} \binom{a}{n-r}\|.$$

$$4. \text{ Hence show that } \sum_r^{[0:n+1]} \|\binom{b}{r} \binom{a}{n-r}\|$$

$$(a) = \sum_r^{[0:\lceil b \rceil]} \|\binom{b}{r} \binom{a}{n-r}\| + \sum_r^{\lceil b \rceil : \lceil n-a \rceil} \|\binom{b}{r} \binom{a}{n-r}\| + \sum_r^{\lceil n-a \rceil : n+1} \|\binom{b}{r} \binom{a}{n-r}\|$$

$$(b) = \sum_r^{[0:\lceil b \rceil]} \|\binom{b}{r} \binom{a}{n-r}\| + \|\sum_r^{\lceil b \rceil : \lceil n-a \rceil} \binom{b}{r} \binom{a}{n-r}\| + \sum_r^{\lceil n-a \rceil : n+1} \|\binom{b}{r} \binom{a}{n-r}\|.$$

### Procedure III:61(wed2407191824)

#### Objective

Choose a rational number  $A > 0$ . The objective of the following instructions is to construct rational numbers  $M > 1$ ,  $N > 0$ , and a procedure  $p(a, n)$  to show that  $\|\binom{a}{n}\|^2 \leq (\frac{M}{n})^{2(\lceil a \rceil)}$  and  $\frac{M}{n} < 1$  when a rational number  $-1 < a < A$  and an integer  $n > N$  are chosen.

#### Implementation

1. Let  $M = 2A$ .
2. Let  $N = 2A$ .
3. Let  $p(a, n)$  be the following procedure:
  - (a) Show that  $\frac{2a}{n} < \frac{2A}{2A} = 1$ 
    - i. given that  $n > N = 2A > 2a$
    - ii. and  $-1 < a < A$ .
  - (b) Show that  $n - \lfloor a \rfloor > n - a > n - \frac{n}{2} = \frac{n}{2}$ 
    - i. given that  $\frac{n}{2} > a$
    - ii. given that  $n > N = 2A > 2a$ .
  - (c) Hence show that  $\|\binom{a}{n}\|^2$ 
    - i.  $= \|\frac{a^n}{n!}\|^2$
    - ii.  $= \|\prod_k^{[0:n]} \frac{a-k}{k+1}\|^2$
    - iii.  $= \prod_k^{[0:n]} \frac{(a-k)^2}{(k+1)^2}$
    - iv.  $= \prod_k^{[0:\lceil a \rceil]} (k-a)^2 \cdot \prod_k^{[0:n]} \frac{(k+\lfloor a \rfloor+1-a)^2}{(k+1)^2}$
    - v.  $\leq (a^{\lceil a \rceil} \cdot 1^n \cdot (\frac{1}{n-\lfloor a \rfloor})^{\lceil a \rceil})^2$

$$\text{vi.} = \left(\frac{a}{n-\lfloor a \rfloor}\right)^{2\lceil a \rceil}$$

$$\text{vii.} \leq \left(\frac{2a}{n}\right)^{2\lceil a \rceil}$$

$$\text{viii.} \leq \left(\frac{M}{n}\right)^{2\lceil a \rceil}.$$

4. **Yield the tuple**  $\langle M, N, p \rangle$ .

### Procedure III:62(sun0812191002)

#### Objective

Choose a positive integer  $A$ . The objective of the following instructions is to construct a rational number  $B$ , an integer  $N$ , and a procedure  $p(a, b, n)$  to show that  $\sum_r^{[0:n+1]} \left\| \binom{b}{r} \binom{a}{n-r} \right\| \leq \frac{B}{n}$  when non-negative rational numbers  $a, b$ , and an integer  $n$  such that  $a < A, b < A$ , and  $n > N$  are chosen.

#### Implementation

1. Execute **procedure III:61** on  $\langle A \rangle$  and let  $\langle M, Q, q \rangle$  receive.
2. Let  $N = \max(2A, Q + A)$ .
3. Let  $B = M^A(M^A + 8A!A)$ .
4. Let  $p(a, b, n)$  be the following procedure:
  - (a) Show that  $n > N \geq Q$ .
  - (b) Now show that  $\left\| \binom{a+b}{n} \right\| \leq \left(\frac{M}{n}\right)^{\lceil a+b \rceil} \leq \frac{M^{\lceil A+b \rceil}}{n} \leq \frac{M^{A+\lceil b \rceil}}{n} \leq \frac{M^{2A}}{n}$  using procedure  $q$ .
  - (c) For  $r$  in  $[0 : \lceil b \rceil]$ , do the following:
    - i. Show that  $n - r \geq N - \lceil b \rceil \geq Q + A - A = Q$ .
    - ii. Now show that  $\left\| \binom{a}{n-r} \right\|^2 \leq \left(\frac{M}{n-r}\right)^{2\lceil a \rceil} \leq \left(\frac{M^A}{r}\right)^2 \leq \left(\frac{M^A}{n-\lfloor a \rfloor}\right)^2 \leq \left(\frac{M^A}{n-\frac{1}{2}N}\right)^2 \leq \left(\frac{M^A}{\frac{1}{2}n}\right)^2 = \left(\frac{2M^A}{n}\right)^2$  using procedure  $q$ .
  - (d) For  $r$  in  $[\lceil n-a \rceil : n+1]$ , do the following:
    - i. Show that  $r \geq \lceil n-a \rceil = n - \lfloor a \rfloor \geq N - \lfloor a \rfloor \geq Q + A - A = Q$ .
    - ii. Now show that  $\left\| \binom{b}{r} \right\|^2 \leq \left(\frac{M}{r}\right)^{2\lceil b \rceil} \leq \left(\frac{M^A}{r}\right)^2 \leq \left(\frac{2M^A}{n}\right)^2$  using procedure  $q$ .

(e) Hence using **procedure III:60**, show that  $\sum_r^{[0:n+1]} \left\| \binom{b}{r} \binom{a}{n-r} \right\|$

$$\text{i.} = \sum_r^{[0:\lceil b \rceil]} \left\| \binom{b}{r} \binom{a}{n-r} \right\| + \left\| \sum_r^{[\lceil b \rceil : \lceil n-a \rceil]} \binom{b}{r} \binom{a}{n-r} \right\| + \sum_r^{[\lceil n-a \rceil : n+1]} \left\| \binom{b}{r} \binom{a}{n-r} \right\|$$

$$\text{ii.} = \sum_r^{[0:\lceil b \rceil]} \left\| \binom{b}{r} \binom{a}{n-r} \right\| + \left\| \sum_r^{[0:n+1]} \binom{b}{r} \binom{a}{n-r} \right\| - \sum_r^{[0:\lceil b \rceil]} \binom{b}{r} \binom{a}{n-r} - \sum_r^{[\lceil n-a \rceil : n+1]} \binom{b}{r} \binom{a}{n-r} + \sum_r^{[\lceil n-a \rceil : n+1]} \left\| \binom{b}{r} \binom{a}{n-r} \right\|$$

$$\text{iii.} = 2 \sum_r^{[0:\lceil b \rceil]} \left\| \binom{b}{r} \binom{a}{n-r} \right\| + \left\| \sum_r^{[0:n+1]} \binom{b}{r} \binom{a}{n-r} \right\| + 2 \sum_r^{[\lceil n-a \rceil : n+1]} \left\| \binom{b}{r} \binom{a}{n-r} \right\|$$

$$\text{iv.} = \left\| \binom{a+b}{n} \right\| + 2 \left( \sum_r^{[0:\lceil b \rceil]} \left\| \binom{b}{r} \binom{a}{n-r} \right\| + \sum_r^{[\lceil n-a \rceil : n+1]} \left\| \binom{b}{r} \binom{a}{n-r} \right\| \right)$$

$$\text{v.} \leq \frac{M^{2A}}{n} + 2 \left( \sum_r^{[0:\lceil b \rceil]} A! \frac{2M^A}{n} + \sum_r^{[\lceil n-a \rceil : n+1]} \frac{2M^A}{n} A! \right)$$

$$\text{vi.} \leq \frac{M^A}{n} (M^A + 8A!A)$$

$$\text{vii.} = \frac{B}{n}.$$

5. **Yield the tuple**  $\langle B, N, p \rangle$ .

### Procedure III:63(thu2507190646)

#### Objective

Choose a rational number  $1 > X \geq 0$ . The objective of the following instructions is to construct rational numbers  $B > 0$ ,  $N > 0$ , and a procedure  $p(x, a, b, n)$  to show that  $(1+x)_n^{a+b} \equiv (1+x)_n^a (1+x)_n^b$  (err  $\frac{B}{n}$ ) when a complex number  $x$ , two positive rational numbers  $a, b$ , and a positive integer  $n$  such that  $\|x\|^2 \leq 1$ ,  $\text{re}(x) \geq -X$ ,  $a < 1$ ,  $b < 1$ , and  $n > N$  are chosen.

#### Implementation

1. Execute **procedure III:62** on  $\langle 1 \rangle$  and let  $\langle M, N, q \rangle$  receive.
2. Let  $B = \frac{2M}{1-X}$ .
3. Let  $p(x, a, b, n)$  be the following procedure:
  - (a) For  $r \in [1 : n]$ , for  $k \in [0 : r]$ , show that  $\binom{a}{k+1+n-r} (-1)^{k+1} - \binom{a}{k+n-r} (-1)^k$ 
    - i.  $= (-1)^{k+1} \left( \binom{a}{k+1+n-r} + \binom{a}{k+n-r} \right)$

- ii.  $= (-1)^{k+1} \binom{a+1}{k+1+n-r}$
- iii.  $= (-1)^{-(k+1)} \binom{a+1}{k+1+n-r} \|(-1)^{k+1+n-r}$
- iv.  $= \binom{a+1}{k+1+n-r} \|(-1)^{n-r}$ .

(b) Now show that  $\sum_r^{[0:n+1]} \binom{b}{r} \binom{a}{n-r} \| \leq \frac{M}{n}$  using procedure  $q$ .

(c) Show that  $\|x+1\|^2 = \text{re}(x+1)^2 + \text{im}(x)^2 \geq (1-X)^2$ .

(d) Hence using **procedure III:51**, show that  $(1+x)_n^{a+b} \equiv (1+x)_n^a (1+x)_n^b$

- i.  $(\text{err } (1+x)_n^a (1+x)_n^b - (1+x)_n^{a+b})$
- ii.  $(\text{err } \sum_k^{[1:n]} \sum_r^{[k:n]} \binom{a}{k+n-1-r} \binom{b}{r} x^{k+n-1})$
- iii.  $(\text{err } x^n \sum_r^{[1:n]} \binom{b}{r} \sum_k^{[0:r]} \binom{a}{k+n-r} x^k)$
- iv.  $(\text{err } x^n \sum_r^{[1:n]} \binom{b}{r} \sum_k^{[0:r]} ((\binom{a}{k+1+n-r} (-1)^{k+1} \cdot \frac{(-x)^{k+1}}{-x-1} - \binom{a}{k+n-r} (-1)^k \cdot \frac{(-x)^k}{-x-1} - \frac{(-x)^{k+1}}{-x-1} ((\binom{a}{k+1+n-r} (-1)^{k+1} - \binom{a}{k+n-r} (-1)^k)))$
- v.  $(\text{err } \frac{x^n}{x+1} \sum_r^{[1:n]} \binom{b}{r} ((\binom{a}{n-r} x^r - \binom{a}{n-r}) - \sum_k^{[0:r]} (-x)^{k+1} ((\binom{a}{k+1+n-r} (-1)^{k+1} - \binom{a}{k+n-r} (-1)^k)))$
- vi.  $(\text{err } \frac{1}{1-X} \sum_r^{[1:n]} \binom{b}{r} (\| \binom{a}{n} \| + \| \binom{a}{n-r} \| + \sum_k^{[0:r]} \binom{a}{k+1+n-r} (-1)^{k+1} - \binom{a}{k+n-r} (-1)^k))$
- vii.  $(\text{err } \frac{1}{1-X} \sum_r^{[1:n]} \binom{b}{r} (\| \binom{a}{n} \| + \| \binom{a}{n-r} \| + \| \sum_k^{[0:r]} ((\binom{a}{k+1+n-r} (-1)^{k+1} - \binom{a}{k+n-r} (-1)^k)))$
- viii.  $(\text{err } \frac{1}{1-X} \sum_r^{[1:n]} \binom{b}{r} (\| \binom{a}{n} \| + \| \binom{a}{n-r} \| + \| \binom{a}{n} (-1)^r - \binom{a}{n-r} ))$
- ix.  $(\text{err } \frac{2}{1-X} \sum_r^{[1:n]} \binom{b}{r} \| \binom{a}{n-r} \|)$
- x.  $(\text{err } \frac{B}{n})$ .

4. Yield the tuple  $\langle B, N, p \rangle$ .

## Procedure III:64(thu2507191017)

### Objective

Choose a rational number  $0 \leq X < 1$ . The objective of the following instructions is to construct a positive rational number  $D$  such that  $D > 1$ , and a

procedure  $p(x, n, a, k)$  to show that  $\|((1+x)_n^a)^k\|^2 < D^2$  when a complex number  $x$ , a rational number  $a$ , and positive integers  $n, k$  such that  $\|x\|^2 \leq 1$ ,  $\text{re}(x) \geq -X$ , and  $(ka)^2 < 1$  are chosen.

### Implementation

1. Execute **procedure III:34** on  $\langle \frac{2}{1-X} \rangle$  and let  $\langle E, N, q \rangle$  receive.

2. Let  $D = \max(E, (1 + \frac{2}{1-X})^{\lfloor N \rfloor})$ .

3. Let  $p(x, n, a, k)$  be the following procedure:

(a) For  $t \in [1 : n]$ , show that  $\binom{a}{t+1} (-1)^{t+1} - \binom{a}{t} (-1)^t$

$$\text{i.} = (-1)^{t+1} (\binom{a}{t+1} + \binom{a}{t})$$

$$\text{ii.} = (-1)^{t+1} \cdot \frac{(a+1)^{t+1}}{(t+1)!}$$

$$\text{iii.} > 0.$$

(b) Hence show that  $\|k \sum_t^{[1:n]} \binom{a}{t} x^t\|^2$

$$\text{i.} = \|k \sum_t^{[1:n]} ((\binom{a}{t+1} (-1)^{t+1} \cdot \frac{(-x)^{t+1}}{-x-1} - \binom{a}{t} (-1)^t \cdot \frac{(-x)^t}{-x-1} - \frac{(-x)^{t+1}}{-x-1} ((\binom{a}{t+1} (-1)^{t+1} - \binom{a}{t} (-1)^t)))\|^2$$

$$\text{ii.} = \frac{k^2}{\|x+1\|^2} \| \binom{a}{n} x^n - \binom{a}{1} x^1 - \sum_t^{[1:n]} (-x)^{t+1} ((\binom{a}{t+1} (-1)^{t+1} - \binom{a}{t} (-1)^t)) \|^2$$

$$\text{iii.} \leq \frac{k^2}{(\text{re}(x)+1)^2 + \text{im}(x)^2} (|\binom{a}{n}| + a + \sum_t^{[1:n]} |(\binom{a}{t+1} (-1)^{t+1} - \binom{a}{t} (-1)^t)|^2)$$

$$\text{iv.} \leq \frac{k^2}{(1-X)^2} (|\binom{a}{n}| + a + \sum_t^{[1:n]} |(\binom{a}{t+1} (-1)^{t+1} - \binom{a}{t} (-1)^t)|^2)$$

$$\text{v.} = \frac{k^2}{(1-X)^2} (|\binom{a}{n}| + a + \binom{a}{n} (-1)^n - \binom{a}{1} (-1)^1)^2$$

$$\text{vi.} = \frac{k^2}{(1-X)^2} (|\binom{a}{n}| + a - |\binom{a}{n}| + a)^2$$

$$\text{vii.} = (\frac{2ak}{1-X})^2$$

$$\text{viii.} \leq (\frac{2}{1-X})^2.$$

(c) If  $k > N$ , then do the following:

i. Using procedure  $q$ , show that  $\|((1+x)_n^a)^k\|^2$

$$\text{A.} = \|(\sum_t^{[0:n]} \binom{a}{t} x^t)^k\|^2$$

$$\text{B.} = \|(1 + \sum_t^{[1:n]} \binom{a}{t} x^t)^k\|^2$$

$$\text{C.} = \|\exp_k(k \sum_t^{[1:n]} \binom{a}{t} x^t)\|^2$$

$$D. \leq E^2.$$

$$E. \leq D^2.$$

(d) Otherwise do the following:

i. Show that  $\|\sum_t^{[1:n]} \binom{a}{t} x^t\|^2$

$$A. \leq \|k \sum_t^{[1:n]} \binom{a}{t} x^t\|^2$$

$$B. \leq (\frac{2}{1-X})^2.$$

ii. Hence show that  $\|((1+x)_n^a)^k\|^2$

$$A. = (\|(1+x)_n^a\|^2)^k$$

$$B. = (\|1 + \sum_t^{[1:n]} \binom{a}{t} x^t\|^2)^k$$

$$C. \leq (1 + \frac{2}{1-X})^{2k}$$

$$D. \leq D^2.$$

4. **Yield the tuple**  $\langle D, p \rangle$ .

### Procedure III:65(thu2507190752)

#### Objective

Choose a rational number  $0 \leq X < 1$ . The objective of the following instructions is to construct positive rational numbers  $G, N$  and a procedure  $p(x, n, a, k)$  to show that  $(1+x)_n^{ka} \equiv ((1+x)_n^a)^k$  (err  $\frac{Gk}{n}$ ) when positive integers  $n, k$ , a rational number  $a$ , and a complex number  $x$  such that  $\|x\|^2 \leq 1$ ,  $\text{re}(x) \geq -X$ ,  $k > 1$ ,  $0 < ka \leq 1$ , and  $n > N$  are chosen.

#### Implementation

1. Execute **procedure III:64** on  $\langle X \rangle$  and let  $\langle D, t \rangle$  receive.
2. Execute **procedure III:63** on  $\langle X \rangle$  and let  $\langle B, N, q \rangle$  receive.
3. Let  $G = DB$ .
4. Let  $p(x, n, a, k)$  be the following procedure:
  - (a) Using procedures  $t, q$ , show that  $(1+x)_n^{ka}$ 
    - i.  $= ((1+x)_n^a)^0 (1+x)_n^{ka}$
    - ii.  $\equiv ((1+x)_n^a)^1 (1+x)_n^{(k-1)a}$  (err  $D\frac{B}{n}$ )
    - iii.  $\equiv ((1+x)_n^a)^2 (1+x)_n^{(k-2)a}$  (err  $D\frac{B}{n}$ )
    - iv.  $\vdots$

$$v. \equiv ((1+x)_n^a)^k (1+x)_n^{(k-k)a} \text{ (err } D\frac{B}{n})$$

$$vi. = ((1+x)_n^a)^k.$$

(b) **Hence show that**  $(1+x)_n^{ka} \equiv ((1+x)_n^a)^k$  (err  $\frac{DBk}{n}$ ) (err  $\frac{Gk}{n}$ ).

5. **Yield the tuple**  $\langle G, D, N, p \rangle$ .

### Procedure III:66(fri2607191210)

#### Objective

Choose a rational number  $1 > X \geq 0$ . The objective of the following instructions is to construct positive rational numbers  $a, c$  such that  $b > 1$ , and a procedure  $p(x, n, k)$  to show that  $\exp_n(n((1+x)_k^{\frac{1}{n}} - 1)) \equiv 1+x$  (err  $\frac{an}{k}$ ) when a complex number  $x$ , and positive integers  $n, k$  such that  $\|x\|^2 \leq 1$ ,  $\text{re}(x) \geq -X$ ,  $n > 1$ , and  $k > c$  are chosen.

#### Implementation

1. Execute **procedure III:65** on  $\langle X \rangle$  and let  $\langle a, c, p_1 \rangle$  receive.
2. Let  $p(x, n, k)$  be the following procedure:
  - (a) Using procedure  $p_1$  and **procedure III:49**, show that  $\exp_n(n((1+x)_k^{\frac{1}{n}} - 1))$ 
    - i.  $= (1 + \frac{1}{n}(n((1+x)_k^{\frac{1}{n}} - 1)))^n$
    - ii.  $= ((1+x)_k^{\frac{1}{n}})^n$
    - iii.  $\equiv (1+x)_k^1$  (err  $\frac{an}{k}$ )
    - iv.  $= (1+x)^1$
    - v.  $= 1+x$ .
  - (b) **Hence show that**  $\exp_n(n((1+x)_k^{\frac{1}{n}} - 1)) \equiv 1+x$  (err  $\frac{an}{k}$ ).
3. **Yield the tuple**  $\langle a, c, p \rangle$ .

### Procedure III:67(fri2607191243)

#### Objective

Choose a rational number  $1 > X \geq 0$ . The objective of the following instructions is to construct a rational number  $a > 0$  and a procedure  $p(x, n, k)$  to show

that  $\|n((1+x)_k^{\frac{1}{n}} - 1)\|^2 \leq a^2$  when positive integers  $n, k$ , and a complex number  $x$  such that  $\|x\|^2 \leq 1$  and  $\text{re}(x) \geq -X$  are chosen.

### Implementation

1. Let  $a = \frac{2}{1-X}$ .

2. Let  $p(x, n, k)$  be the following procedure:

- (a) Show that  $\|n((1+x)_k^{\frac{1}{n}} - 1)\|^2$ 
  - i.  $= \|n(\sum_r^{[0:k]} (\frac{1}{r}) x^r - 1)\|^2$
  - ii.  $= \|n \sum_r^{[1:k]} (\frac{1}{r}) (-1)^r (-x)^r\|^2$
  - iii.  $= n^2 \|\sum_r^{[1:k]} ((\frac{1}{r+1}) (-1)^{r+1} \cdot \frac{(-x)^{r+1}}{-x-1} - (\frac{1}{r}) (-1)^r \cdot \frac{(-x)^r}{-x-1} - ((\frac{1}{r+1}) (-1)^{r+1} - (\frac{1}{r}) (-1)^r) \frac{(-x)^{r+1}}{-x-1}\|^2$
  - iv.  $= \frac{n^2}{\|x+1\|^2} \|(\frac{1}{k}) x^k - (\frac{1}{1}) x^1 - \sum_r^{[1:k]} ((\frac{1}{r+1}) (-1)^{r+1} - (\frac{1}{r}) (-1)^r) (-x)^{r+1}\|^2$
  - v.  $\leq \frac{n^2}{\|x+1\|^2} \|(\frac{1}{k}) (-1)^{k-1} + \frac{1}{n} + \sum_r^{[1:k]} ((\frac{1}{r+1}) (-1)^{r+1} - (\frac{1}{r}) (-1)^r)\|^2$
  - vi.  $= \frac{n^2}{(\text{re}(x)+1)^2 + \text{im}(x)^2} \|(\frac{1}{k}) (-1)^{k-1} + \frac{1}{n} + (\frac{1}{k}) (-1)^k - (\frac{1}{1}) (-1)^1\|^2$
  - vii.  $\leq \frac{n^2}{(1-X)^2} (\frac{2}{n})^2$
  - viii.  $= a^2$ .

3. Yield the tuple  $\langle a, p \rangle$ .

### Declaration III:17

The notation  $\omega(r)$  will be used as a shorthand notation for  $\frac{1}{r}(1 - \prod_t^{[1:r]}(1 - \frac{1}{nt}))$ .

### Procedure III:68(thu0108191318)

#### Objective

Choose two positive integers  $r, n$  such that  $r > 1$ . The objective of the following instructions is to show that  $\frac{\omega(r+1)}{\omega(r)} \leq 1$ .

### Implementation

1. Using [procedure II:32](#), show that  $\frac{\omega(r+1)}{\omega(r)}$

- (a)  $= \frac{\frac{1}{r+1}(1 - \prod_t^{[1:r+1]}(1 - \frac{1}{nt}))}{\frac{1}{r}(1 - \prod_t^{[1:r]}(1 - \frac{1}{nt}))}$
- (b)  $= \frac{r}{r+1} \cdot \frac{1 - (1 - \frac{1}{nr}) \prod_t^{[1:r]}(1 - \frac{1}{nt})}{1 - \prod_t^{[1:r]}(1 - \frac{1}{nt})}$
- (c)  $= \frac{r}{r+1} \left( 1 + \frac{\frac{1}{nr} \prod_t^{[1:r]}(1 - \frac{1}{nt})}{1 - \prod_t^{[1:r]}(1 - \frac{1}{nt})} \right)$
- (d)  $= \frac{r}{r+1} \left( 1 + \frac{\frac{1}{nr}}{(\prod_t^{[1:r]}(1 - \frac{1}{nt}))^{-1} - 1} \right)$
- (e)  $\leq \frac{r}{r+1} \left( 1 + \frac{\frac{1}{nr}}{(1 - \frac{1}{n(r-1)})^{-(r-1)} - 1} \right)$
- (f)  $= \frac{r}{r+1} \left( 1 + \frac{\frac{1}{nr}}{(1 + \frac{1}{nr - n - 1})^{r-1} - 1} \right)$
- (g)  $\leq \frac{r}{r+1} \left( 1 + \frac{\frac{1}{nr}}{(1 + \frac{1}{n(r-1)})^{r-1} - 1} \right)$
- (h)  $\leq \frac{r}{r+1} \left( 1 + \frac{\frac{1}{nr}}{1 + \frac{r-1}{n(r-1)} - 1} \right)$
- (i)  $= \frac{r}{r+1} (1 + \frac{1}{r})$
- (j)  $= 1$ .

### Declaration III:18

The notation  $\ln_k(1+x)$  will be used as a shorthand for  $\sum_r^{[1:k]} \frac{(-1)^{r-1}}{r} x^r$ .

### Procedure III:69(fri2607191450)

#### Objective

Choose a rational number  $1 > X \geq 0$ . The objective of the following instructions is to construct a positive rational number  $a$  and a procedure  $p(x, n, k)$  to show that  $\ln_k(1+x) \equiv n((1+x)_k^{\frac{1}{n}} - 1)$  (err  $\frac{a}{n}$ ) when positive integers  $n, k$  and a complex number  $x$  such that  $\|x\|^2 \leq 1$  and  $\text{re}(x) \geq -X$  are chosen.

### Implementation

1. Let  $a = \frac{1}{1-X}$ .
2. Let  $p(x, n, k)$  be the following procedure:
  - (a) For  $r \in [2 : k]$ , show that  $\frac{\omega(r+1)}{\omega(r)} \leq 1$  using [procedure III:68](#).
  - (b) Also show that  $\|x + 1\|^2 \geq \text{re}(x + 1)^2 + \text{im}(x)^2 \geq (1 - X)^2$ .
  - (c) Hence show that  $\ln_k(1+x) \equiv n((1+x)^{\frac{1}{k}} - 1)$ 
    - i. (err  $\ln_k(1+x) - n((1+x)^{\frac{1}{k}} - 1)$ )
    - ii. (err  $\sum_r^{[1:k]} \frac{(-1)^{r-1}}{r} x^r - n(\sum_r^{[0:k]} (\frac{1}{r}) x^r - 1)$ )
    - iii. (err  $\sum_r^{[1:k]} \frac{(-1)^{r-1}}{r} x^r - n \sum_r^{[1:k]} (\frac{1}{r}) x^r$ )
    - iv. (err  $\sum_r^{[1:k]} \frac{(-1)^{r-1}}{r!} x^r - \sum_r^{[1:k]} \frac{(\frac{1}{n}-1)^{r-1}}{r!} x^r$ )
    - v. (err  $\sum_r^{[1:k]} \frac{1}{r!} ((-1)^{r-1} - (\frac{1}{n}-1)^{r-1}) x^r$ )
    - vi. (err  $\sum_r^{[1:k]} \frac{(-1)^{r-1}}{r!} (1 - (\frac{1}{n}-1)^{r-1}) x^r$ )
    - vii. (err  $\sum_r^{[1:k]} \frac{(-1)^{r-1}}{r} (1 - \prod_t^{[1:r]} \frac{1}{n-t}) x^r$ )
    - viii. (err  $\sum_r^{[1:k]} \omega(r)(-x)^r$ )
    - ix. (err  $\sum_r^{[1:k]} (\omega(r+1) \cdot \frac{(-x)^{r+1}}{-x-1} - \omega(r) \cdot \frac{(-x)^r}{-x-1} - (\omega(r+1) - \omega(r)) \cdot \frac{(-x)^{r+1}}{-x-1})$ )
    - x. (err  $\frac{1}{x+1} (\omega(k)(-x)^k - \omega(1)(-x)^1 - \sum_r^{[1:k]} (\omega(r+1) - \omega(r))(-x)^{r+1})$ )
    - xi. (err  $\frac{1}{x+1} (\omega(k) + \omega(1) + \sum_r^{[2:k]} (\omega(r) - \omega(r+1)) + \omega(2) - \omega(1))$ )
    - xii. (err  $\frac{1}{1-X} (\omega(k) - \omega(k) + \omega(2) + \omega(2) + \omega(1) - \omega(1))$ )
    - xiii. (err  $\frac{a}{n}$ ).
3. Yield the tuple  $\langle a, p \rangle$ .

### Procedure III:70(fri2607191736)

#### Objective

Choose a rational number  $1 > X \geq 0$ . The objective of the following instructions is to construct a rational number  $a > 0$  and a procedure  $p(x, k)$  to show that  $\|\ln_k(1+x)\|^2 \leq a^2$  when a positive integer  $k$

and a complex number  $x$  such that  $\|x\|^2 \leq 1$  and  $\text{re}(x) \geq -X$  are chosen.

### Implementation

1. Let  $a = \frac{2}{1-X}$ .
2. Let  $p(x, k)$  be the following procedure:
  - (a) Show that  $\|\ln_k(1+x)\|^2$ 
    - i.  $= \|\sum_r^{[1:k]} \frac{(-1)^{r-1}}{r} x^r\|^2$
    - ii.  $= \|\sum_r^{[1:k]} \frac{1}{r} (-x)^r\|^2$
    - iii.  $= \|\sum_r^{[1:k]} (\frac{1}{r+1} \cdot \frac{(-x)^{r+1}}{-x-1} - \frac{1}{r} \cdot \frac{(-x)^r}{-x-1} - (\frac{1}{r+1} - \frac{1}{r}) \cdot \frac{(-x)^{r+1}}{-x-1})\|^2$
    - iv.  $= \frac{1}{\|x+1\|^2} \|\frac{1}{k} (-x)^k - \frac{1}{1} (-x)^1 - \sum_r^{[1:k]} (\frac{1}{r+1} - \frac{1}{r}) (-x)^{r+1}\|^2$
    - v.  $\leq \frac{1}{\|x+1\|^2} (\frac{1}{k} + 1 + \sum_r^{[1:k]} (\frac{1}{r} - \frac{1}{r+1}))^2$
    - vi.  $= \frac{1}{\|x+1\|^2} (\frac{1}{k} + 1 - \frac{1}{k} + 1)^2$
    - vii.  $= \frac{4}{(\text{re}(x)+1)^2 + \text{im}(x)^2}$
    - viii.  $\leq a^2$
3. Yield the tuple  $\langle a, p \rangle$ .

### Procedure III:71(fri2607191801)

#### Objective

Choose a rational number  $1 > X \geq 0$ . The objective of the following instructions is to construct positive rational numbers  $a, c, d, e$  such that  $b > 1$ , and a procedure  $p(x, n, k)$  to show that  $\exp_n(\ln_k(1+x)) \equiv 1+x$  (err  $\frac{an}{k} + \frac{c}{n}$ ) when positive integers  $n, k$ , and a complex number  $x$  such that  $\|x\|^2 \leq 1$ ,  $\text{re}(x) \geq -X$ ,  $k > d$ , and  $n > e$  are chosen.

### Implementation

1. Execute [procedure III:67](#) on  $\langle X \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Execute [procedure III:70](#) on  $\langle X \rangle$  and let  $\langle a_2, p_2 \rangle$  receive.
3. Execute [procedure III:39](#) on  $\langle \max(a_1, a_2) \rangle$  and let  $\langle a_3, e, p_3 \rangle$  receive.



4. Execute **procedure III:69** on  $\langle X \rangle$  and let  $\langle a_4, p_4 \rangle$  receive.
5. Execute **procedure III:66** on  $\langle X \rangle$  and let  $\langle a, d, p_5 \rangle$  receive.
6. Let  $c = a_4 a_3$ .
7. Let  $p(x, n, k)$  be the following procedure:
  - (a) Show that  $\|n((1+x)_k^{\frac{1}{n}} - 1)\|^2 \leq a_1^2$  using procedure  $p_1$ .
  - (b) Show that  $\|\ln_k(1+x)\|^2 \leq a_2^2$  using procedure  $p_2$ .
  - (c) Show that  $\|\ln_k(1+x) - n((1+x)_k^{\frac{1}{n}} - 1)\|^2 \leq (\frac{a_4}{n})^2$  using procedure  $p_4$ .
  - (d) Now using procedures  $p_3, p_5$ , show that  $\exp_n(\ln_k(1+x))$ 
    - i.  $\equiv \exp_n(n((1+x)_k^{\frac{1}{n}} - 1))$ 
      - A. (err  $a_3(\ln_k(1+x) - n((1+x)_k^{\frac{1}{n}} - 1))$ )
      - B. (err  $a_3 \cdot \frac{a_4}{n}$ )
    - ii.  $\equiv 1 + x$  (err  $\frac{an}{k}$ )
  - (e) **Hence show that**  $\exp_n(\ln_k(1+x)) \equiv 1 + x$  (err  $\frac{a_3 a_4}{n} + \frac{an}{k}$ ) (err  $\frac{c}{n} + \frac{an}{k}$ ).
8. Yield the tuple  $\langle a, c, d, e, p \rangle$ .

# Chapter 11

## Gregory-Leibniz Series

### Declaration III:19

The notation  $\tau_n$ , where  $n$  is a positive integer, will be used as a shorthand for  $8 \operatorname{im}(\ln_n(1+i))$ .

### Procedure III:72(3.47)

#### Objective

Choose a positive integer  $k$ . The objective of the following instructions is to show that  $\tau_k = 8 \sum_r^{[0: \lfloor \frac{k}{2} \rfloor]} \frac{(-1)^r}{2r+1}$ .

#### Implementation

1. Using **declaration III:19**, show that  $\tau_k$

$$(a) = 8 \operatorname{im}(\sum_r^{[1:k]} \frac{(-1)^{r-1}}{r} i^r)$$

$$(b) = 8 \operatorname{im}(\sum_r^{[0: \lfloor \frac{k}{2} \rfloor]} \frac{(-1)^{2r}}{2r+1} i^{2r+1})$$

$$(c) = 8 \sum_r^{[0: \lfloor \frac{k}{2} \rfloor]} \frac{i^{2r}}{2r+1}$$

$$(d) = 8 \sum_r^{[0: \lfloor \frac{k}{2} \rfloor]} \frac{(-1)^r}{2r+1}.$$

### Procedure III:73(3.49)

#### Objective

The objective of the following instructions is to construct positive rational numbers  $a, b$  such that  $a \geq 4$ , and a procedure,  $p(n)$ , to show that  $\tau_n \geq a$  when a positive integer  $n \geq b$  is chosen.

#### Implementation

1. Let  $a = \frac{16}{3}$ .

2. **Show that**  $a \geq 4$ .

3. Let  $b = 4$ .

4. Let  $p(n)$  be the following procedure:

(a) Let  $d = n \operatorname{div} 4$ .

(b) Let  $g = n \bmod 4$ .

(c) Hence show that  $n = 4d + g$ .

(d) If  $g = 0$  or  $g = 1$ , then do the following:

i. Using **procedure III:72**, show that  $\tau_n$

$$A. = 8 \sum_r^{[0: \lfloor \frac{4d+g}{2} \rfloor]} \frac{(-1)^r}{2r+1}$$

$$B. = 8 \sum_r^{[0:2d]} \frac{(-1)^r}{2r+1}$$

$$C. = 8(1 - \frac{1}{3} + \sum_r^{[2:2d]} \frac{(-1)^r}{2r+1})$$

$$D. = \frac{16}{3} + 8 \sum_r^{[1:d]} (\frac{1}{4r+1} - \frac{1}{4r+3})$$

$$E. \geq \frac{16}{3}.$$

(e) Otherwise do the following:

i. Show that  $g = 2$  or  $g = 3$ .

ii. Hence show that  $\tau_n$

$$A. = 8 \sum_r^{[0: \lfloor \frac{4d+g}{2} \rfloor]} \frac{(-1)^r}{2r+1}$$

$$B. = 8 \sum_r^{[0:2d+1]} \frac{(-1)^r}{2r+1}$$

$$C. = 8(1 - \frac{1}{3} + \sum_r^{[0:2d]} \frac{(-1)^r}{2r+1} + \frac{(-1)^{2d}}{4d+1})$$

$$D. \frac{16}{3} + 8 \sum_r^{[1:d]} (\frac{1}{4r+1} - \frac{1}{4r+3}) + \frac{8}{4d+1}$$

$$E. \geq \frac{16}{3}.$$

5. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:74(3.50)

#### Objective

The objective of the following instructions is to construct rational numbers  $a, b$  such that  $a \geq 4$  and  $a^2 < 48$ , and a procedure,  $p(n)$ , to show that  $\tau_n \leq a$  when a positive integer  $n$  such that  $n \geq b$  is chosen.

#### Implementation

1. Let  $a = \frac{2104}{315}$ .
2. Show that  $a \geq 4$ .
3. Show that  $a^2 = \frac{4426816}{99225} < 48$ .
4. Let  $b = 10$ .
5. Let  $p(n)$  be the following procedure:
  - (a) Let  $d = n \div 4$ .
  - (b) Let  $g = n \bmod 4$ .
  - (c) Hence verify that  $n = 4d + g$ .
  - (d) If  $g = 0$  or  $g = 1$ , then do the following:
    - i. Show that  $\tau_n$ 
      - A.  $= 8 \sum_r^{[0: \lfloor \frac{n}{2} \rfloor]} \frac{(-1)^r}{2r+1}$
      - B.  $= 8 \sum_r^{[0:5]} \frac{(-1)^r}{2r+1} + 8 \sum_r^{[5: \lfloor \frac{4d+g}{2} \rfloor]} \frac{(-1)^r}{2r+1}$
      - C.  $= a + 8 \sum_r^{[5:2d]} \frac{(-1)^r}{2r+1}$
      - D.  $= a + 8 \sum_r^{[5:2d-1]} \frac{(-1)^r}{2r+1} + \frac{8(-1)^{2d-1}}{4d-1}$
      - E.  $= a - 8 \sum_r^{[3:d]} (\frac{1}{4r-1} - \frac{1}{4r+1}) - \frac{8}{4d-1}$
      - F.  $\leq a$ .
  - (e) Otherwise do the following:
    - i. Show that  $g = 2$  or  $g = 3$ .
    - ii. Hence show that  $\tau_n$ 
      - A.  $= 8 \sum_r^{[0: \lfloor \frac{n}{2} \rfloor]} \frac{(-1)^r}{2r+1}$
      - B.  $= 8 \sum_r^{[0:5]} \frac{(-1)^r}{2r+1} + 8 \sum_r^{[5: \lfloor \frac{4d+g}{2} \rfloor]} \frac{(-1)^r}{2r+1}$
      - C.  $= a + 8 \sum_r^{[5:2d+1]} \frac{(-1)^r}{2r+1}$

$$D. = a - 8 \sum_r^{[2:d]} (\frac{1}{4r+3} - \frac{1}{4r+5})$$

$$E. \leq a.$$

6. Yield the tuple  $\langle a, b, p \rangle$ .

### Procedure III:75(3.53)

#### Objective

The objective of the following instructions is to construct positive rational numbers  $a, c, d, e$ , and a procedure  $p(n, k)$  to show that  $\exp_n(\frac{1}{4}\tau_k i) \equiv i$  (err  $\frac{an}{k} + \frac{c}{n}$ ) when integers  $k, n$  such that  $n > e$  and  $k > d$  are chosen.

#### Implementation

1. Execute **procedure III:70** on  $\langle 0 \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Execute **procedure III:37** on  $\langle a_1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:35** on  $\langle a_1 \rangle$  and let  $\langle a_3, b_3, p_3 \rangle$  receive.
4. Execute **procedure III:71** on  $\langle 0 \rangle$  and let  $\langle a_4, c_4, d, e_4, p_4 \rangle$  receive.
5. Let  $a = \frac{2a_4}{a_3}$ .
6. Let  $c = \frac{2c_4}{a_3} + a_2$ .
7. Let  $e = \max(b_2, b_3, e_4)$ .
8. Let  $p(n, k)$  be the following procedure:
  - (a) Show that  $\|\ln_k(1+i)\|^2 \leq a_1^2$  using procedure  $p_1$ .
  - (b) Hence using procedures  $p_2, p_3, p_4$ , show that  $\exp_n(\frac{1}{4}\tau_k i)$ 
    - i.  $= \exp_n(2 \operatorname{im}(\ln_k(1+i))i)$
    - ii.  $= \exp_n(\ln_k(1+i) - \overline{\ln_k(1+i)})$
    - iii.  $\equiv \frac{\exp_n(\ln_k(1+i))}{\exp_n(\ln_k(1+i))} \text{ (err } \frac{a_2}{n} \text{)}$
    - iv.  $\equiv \frac{1+i}{\exp_n(\ln_k(1+i))} \text{ (err } \frac{1}{a_3}(\frac{a_4 n}{k} + \frac{c_4}{n}) \text{)}$
    - v.  $= \frac{1+i}{\exp_n(\ln_k(1+i))}$
    - vi.  $\equiv \frac{1+i}{1+i}$ 
      - A. (err  $\frac{(1+i)(\frac{a_4 n}{k} + \frac{c_4}{n})}{\exp_n(\ln_k(1+i)) \cdot 1+i}$ )

B.  $(\text{err } \frac{1}{a_3}(\frac{a_4 n}{k} + \frac{c_4}{n}))$

vii.  $= i$ .

(c) **Hence show that**  $\exp_n(\frac{1}{4}\tau_k i) \equiv i (\text{err } \frac{a_2}{n} + \frac{2}{a_3}(\frac{a_4 n}{k} + \frac{c_4}{n})) (\text{err } \frac{a_n}{k} + \frac{c}{n})$ .

9. **Yield the tuple**  $\langle a, c, d, e, p \rangle$ .

### Procedure III:76(3.54)

#### Objective

The objective of the following instructions is to construct positive rational numbers  $a, c, d, e$  such that  $b > 1$ , and a procedure,  $p(n, k)$ , to show that  $\exp_n(-\frac{1}{4}\tau_k i) \equiv -i (\text{err } \frac{a_n}{k} + \frac{c}{n})$  when integers  $k, n$  such that  $n > e$  and  $k > d$  are chosen.

#### Implementation

Implementation is analogous to that of [procedure III:75](#).

### Procedure III:77(mon2608190753)

#### Objective

Choose a rational number  $X \geq 0$  and an integer  $K \geq 0$ . The objective of the following instructions is to construct a rational number  $a$ , and a procedure  $p(x, y, k)$  to show that  $x^k \equiv y^k (\text{err } a(x - y))$  when two complex numbers  $x, y$  and a non-negative integer  $k$  such that  $\|x\|^2 \leq X^2$ ,  $\|y\|^2 \leq X^2$ , and  $k \leq K$  are chosen.

#### Implementation

1. Let  $a = K \max(1, X)^{K-1}$ .
2. Let  $p(x, y, k)$  be the following procedure:
  - (a) Show that  $x^k \equiv y^k$ 
    - i.  $(\text{err } y^k - x^k)$
    - ii.  $(\text{err } (y - x) \sum_r^{[0:k]} x^r y^{k-1-r})$
    - iii.  $(\text{err } (y - x) \sum_r^{[0:k]} X^{k-1})$
    - iv.  $(\text{err } (y - x) K X^{k-1})$
    - v.  $(\text{err } a(y - x))$

3. **Yield the tuple**  $\langle a, p \rangle$ .

### Procedure III:78(3.55)

#### Objective

Choose an integer  $K \geq 0$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(n, m, k)$ , to show that  $\exp_n(\frac{k}{4}\tau_m i) \equiv i^k (\text{err } \frac{a_n}{m} + \frac{b}{n})$  when a non-negative integer  $k$  and two positive integers  $n, m$  such that  $k \leq K$ ,  $n > c$ , and  $m > d$  are chosen.

#### Implementation

1. Execute [procedure III:74](#) and let  $\langle a_1, d, p_1 \rangle$  receive.
2. Execute [procedure III:38](#) on  $\langle (\frac{a_1}{4})^2 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute [procedure III:75](#) and let  $\langle a_3, b_3, c_3, p_3 \rangle$  receive.
4. Execute [procedure III:34](#) on  $\langle (\frac{a_1}{4})^2 \rangle$  and let  $\langle a_4, b_4, p_4 \rangle$  receive.
5. Execute [procedure III:77](#) on  $\langle \max(1, \frac{a_1}{4}), K \rangle$  and let  $\langle a_5, p_5 \rangle$  receive.
6. Let  $a = a_3 a_5$ .
7. Let  $b = a_2 K + b_3 a_5$ .
8. Let  $c = \max(b_2, b_4, c_3)$ .
9. Let  $p(n, k, m)$  be the following procedure:
  - (a) Show that  $\tau_m \leq a_1$  using procedure  $p_1$ .
  - (b) Hence show that  $\|\frac{1}{4}\tau_m i\|^2 = \|\frac{1}{4}\tau_m\|^2 \leq (\frac{a_1}{4})^2$ .
  - (c) Hence show that  $\|\exp_n(\frac{1}{4}\tau_m i) - i\|^2 \leq (\frac{a_3 n}{m} + \frac{b_3}{n})^2$  using procedure  $p_3$ .
  - (d) Hence show that  $\|\exp_n(\frac{1}{4}\tau_m i)\|^2 \leq a_4$  using procedure  $p_4$ .
  - (e) Hence using procedures  $p_2, p_5$ , show that  $\exp_n(\frac{k}{4}\tau_m i)$ 
    - i.  $\equiv \exp_n(\frac{1}{4}\tau_m i)^k$ 
      - A.  $(\text{err } \frac{a_2 k}{n})$
      - B.  $(\text{err } \frac{a_2 K}{n})$

$$\text{ii.} \equiv i^k$$

$$\text{A. } (\text{err } a_5(\exp_n(\frac{1}{4}\tau_m i) - i))$$

$$\text{B. } (\text{err } a_5(\frac{a_3 n}{m} + \frac{b_3}{n}))$$

$$\text{(f) Hence show that } \exp_n(\frac{k}{4}\tau_m i) \equiv i^k (\text{err } \frac{a_2 K}{n} + a_5(\frac{a_3 n}{m} + \frac{b_3}{n})) (\text{err } \frac{a n}{m} + \frac{b}{n}).$$

10. Yield the tuple  $\langle a, b, c, d, p \rangle$ .

### Procedure III:79(3.56)

#### Objective

Choose an integer  $K \geq 0$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(n, m, k)$ , to show that  $\exp_n(\frac{k}{4}\tau_m i) \equiv i^k (\text{err } \frac{a n}{m} + \frac{b}{n})$  when an integer  $k$  and two positive integers  $n, m$  such that  $|k| \leq K$ ,  $n > c$ , and  $m > d$  are chosen.

#### Implementation

Implementation is an extension of that of [procedure III:78](#) using [procedure III:76](#).

### Procedure III:80(3.57)

#### Objective

Choose an integer  $K \geq 0$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(n, m, k)$ , to show that  $\cos_n(\frac{k}{4}\tau_m) \equiv \frac{i^k + (-i)^k}{2} (\text{err } \frac{a n}{m} + \frac{b}{n})$  when an integer  $k$  and two positive integers  $n, m$  such that  $|k| \leq K$ ,  $n > c$ , and  $m > d$  are chosen.

#### Implementation

1. Execute [procedure III:79](#) on  $\langle K \rangle$  and let  $\langle a, b, c, d, q \rangle$  receive.
2. Let  $p(n, m, k)$  be the following procedure:
  - (a) Using procedure  $q$ , show that  $\cos_n(\frac{k}{4}\tau_m)$ 
    - i.  $= \frac{\exp_n(\frac{k}{4}\tau_m i) + \exp_n(-\frac{k}{4}\tau_m i)}{2}$
    - ii.  $= \frac{\exp_n(\frac{k}{4}\tau_m i)}{2} + \frac{\exp_n(-\frac{k}{4}\tau_m i)}{2}$

$$\text{iii.} \equiv \frac{i^k}{2} + \frac{\exp_n(-\frac{k}{4}\tau_m i)}{2} (\text{err } \frac{1}{2}(\frac{a n}{m} + \frac{b}{n}))$$

$$\text{iv.} \equiv \frac{i^k}{2} + \frac{i^{-k}}{2} (\text{err } \frac{1}{2}(\frac{a n}{m} + \frac{b}{n}))$$

$$\text{v.} = \frac{i^k + i^{-k}}{2}.$$

$$\text{(b) Hence show that } \cos_n(\frac{k}{4}\tau_m) \equiv \frac{i^k + i^{-k}}{2} (\text{err } \frac{a n}{m} + \frac{b}{n}).$$

3. Yield the tuple  $\langle a, b, c, d, p \rangle$ .

### Procedure III:81(3.58)

#### Objective

Choose an integer  $K \geq 0$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(n, m, k)$ , to show that  $\sin_n(\frac{k}{4}\tau_m) \equiv \frac{i^k - (-i)^k}{2i} (\text{err } \frac{a n}{m} + \frac{b}{n})$  when an integer  $k$  and two positive integers  $n, m$  such that  $|k| \leq K$ ,  $n > c$ , and  $m > d$  are chosen.

#### Implementation

Implementation is analogous to that of [procedure III:80](#).

### Procedure III:82(3.59)

#### Objective

Choose two integers  $X \geq 0, K \geq 0$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(x, n, m, k)$ , to show that  $\exp_n(x + \frac{k}{4}\tau_m i) \equiv i^k \exp_n(x) (\text{err } \frac{a n}{m} + \frac{b}{n})$  when an integer  $k$  and two positive integers  $n, m$  such that  $\|x\|^2 \leq X$ ,  $|k| \leq K$ ,  $n > c$ , and  $m > d$  are chosen.

#### Implementation

1. Execute [procedure III:74](#) and let  $\langle a_1, b_1, p_1 \rangle$  receive.
2. Let  $H = \max(X, \frac{K a_1}{4})$ .
3. Execute [procedure III:36](#) on  $\langle H \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
4. Execute [procedure III:34](#) on  $\langle X \rangle$  and let  $\langle a_3, b_3, p_3 \rangle$  receive.

5. Execute **procedure III:79** on  $\langle K \rangle$  and let  $\langle a_4, b_4, c_4, d_4, p_4 \rangle$  receive.
6. Let  $a = a_3 a_4$ .
7. Let  $b = a_2 H^2 + a_3 b_4$ .
8. Let  $c = \max(b_2, b_3, c_4)$ .
9. Let  $d = \max(b_1, d_4)$ .
10. Let  $p(x, n, k, m)$  be the following procedure:
  - (a) Show that  $\tau_m \leq a_1$  using procedure  $p_1$ .
  - (b) Hence show that  $\|\frac{k}{4}\tau_m i\|^2 = (\frac{k\tau_m}{4})^2$ .
  - (c) Hence using procedures  $p_2, p_3, p_4$ , show that  $\exp_n(x + \frac{k}{4}\tau_m i)$ 
    - i.  $\equiv \exp_n(\frac{k}{4}\tau_m i) \exp_n(x)$  (err  $\frac{a_2 H^2}{n}$ )
    - ii.  $\equiv i^k \exp_n(x)$  (err  $a_3(\frac{a_4 n}{m} + \frac{b_4}{n})$ )
  - (d) **Hence show that**  $\exp_n(x + \frac{k}{4}\tau_m i) \equiv i^k \exp_n(x)$  (err  $\frac{a_2 H^2}{n} + a_3(\frac{a_4 n}{m} + \frac{b_4}{n})$ ) (err  $\frac{an}{m} + \frac{b}{n}$ ).
11. **Yield the tuple**  $\langle a, b, c, d, p \rangle$ .

### Procedure III:83(3.89)


#### Objective

Choose a positive integer  $K$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(n, m, k)$ , to show that  $\exp_n(\frac{k}{K}\tau_m i)^K \equiv 1$  (err  $\frac{an}{m} + \frac{b}{n}$ ) when an integer  $k$  and positive integers  $n, m$  such that  $0 \leq k < K$ ,  $n \geq c$ , and  $m > d$  are chosen.

#### Implementation

1. Execute **procedure III:74** and let  $\langle a_1, b_1, p_1 \rangle$  receive.
2. Execute **procedure III:38** on  $\langle K a_1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:79** on  $\langle 4K \rangle$  and let  $\langle a_3, b_3, c_3, d_3, p_3 \rangle$  receive.
4. Let  $a = a_3$ .
5. Let  $b = a_2 K + b_3$ .
6. Let  $c = \max(b_2, c_3)$ .
7. Let  $d = \max(b_1, d_3)$ .
8. Let  $p(n, m, k)$  be the following procedure:
  - (a) Show that  $\tau_m \leq a_1$  using procedure  $p_1$ .
  - (b) Hence show that  $\|K \frac{k}{K}\tau_m i\| = \|k\tau_m\|^2 \leq (K a_1)^2$ .
  - (c) Now using procedures  $p_2, p_3$ , show that  $\exp_n(\frac{k}{K}\tau_m i)^K$ 
    - i.  $\equiv \exp_n(K \frac{k}{K}\tau_m i)$  (err  $\frac{a_2 K}{n}$ )
    - ii.  $\equiv \exp_n(\frac{4k}{4}\tau_m i)$
    - iii.  $\equiv i^{4k}$  (err  $\frac{a_3 n}{m} + \frac{b_3}{n}$ )
  - (d) **Hence show that**  $\exp_n(\frac{k}{K}\tau_m i)^K \equiv i^{4k}$  (err  $\frac{a_2 K}{n} + \frac{a_3 n}{m} + \frac{b_3}{n}$ ) (err  $\frac{an}{m} + \frac{b}{n}$ ).
9. **Yield the tuple**  $\langle a, b, c, d, p \rangle$ .

**Figure III:1**



A plot of the list of complex numbers  $\exp_{30}(\frac{[0:11]}{10}\tau_{100}i)$ . Notice that when measurements are done relative to the complex number 1,  $\exp_{30}(\frac{1}{10}\tau_{100}i)$  is roughly  $\frac{1}{10}$ th of a revolution, and also that each complex number has an angle that is roughly an integral multiple of that of  $\exp_{30}(\frac{1}{10}\tau_{100}i)$ .

### Procedure III:84(3.90)

#### Objective

Choose a two rationals  $M, N$  such that  $0 < M$  and  $N^2 < 12$ . The objective of the following instructions is to construct rational numbers  $a, b$  such that  $a > 0$ , and a procedure,  $p(x, n)$ , to show that  $\|\cos_n(x) - 1\|^2 \geq a^2$  when a rational number  $x$  and a positive integer  $n$  such that  $M \leq |x| \leq N$  and  $n > b$  are chosen.

#### Implementation

1. Let  $a = \frac{M^2}{4}(1 - \frac{N^2}{12})$ .
2. **Show that**  $a > 0$ .
3. Let  $b = 4$ .
4. Let  $p(x, n)$  be the following procedure:
  - (a) Using **procedure III:41**, show that  $(\cos_n(x) - 1)^2$

$$\text{i.} = (\frac{1}{2}((1 + \frac{xi}{n})^n + (1 - \frac{xi}{n})^n - 1))^2$$

$$\text{ii.} = (\frac{1}{2}(\sum_r^{[0:n+1]} \frac{n^r}{r!} (\frac{x}{n})^r i^r + \sum_r^{[0:n+1]} \frac{n^r}{r!} (\frac{x}{n})^r (-i)^r - 1))^2$$

$$\text{iii.} = (\sum_r^{[0:\lfloor \frac{n}{2} \rfloor + 1]} \frac{n^{2r}}{(2r)!} (\frac{x}{n})^{2r} (-1)^r - 1)^2$$

$$\text{iv.} = (\sum_r^{[1:\lfloor \frac{n}{2} \rfloor + 1]} \frac{n^{2r}}{(2r)!} (\frac{x}{n})^{2r} (-1)^r)^2$$

$$\text{v.} = (\sum_r^{[1:\lfloor \frac{n}{2} \rfloor + 1]} (-\frac{n^{4r-2}}{(4r-2)!} (\frac{x}{n})^{4r-2} + \frac{n^{4r}}{(4r)!} (\frac{x}{n})^{4r}) - \frac{n^{2\lfloor \frac{n}{2} \rfloor}}{(2\lfloor \frac{n}{2} \rfloor)!} (\frac{x}{n})^{2\lfloor \frac{n}{2} \rfloor} [\lfloor \frac{n}{2} \rfloor \bmod 2 = 1])^2$$

$$\text{vi.} \geq (\sum_r^{[1:\lfloor \frac{n}{2} \rfloor + 1]} \frac{n^{4r-2}}{(4r-2)!} (\frac{x}{n})^{4r-2} (-1 + \frac{(n-4r+2)^2}{(4r)^2} (\frac{x}{n})^2))^2$$

$$\text{vii.} \geq (\sum_r^{[1:\lfloor \frac{n}{2} \rfloor + 1]} \frac{n^{4r-2}}{(4r-2)!} (\frac{x}{n})^{4r-2} (-1 + \frac{1}{(4r)^2} (x^2))^2$$

$$\text{viii.} \geq (\sum_r^{[1:\lfloor \frac{n}{2} \rfloor + 1]} \frac{n^{4r-2}}{(4r-2)!} (\frac{x}{n})^{4r-2} (-1 + \frac{1}{12} x^2))^2$$

$$\text{ix.} \geq (\sum_r^{[1:\lfloor \frac{n}{2} \rfloor + 1]} \frac{n^{4r-2}}{(4r-2)!} (\frac{x}{n})^{4r-2} (-1 + \frac{N^2}{12}))^2$$

$$\text{x.} \geq (\frac{n^2}{2} (\frac{x}{n})^2 (-1 + \frac{N^2}{12}))^2$$

$$\text{xi.} \geq (\frac{1}{4} x^2 (-1 + \frac{N^2}{12}))^2$$

$$\text{xii.} \geq (\frac{M^2}{4} (-1 + \frac{N^2}{12}))^2$$

$$\text{xiii.} = a^2$$

5. **Yield the tuple**  $\langle a, b, p \rangle$ .

### Procedure III:85(3.60)

#### Objective

Choose a positive integer  $K$ . The objective of the following instructions is to construct rational numbers  $a, b, c$  such that  $a > 0$ , and a procedure,  $p(n, m, k)$ , to show that  $\|\exp_n(\frac{k}{K}\tau_m i) - 1\|^2 \geq a^2$  when an integer  $k$  and positive integers  $n, m$  such that  $0 < |k| \leq \frac{K}{2}$ ,  $n > b$ , and  $m > c$  are chosen.

#### Implementation

1. Execute **procedure III:74** and let  $\langle a_1, c, p_1 \rangle$  receive.
2. Show that  $(\frac{a_1}{2})^2 < 12$ .
3. Execute **procedure III:73** and let  $\langle a_2, p_2 \rangle$  receive.
4. Show that  $a_2 > 0$ .
5. Execute **procedure III:84** on  $\langle \frac{a_2}{K}, \frac{a_1}{2} \rangle$  and let  $\langle a, b, p_3 \rangle$  receive.
6. **Show that**  $a > 0$ .
7. Let  $p(n, m, k)$  be the following procedure:
  - (a) Show that  $\frac{1}{K} \leq \frac{|k|}{K} \leq \frac{1}{2}$  given that  $1 \leq |k| \leq \frac{K}{2}$ .
  - (b) Hence show that  $0 < \frac{a_2}{K} \leq \frac{1}{K}\tau_m \leq \frac{|k|}{K}\tau_m \leq \frac{1}{2}\tau_m \leq \frac{a_1}{2}$  using procedures  $p_1$  and  $p_2$ .
  - (c) Hence show that  $(\cos_n(\frac{k}{K}\tau_m) - 1)^2 \geq a^2$  using procedure  $p_3$ .
  - (d) Using **procedure III:41**, show that
    - i.  $\geq \text{re}(\exp_n(\frac{k}{K}\tau_m i) - 1)^2$
    - ii.  $= (\cos_n(\frac{k}{K}\tau_m) - 1)^2$
    - iii.  $\geq a^2$
8. **Yield the tuple**  $\langle a, b, c, p \rangle$ .

### Procedure III:86(3.61)

#### Objective

Choose a positive integer  $K$ . The objective of the following instructions is to construct rational num-

bers  $a, b, c$  such that  $a > 0$ , and a procedure,  $p(n, m, j, k)$ , to show that  $\|\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2 \geq a^2$  when positive integers  $n, j, k, m$  such that  $-K < j \leq k < K$ ,  $0 < k - j \leq \frac{K}{2}$ ,  $n \geq b$ , and  $m \geq c$  are chosen.

#### Implementation

1. Execute **procedure III:74** and let  $\langle a_1, b_1, p_1 \rangle$  receive.
2. Execute **procedure III:35** on  $\langle a_1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:85** on  $\langle K \rangle$  and let  $\langle a_3, b_3, c_3, p_3 \rangle$  receive.
4. Execute **procedure III:36** on  $\langle a_1 \rangle$  and let  $\langle a_4, b_4, p_4 \rangle$  receive.
5. Let  $a = \frac{1}{2}a_2a_3$ .
6. Let  $b = \max(\frac{2a_4a_1^2}{a_2a_3}, b_3, b_4, b_2)$ .
7. Let  $c = \max(b_1, c_3)$ .
8. Let  $p(n, m, j, k)$  be the following procedure:
  - (a) Show that  $\|\frac{j}{K}\|^2 < 1$ 
    - i. given that  $-1 < \frac{j}{K} < 1$
    - ii. given that  $-K < j < K$ .
  - (b) Hence show that  $\|\frac{j}{K}\tau_m i\|^2 = \|\frac{j}{K}\|^2\|\tau_m\|^2 \leq \|\tau_m\|^2 \leq a_1^2$  using procedure  $p_1$ .
  - (c) Hence show that  $\|\exp_n(\frac{j}{K}\tau_m i)\|^2 \geq a_2^2 > 0$  using procedure  $p_2$ .
  - (d) Hence show that  $\|\exp_n(\frac{k-j}{K}\tau_m i) - 1\|^2 \geq a_3^2 > 0$  using procedure  $p_3$ .
  - (e) Show that  $\|\frac{k-j}{K}\tau_m i\|^2 \leq \|\frac{k-j}{K}\|^2\|\tau_m\|^2 \leq \|\tau_m\|^2 \leq a_1^2$  given that  $0 < \frac{k-j}{K} \leq \frac{1}{2}$ .
  - (f) Show that  $n \geq b \geq \frac{2a_4a_1^2}{a_2a_3}$ .
  - (g) Hence show that  $\|\exp_n(\frac{k-j}{K}\tau_m i) \exp_n(\frac{j}{K}\tau_m i) - \exp_n(\frac{k-j}{K}\tau_m i + \frac{j}{K}\tau_m i)\|^2 \leq \frac{a_4^2 \|\frac{k-j}{K}\tau_m i\|^2 \|\frac{j}{K}\tau_m i\|^2}{n^2} \leq \frac{a_4^2 a_1^4}{n^2} \leq (\frac{a_2a_3}{2})^2$  using procedure  $p_4$ .
  - (h) Hence using **procedure III:19**, show that
    - i.  $= \|\exp_n(\frac{k-j}{K}\tau_m i) + \frac{j}{K}\tau_m i - \exp_n(\frac{k-j}{K}\tau_m i) \exp_n(\frac{j}{K}\tau_m i) + \exp_n(\frac{k-j}{K}\tau_m i) \exp_n(\frac{j}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2$



- ii. =  $\|\exp_n(\frac{j}{K}\tau_m i)(\exp_n(\frac{k-j}{K}\tau_m i) - 1) - (\exp_n(\frac{k-j}{K}\tau_m i) + \frac{j}{K}\tau_m i) - \exp_n(\frac{k-j}{K}\tau_m i)\exp_n(\frac{j}{K}\tau_m i)\|^2$
- iii.  $\geq (a_2 a_3 - \frac{a_2 a_3}{2})^2$
- iv.  $\geq a^2$ .

9. Yield the tuple  $\langle a, b, c, p \rangle$ .

### Procedure III:87(3.62)

#### Objective

Choose a positive integer  $K$ . The objective of the following instructions is to construct rational numbers  $a, b, c$  such that  $a > 0$ , and a procedure,  $p(n, m, j, k)$ , to show that  $\|\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2 \geq a^2$  when positive integers  $n, j, k, m$  such that  $0 \leq j \leq k < K$ ,  $\frac{K}{2} \leq k - j < K$ ,  $n \geq b$ , and  $\frac{m}{n} \geq c$  are chosen.

#### Implementation

1. Execute **procedure III:86** on  $\langle K \rangle$  and let  $\langle a_1, b_1, c_1, p_1 \rangle$  receive.
2. Execute **procedure III:74** and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:82** on  $\langle a_2, 4 \rangle$  and let  $\langle a_3, b_3, c_3, d_3, p_3 \rangle$  receive.
4. Let  $a = \frac{1}{2}a_1$ .
5. Let  $b = \max(\frac{4b_3}{a_1}, b_1, c_3)$ .
6. Let  $c = \max(\frac{4a_3}{a_1}, \frac{c_1}{b}, \frac{b_2}{b}, \frac{d_3}{b})$ .
7. Let  $p(n, m, j, k)$  be the following procedure:
  - (a) Show that  $-\frac{K}{2} \leq k - K < j < \frac{K}{2}$ .
  - (b) Also show that  $0 < j - (k - K) \leq \frac{K}{2}$ .
  - (c) Show that  $m \geq cn \geq \frac{c_1}{b}b = c_1$ .
  - (d) Hence show that  $\|\exp_n(\frac{j}{K}\tau_m i) - \exp_n(\frac{k-K}{K}\tau_m i)\|^2 \geq a_1^2$  using procedure  $p_1$ .
  - (e) Show that  $m \geq cn \geq \frac{b_2}{b}b = b_2$ .
  - (f) Hence show that  $\tau_m \leq a_2$  using procedure  $p_2$ .
  - (g) Hence show that  $\|\frac{k}{K}\tau_m i\|^2 = \|\frac{k}{K}\|^2 \|\tau_m\|^2 \leq \|\tau_m\|^2 \leq a_2^2$ .

(h) Also show that  $m \geq cn \geq \frac{d_3}{b}b = d_3$ .

(i) Hence show that  $\|i^{-4}\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{k}{K}\tau_m i - \frac{4}{4}\tau_m i)\|^2 \leq (\frac{a_3 n}{m} + \frac{b_3}{n})^2 \leq (\frac{a_1}{2})^2$  using procedure  $p_3$ .

(j) Now show that  $\|\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2$

$$i. = \|\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{k}{K}\tau_m i - \tau_m i) + \exp_n(\frac{k-K}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2$$

$$ii. \geq \frac{1}{2}\|\exp_n(\frac{k-K}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2 - \|\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{k}{K}\tau_m i - \tau_m i)\|^2$$

$$iii. \geq \frac{1}{2}a_1^2 - (\frac{a_1}{2})^2$$

$$iv. \geq a^2.$$

8. Yield the tuple  $\langle a, b, c, p \rangle$ .

### Procedure III:88(3.63)

#### Objective

Choose a positive integer  $K$ . The objective of the following instructions is to construct rational numbers  $a, b, c$  such that  $a > 0$ , and a procedure,  $p(n, m, j, k)$ , to show that  $\|\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2 \geq a^2$  when positive integers  $n, j, k, m$  such that  $0 \leq j \leq k < K$ ,  $0 < k - j < K$ ,  $n \geq b$ , and  $\frac{m}{n} \geq c$  are chosen.

#### Implementation

1. Execute **procedure III:86** on  $\langle K \rangle$  and let  $\langle a_1, b_1, c_1, p_1 \rangle$  receive.
2. Execute **procedure III:87** on  $\langle K \rangle$  and let  $\langle a_2, b_2, c_2, p_2 \rangle$  receive.
3. Let  $a = \min(a_1, a_2)$ .
4. **Show that**  $a > 0$ .
5. Let  $b = \max(b_1, b_2)$ .
6. Let  $c = \max(\frac{c_1}{b}, c_2)$ .
7. Let  $p(n, m, j, k)$  be the following procedure:
  - (a) If  $k - j \leq \frac{K}{2}$ , then do the following:
    - i. Show that  $m \geq cn \geq \frac{c_1}{b}b = c_1$ .

- ii. **Hence show that**  $\|\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2 \geq a_1 \geq a$  **using procedure**  $p_1$ .
- (b) Otherwise if  $k - j > \frac{K}{2}$ , then do the following:
  - i. **Show that**  $\|\exp_n(\frac{k}{K}\tau_m i) - \exp_n(\frac{j}{K}\tau_m i)\|^2 \geq a_2 \geq a$  **using procedure**  $p_2$ .
8. **Yield the tuple**  $\langle a, b, c, p \rangle$ .

### Declaration III:20

The phrase "**complex polynomial**" will be used to indicate that the declarations and procedures pertaining to polynomials are being used but with the provision that all uses of rational numbers therein are substituted with uses of complex numbers.

### Procedure III:89(3.64)

#### Objective

Choose a positive integer  $K$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(n, m)$ , to construct a list of complex numbers  $z$  and a list of complex polynomials  $q$  such that,

1.  $z_k = \exp_n(\frac{K-k-1}{K}\tau_m i)$  for  $k \in [0 : K]$
2.  $q_K = \lambda^K - 1$
3.  $q_{K-1} = \sum_r^{[0:K]} \lambda^r$
4.  $q_{k+1} = (\lambda - z_k)q_k + \Lambda(q_{k+1}, z_k)$  for  $k \in [0 : K]$
5.  $(q_k)_{\deg(q_k)} = 1$  for  $k \in [0 : K + 1]$
6.  $\Lambda(q_k, z_j) \equiv 0 \pmod{\frac{an}{m} + \frac{b}{n}}$  for  $j \in [0 : k]$ , for  $k \in [0 : K + 1]$

when two positive integers  $n, m$  such that  $n > c$  and  $\frac{m}{n} > d$  are chosen.

#### Implementation

1. Execute **procedure III:83** on  $\langle K \rangle$  and let  $\langle a_1, b_1, c_1, d_1, p_1 \rangle$  receive.
2. Execute **procedure III:88** on  $\langle K \rangle$  and let  $\langle a_2, b_2, c_2, p_2 \rangle$  receive.
3. Let  $a = \max(1, \frac{2}{a_2})^K a_1$ .

4. Let  $b = \max(1, \frac{2}{a_2})^K b_1$ .
5. Let  $c = \max(c_1, b_2)$ .
6. Let  $d = \max(d_1, c_2)$ .
7. Let  $p(n, m)$  be the following procedure:
  - (a) **Let**  $q_K = \lambda^K - 1$ .
  - (b) For  $k \in [K : 0]$ , do the following:
    - i. **Let**  $z_k = \exp_n(\frac{K-k-1}{K}\tau_m i)$ .
    - ii. **Now show that**  $\|\Lambda(q_K, z_k)\|^2 \leq (\frac{a_1 n}{m} + \frac{b_1}{n})^2$  **using procedure**  $p_1$ .
  - (c) For  $k \in [K : 0]$ , do the following:
    - i. Let  $q_k = q_{k+1} \operatorname{div}(\lambda - z_k)$ .
    - ii. Let  $r_k = q_{k+1} \bmod (\lambda - z_k)$ .
    - iii. Show that  $\deg(r_k) = 0$  given that  $\deg(r_k) < \deg(\lambda - z_k) = 1$ .
    - iv. **Show that**  $1 = (q_{k+1})_{\deg(q_{k+1})} = ((\lambda - z_k)q_k + r_k)_{\deg(q_{k+1})} = (q_k)_{\deg(q_k)}$  **given that**  $q_{k+1} = (\lambda - z_k)q_k + r_k$ .
    - v. **Show that**  $q_{k+1} = (\lambda - z_k)q_k + \Lambda(q_{k+1}, z_k)$  **given that**  $\Lambda(q_{k+1}, z_k) = \Lambda(\lambda - z_k, z_k)\Lambda(q_k, z_k) + \Lambda(r_k, z_k) = (z_k - z_k)\Lambda(q_k, z_k) + r_k = r_k$ .
    - vi. **Execute the subprocedure III:90:0 on**  $\langle k, q_{k+1}, z \rangle$ .
  - (d) Now using (cv), verify that  $(\lambda - 1) \sum_r^{[0:K]} \lambda^r$ 
    - i.  $= q_K$
    - ii.  $= (\lambda - z_{K-1})q_{K-1} + \Lambda(q_K, z_{K-1})$
    - iii.  $= (\lambda - 1)q_{K-1} + \Lambda(\lambda^K - 1, 1)$
    - iv.  $= (\lambda - 1)q_{K-1}$ .
  - (e) **Hence show that**  $\sum_r^{[0:K]} \lambda^r = q_{K-1}$ .
  - (f) **Yield the tuple**  $\langle z, q \rangle$ .
8. **Yield the tuple**  $\langle a, b, c, d, p \rangle$ .

#### Subprocedure III:90:0

**Objective** Choose a non-negative integer  $k$ , a complex polynomial  $q_{k+1}$ , and a list of complex numbers  $z$  such that  $z_j = \exp_n(\frac{j}{K}\tau_m i)$  and  $\Lambda(q_{k+1}, z_j) \equiv 0 \pmod{(\frac{2}{a_2})^{K-(k+1)}(\frac{a_1 n}{m} + \frac{b_1}{n})}$  for  $j \in [k+1 : 0]$ . Let  $q_k = q_{k+1} \operatorname{div}(\lambda - z_k)$ . The objective of the

following instructions is to show that  $\Lambda(q_k, z_j) \equiv 0 \pmod{(\frac{2}{a_2})^{K-k}(\frac{a_1 n}{m} + \frac{b_1}{n})}$  for  $j \in [k : 0]$ .

### Implementation

1. For  $j \in [k : 0]$ , do the following:
  - (a) Show that  $\Lambda(q_{k+1}, z_j) - \Lambda(q_{k+1}, z_k) = (z_j - z_k)\Lambda(q_k, z_j)$  given that  $\Lambda(q_{k+1}, z_j) = \Lambda(\lambda - z_k, z_j)\Lambda(q_k, z_j) + \Lambda(q_{k+1}, z_k)$ .
  - (b) Show that  $\|z_j - z_k\|^2 \geq a_2^2$  using procedure  $p_2(n, m, \min(j, k), \max(j, k))$ .
  - (c) Hence show that  $a_2^2 \|\Lambda(q_k, z_j)\|^2$ 
    - i.  $\leq \|z_j - z_k\|^2 \|\Lambda(q_k, z_j)\|^2$
    - ii.  $= \|(z_j - z_k)\Lambda(q_k, z_j)\|^2$
    - iii.  $= \|\Lambda(q_{k+1}, z_j) - \Lambda(q_{k+1}, z_k)\|^2$
    - iv.  $\leq ((\frac{2}{a_2})^{K-k-1}(\frac{a_1 n}{m} + \frac{b_1}{n}) + (\frac{2}{a_2})^{K-k-1}(\frac{a_1 n}{m} + \frac{b_1}{n}))^2$
    - v.  $= (2(\frac{2}{a_2})^{K-k-1}(\frac{a_1 n}{m} + \frac{b_1}{n}))^2$
    - vi.  $= a_2^2 ((\frac{2}{a_2})^{K-k}(\frac{a_1 n}{m} + \frac{b_1}{n}))^2$ .
  - (d) **Hence show that**  $\|\Lambda(q_k, z_j)\|^2 \leq ((\frac{2}{a_2})^{K-k}(\frac{a_1 n}{m} + \frac{b_1}{n}))^2 \leq (\frac{an}{m} + \frac{b}{n})^2$ .

### Procedure III:90(3.65)

#### Objective

Choose a rational number  $X$  and a positive integer  $K$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(x, n, m)$ , to show that  $\sum_r^{[0:K]} x^r \equiv \prod_r^{[1:K]} (x - \exp_n(\frac{x}{K} \tau_m i)) \pmod{(\frac{an}{m} + \frac{b}{n})}$  when a complex number  $x$  and positive integers  $n, m$  such that  $n > c$ ,  $\frac{m}{n} > d$ , and  $\|x\|^2 \leq X$  are chosen.

### Implementation

1. Execute **procedure III:89** on  $\langle K \rangle$  and let  $\langle a_1, b_1, c_1, d_1, p_1 \rangle$  receive.
2. Execute **procedure III:74** and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:34** on  $\langle a_2 \rangle$  and let  $\langle a_3, b_3, p_3 \rangle$  receive.

$$4. \text{ Let } l = \sum_k^{[0:K-1]} \prod_j^{[k+1:K-1]} (X + a_3).$$

$$5. \text{ Let } a = a_1 l.$$

$$6. \text{ Let } b = b_1 l.$$

$$7. \text{ Let } c = \max(c_1, b_3).$$

$$8. \text{ Let } d = \max(d_1, b_2).$$

9. Let  $p(x, n, m)$  be the following procedure:

(a) Show that  $\tau_m \leq a_2$  using procedure  $p_2$ .

(b) Execute procedure  $p_1$  on  $\langle n, m \rangle$  and let  $\langle z, t \rangle$  receive.

(c) For  $j \in [1 : K]$ , do the following:

$$\text{i. Show that } \|\frac{j}{K} \tau_m i\|^2 = \|\frac{j}{K}\|^2 \|\tau_m\|^2 \leq \|\tau_m\|^2 \leq a_2.$$

$$\text{ii. Hence show that } \|z_j\|^2 = \|\exp_n(\frac{j}{K} \tau_m i)\|^2 \leq a_3 \text{ using procedure } p_3.$$

$$\text{(d) Hence show that } \|\sum_r^{[0:K]} x^r - \prod_r^{[1:K]} (x - z_r)\|^2$$

$$\text{i. } = \|\Lambda(\sum_r^{[0:K]} \lambda^r, x) - \prod_r^{[1:K]} (x - z_r)\|^2$$

$$\text{ii. } = \|\Lambda(t_{K-1}, x) - \prod_r^{[1:K]} (x - z_r)\|^2$$

$$\text{iii. } = \|\Lambda(\prod_j^{[0:K-1]} (\lambda - z'_j) + \sum_k^{[0:K-1]} \Lambda(t_{k+1}, z'_k) \prod_j^{[k+1:K-1]} (\lambda - z'_j), x) - \prod_r^{[1:K]} (x - z_r)\|^2$$

$$\text{iv. } = \|\prod_j^{[0:K-1]} (x - z'_j) + \sum_k^{[0:K-1]} \Lambda(t_{k+1}, z'_k) \prod_j^{[k+1:K-1]} (x - z'_j) - \prod_r^{[1:K]} (x - z_r)\|^2$$

$$\text{v. } = \|\sum_k^{[0:K-1]} \Lambda(t_{k+1}, z'_k) \prod_j^{[k+1:K-1]} (x - z'_j)\|^2$$

$$\text{vi. } \leq (\sum_k^{[0:K-1]} (\frac{a_1 n}{m} + \frac{b_1}{n}) \prod_j^{[k+1:K-1]} (X + a_3))^2$$

$$\text{vii. } = ((\frac{a_1 n}{m} + \frac{b_1}{n}) \sum_k^{[0:K-1]} \prod_j^{[k+1:K-1]} (X + a_3))^2$$

$$\text{viii. } = (\frac{an}{m} + \frac{b}{n})^2.$$

10. **Yield the tuple**  $\langle a, b, c, d, p \rangle$ .

### Procedure III:91(3.66)

#### Objective

Choose a rational number  $X$  and a positive integer  $K$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(x, n, m)$ , to show that  $x^K - 1 \equiv \prod_r^{[0:K]} (x - \exp_n(\frac{r}{K} \tau_m i))$  (err  $\frac{an}{m} + \frac{b}{n}$ ) when a complex number  $x$  and positive integers  $n, m$  such that  $n > c$ ,  $\frac{m}{n} > d$ , and  $\|x\|^2 \leq X$  are chosen.

#### Implementation

1. Execute **procedure III:90** on  $\langle X, K \rangle$  and let  $\langle a_1, b_1, c, d, p_1 \rangle$  receive.
2. Let  $a = (X + 1)a_1$ .
3. Let  $b = (X + 1)b_1$ .
4. Let  $p(x, n, m)$  be the following procedure:
  - (a) Show that  $\|\sum_r^{[0:K]} x^r - \prod_r^{[1:K]} (x - \exp_n(\frac{r}{K} \tau_m i))\|^2 \leq (\frac{a_1 n}{m} + \frac{b_1}{n})^2$  using procedure  $p_1$ .
  - (b) Hence show that  $\|x^K - 1 - \prod_r^{[0:K]} (x - \exp_n(\frac{r}{K} \tau_m i))\|^2$ 
    - i.  $= \|(x - 1) \sum_r^{[0:K]} x^r - (x - 1) \prod_r^{[1:K]} (x - \exp_n(\frac{r}{K} \tau_m i))\|^2$
    - ii.  $= \|x - 1\|^2 \|\sum_r^{[0:K]} x^r - \prod_r^{[1:K]} (x - \exp_n(\frac{r}{K} \tau_m i))\|^2$
    - iii.  $\leq (X + 1)^2 (\frac{a_1 n}{m} + \frac{b_1}{n})^2$
    - iv.  $= (\frac{an}{m} + \frac{b}{n})^2$ .
5. Yield the tuple  $\langle a, b, c, d, p \rangle$ .

### Procedure III:92(3.67)

#### Objective

Choose a rational number  $X$  and a positive integer  $K$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , and a procedure,  $p(x, n, m)$ , to show that  $\exp_K(x) - 1 \equiv x \prod_r^{[1:K]} (1 - \frac{x}{\exp_n(\frac{r}{K} \tau_m i) - 1})$  (err  $\frac{an}{m} + \frac{b}{n}$ ) when a complex number  $x$  and positive integers  $n, m$  such that  $n > c$ ,  $\frac{m}{n} > d$ , and  $\|x\|^2 \leq X$  are chosen.

#### Implementation

1. Execute **procedure III:91** on  $\langle 1 + \frac{X}{K}, K \rangle$  and let  $\langle a_1, b_1, c_1, d_1, p_1 \rangle$  receive.
2. Execute **procedure III:90** on  $\langle 1, K \rangle$  and let  $\langle a_2, b_2, c_2, d_2, p_2 \rangle$  receive.
3. Execute **procedure III:88** on  $\langle K \rangle$  and let  $\langle a_3, b_3, c_3, p_3 \rangle$  receive.
4. Let  $l = \frac{X}{K} (1 + \frac{X}{Ka_3})^{K-1}$ .
5. Let  $a = a_1 + la_2$ .
6. Let  $b = b_1 + lb_2$ .
7. Let  $c = \max(c_1, c_2, b_3)$ .
8. Let  $d = \max(d_1, d_2, c_3)$ .
9. Let  $p(x, n, m)$  be the following procedure:
  - (a) Show that  $\|1 + \frac{x}{K}\|^2 \leq (1 + \frac{X}{K})^2$ .
  - (b) Hence show that  $\|(1 + \frac{x}{K})^K - 1 - \prod_r^{[0:K]} (1 + \frac{x}{K} - \exp_n(\frac{r}{K} \tau_m i))\|^2 \leq (\frac{a_1 n}{m} + \frac{b_1}{n})^2$  using procedure  $p_1$ .
  - (c) Hence show that  $\|K - \prod_r^{[1:K]} (1 - \exp_n(\frac{r}{K} \tau_m i))\|^2 = \sum_r^{[0:K]} 1^r - \prod_r^{[1:K]} (1 - \exp_n(\frac{r}{K} \tau_m i))\|^2 \leq (\frac{a_2 n}{m} + \frac{b_2}{n})^2$  using procedure  $p_2$ .
  - (d) For  $j \in [1 : K]$ , do the following:
    - i. Show that  $\|\exp_n(\frac{j}{K} \tau_m i) - 1\|^2 \geq a_3^2$  using procedure  $p_3$ .
    - ii. Let  $z_j = K(\exp_n(\frac{j}{K} \tau_m i) - 1)$ .
  - (e) Hence show that  $\|\exp_K(x) - 1 - x \prod_r^{[1:K]} (1 - \frac{x}{z_r})\|^2$ 
    - i.  $= \|\exp_K(x) - 1 - \prod_r^{[0:K]} (1 + \frac{x}{K} - \exp_n(\frac{r}{K} \tau_m i)) + \prod_r^{[0:K]} (1 + \frac{x}{K} - \exp_n(\frac{r}{K} \tau_m i)) - x \prod_r^{[1:K]} (1 - \frac{x}{z_r})\|^2$
    - ii.  $= \|\exp_K(x) - 1 - \prod_r^{[0:K]} (1 + \frac{x}{K} - \exp_n(\frac{r}{K} \tau_m i)) + \frac{x}{K} \prod_r^{[1:K]} (1 + \frac{x}{K} - \exp_n(\frac{r}{K} \tau_m i)) - x \prod_r^{[1:K]} (1 - \frac{x}{z_r})\|^2$
    - iii.  $= \|\exp_K(x) - 1 - \prod_r^{[0:K]} (1 + \frac{x}{K} - \exp_n(\frac{r}{K} \tau_m i)) + \frac{x}{K} \prod_r^{[1:K]} (1 - \exp_n(\frac{r}{K} \tau_m i)) \prod_r^{[1:K]} (1 - \frac{x}{z_r}) - x \prod_r^{[1:K]} (1 - \frac{x}{z_r})\|^2$

$$\text{iv.} = \left\| \left( \exp_K(x) - 1 - \prod_r^{[0:K]} \left( 1 + \frac{x}{K} - \exp_n\left(\frac{r}{K} \tau_m i\right) \right) + \frac{x}{K} \prod_r^{[1:K]} \left( 1 - \frac{x}{z_r} \right) \left( \prod_r^{[1:K]} (1 - \exp_n\left(\frac{r}{K} \tau_m i\right)) - K \right) \right\|^2$$

$$\text{v.} \leq \left( \left( \frac{a_1 n}{m} + \frac{b_1}{n} \right) + \frac{X}{K} \left( \prod_r^{[1:K]} \left( 1 + \frac{X}{K a_3} \right) \right) \left( \frac{a_2 n}{m} + \frac{b_2}{n} \right) \right)^2$$

$$\text{vi.} = \left( \left( \frac{a_1 n}{m} + \frac{b_1}{n} \right) + \frac{X}{K} \left( 1 + \frac{X}{K a_3} \right)^{K-1} \left( \frac{a_2 n}{m} + \frac{b_2}{n} \right) \right)^2$$

$$\text{vii.} = \left( \frac{a n}{m} + \frac{b}{n} \right)^2.$$

10. **Yield the tuple**  $\langle a, b, c, d, p \rangle$ .

## Part IV

# Differential Arithmetic

## Chapter 12

# Differential Arithmetic

### Procedure IV:0(tue2008191129)

#### Objective

Choose the following:

1. A procedure  $q_1(x, n)$  to show that  $p_n(x) \equiv 0$  (err  $a_1$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > c_1$  are chosen.
2. A procedure  $q_2(x, n)$  to show that  $t_n(x) \equiv 0$  (err  $a_2$ )<sup>2</sup> when a complex number  $x$  and a positive integer  $n$  such that  $R(x)$  and  $n > c_2$  are chosen.

The objective of the following instructions is to construct the following:

1. Rational numbers  $a_3, b_3$ .
2. A procedure  $q_3(x, n)$  to show that  $p_n(x) + t_n(x) \equiv 0$  (err  $a_3$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x), R(x)$ , and  $n > b_3$  are chosen.

#### Implementation

1. Let  $a_3 = a_1 + a_2$ .
2. Let  $b_3 = \max(c_1, c_2)$ .
3. Let  $q_3(x, n)$  be the following procedure:
  - (a) Show that  $p_n(x) \equiv 0$  (err  $a_1$ ) using procedure  $q_1$ .
  - (b) Show that  $t_n(x) \equiv 0$  (err  $a_2$ ) using procedure  $q_2$ .

- (c) Hence show that  $p_n(x) + t_n(x) \equiv 0$  (err  $a_1 + a_2$ ) (err  $a_3$ ).

4. Yield the tuple  $\langle a_3, b_3, q_3 \rangle$ .

### Procedure IV:1(tue2008191139)

#### Objective

Choose the following:

1. A procedure  $q_1(x, n)$  to show that  $p_n(x) \equiv 0$  (err  $a_1$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > c_1$  are chosen.
2. A procedure  $q_2(x, n)$  to show that  $t_n(x) \equiv 0$  (err  $a_2$ ) when a complex number  $x$  and a positive integer  $n$  such that  $R(x)$  and  $n > c_2$  are chosen.

The objective of the following instructions is to construct the following:

1. Rational numbers  $a_3, b_3$ .
2. A procedure  $q_3(x, n)$  to show that  $p_n(x)t_n(x) \equiv 0$  (err  $a_3$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x), R(x)$ , and  $n > b_3$  are chosen.

#### Implementation

Implementation is analogous to that of **procedure IV:0**.

### Declaration IV:0

The notation  $\{x\}$ , where  $x$  is a complex number, will be used as a shorthand for  $|\operatorname{re}(x)| + |\operatorname{im}(x)|$ .

### Procedure IV:2(tue2008190655)

#### Objective

Choose a complex number  $a$  such that  $\{a\} = 0$ . The objective of the following instructions is to show that  $a = 0$ .

#### Implementation

1. Using **declaration IV:0**, show that  $|\operatorname{re}(a)| + |\operatorname{im}(a)| = 0$ .
2. Hence show that  $\operatorname{re}(a) = 0$ 
  - (a) given that  $|\operatorname{re}(a)| = 0$
  - (b) given that  $0 \geq |\operatorname{re}(a)| \geq 0$
  - (c) given that  $|\operatorname{im}(a)| \geq 0$ .
3. Also show that  $\operatorname{im}(a) = 0$ 
  - (a) given that  $|\operatorname{im}(a)| = 0$
  - (b) given that  $0 \geq |\operatorname{im}(a)| \geq 0$
  - (c) given that  $|\operatorname{re}(a)| \geq 0$ .
4. **Hence show that  $a = 0$ .**

### Procedure IV:3(tue2008190520)

#### Objective

Choose a complex number  $a$  and a rational number  $b$ . The objective of the following instructions is to show that  $\{ba\} = |b|\{a\}$ .

#### Implementation

1. Using **declaration IV:0**, show that  $\{ba\}$ 
  - (a)  $= |\operatorname{re}(ba)| + |\operatorname{im}(ba)|$
  - (b)  $= |b \operatorname{re}(a)| + |b \operatorname{im}(a)|$
  - (c)  $= |b|(|\operatorname{re}(a)| + |\operatorname{im}(a)|)$
  - (d)  $= |b|\{a\}$ .

### Procedure IV:4(tue2008190540)

#### Objective

Choose two complex numbers  $a, b$ . The objective of the following instructions is to show that  $\{a + b\} \leq \{a\} + \{b\}$ .

#### Implementation

1. Using **declaration IV:0**, show that  $\{a + b\}$ 
  - (a)  $= |\operatorname{re}(a + b)| + |\operatorname{im}(a + b)|$
  - (b)  $= |\operatorname{re}(a) + \operatorname{re}(b)| + |\operatorname{im}(a) + \operatorname{im}(b)|$
  - (c)  $\leq |\operatorname{re}(a)| + |\operatorname{re}(b)| + |\operatorname{im}(a)| + |\operatorname{im}(b)|$
  - (d)  $= \{a\} + \{b\}$ .

### Procedure IV:5(tue2008190546)

#### Objective

Choose two complex numbers  $a, b$ . The objective of the following instructions is to show that  $\{ab\} \leq \{a\}\{b\}$ .

#### Implementation

1. Using **procedure IV:4**, show that  $\{ab\}$ 
  - (a)  $= \{(\operatorname{re}(a) + \operatorname{im}(b)i)b\}$
  - (b)  $= \{\operatorname{re}(a)b + \operatorname{im}(a)bi\}$
  - (c)  $\leq \{\operatorname{re}(a)b\} + \{\operatorname{im}(a)b\}$
  - (d)  $= (|\operatorname{re}(a)| + |\operatorname{im}(a)|)\{b\}$
  - (e)  $= \{a\}\{b\}$ .

### Procedure IV:6(tue2008190632)

#### Objective

Choose a complex number  $a$ . The objective of the following instructions is to show that  $\|a\|^2 \leq \{a\}^2$ .



### Implementation

- Using **procedure III:18**, show that  $\|a\|^2$ 
  - $\| \operatorname{re}(a) + \operatorname{im}(a)i \|^2$
  - $\leq (|\operatorname{re}(a)| + |\operatorname{im}(a)|)^2$
  - $= \{a\}^2$ .

### Procedure IV:7(tue2008190639)

#### Objective

Choose a complex number  $a$ . The objective of the following instructions is to show that  $\{a\}^2 \leq 2\|a\|^2$ .

#### Implementation

- Show that  $2\|a\|^2 - \{a\}^2$ 
  - $= 2\operatorname{re}(a)^2 + 2\operatorname{im}(a)^2 - (|\operatorname{re}(a)| + |\operatorname{im}(a)|)^2$
  - $= 2\operatorname{re}(a)^2 + 2\operatorname{im}(a)^2 - \operatorname{re}(a)^2 - 2|\operatorname{re}(a)||\operatorname{im}(a)| - \operatorname{im}(a)^2$
  - $= \operatorname{re}(a)^2 - 2|\operatorname{re}(a)||\operatorname{im}(a)| + \operatorname{im}(a)^2$
  - $= (|\operatorname{re}(a)| - |\operatorname{im}(a)|)^2$
  - $\geq 0$ .
- Hence show that  $\{a\}^2 \leq 2\|a\|^2$ .

#### Declaration IV:1

The notation  $\Delta_{x=y}^z f(x)$ , where  $x, z$  are complex numbers such that  $z \neq 0$  and  $f[x]$  is a function of  $x$ , will be used as a shorthand for  $\frac{f(y+z)-f(y)}{z}$ .

### Procedure IV:8(3.83)

#### Objective

Choose two functions  $f[x], g[x]$  and two complex numbers  $y, z$  such that  $z \neq 0$ . The objective of the following instructions is to show that  $\Delta_{x=y}^z(f(x) + g(x)) = \Delta_{x=y}^z f(x) + \Delta_{x=y}^z g(x)$ .

### Implementation

- Show that  $\Delta_{x=y}^z(f(x) + g(x))$ 
  - $= \frac{(f(y+z)+g(y+z))-(f(y)+g(y))}{z}$
  - $= \frac{f(y+z)-f(y)}{z} + \frac{g(y+z)-g(y)}{z}$
  - $= \Delta_{x=y}^z f(x) + \Delta_{x=y}^z g(x)$ .

### Procedure IV:9(3.84)

#### Objective

Choose a functions  $f[x]$  and complex numbers  $a, y, z$  such that  $z \neq 0$ . The objective of the following instructions is to show that  $\Delta_{x=y}^z(af(x)) = a\Delta_{x=y}^z f(x)$ .

#### Implementation

- Show that  $\Delta_{x=y}^z(af(x))$ 
  - $= \frac{af(y+z)-af(y)}{z}$
  - $= a\frac{f(y+z)-f(y)}{z}$
  - $= a\Delta_{x=y}^z f(x)$ .

### Procedure IV:10(mon1908191506)

#### Objective

Choose the following:

- A procedure  $q_0(x, n)$  to show that  $p'_n(x) \equiv 0$  (err  $a_0$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > b_0$  are chosen.
- A procedure  $q_1(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv p'_n(x)$  (err  $\frac{a_1}{n} + b_1\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > b_0$ , and  $0 < \|\delta\|^2 < c_1^2$  are chosen.
- A procedure  $q_2(x, n)$  to show that  $t'_n(x) \equiv 0$  (err  $a_2$ ) when a complex number  $x$  and a positive integer  $n$  such that  $R(x)$  and  $n > b_2$  are chosen.

4. A procedure  $q_3(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} t_n(y) \equiv t'_n(x)$  (err  $\frac{a_3}{n} + b_3\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $R(x)$ ,  $n > b_2$ , and  $0 < \|\delta\|^2 < c_3^2$  are chosen.

The objective of the following instructions is to construct the following:

1. Rational numbers  $a_4, b_4, a_5, b_5, c_5$ .
2. A procedure  $q_4(x, n)$  to show that  $p'_n(x) + t'_n(x) \equiv 0$  (err  $a_4$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$ ,  $R(x)$ , and  $n > b_4$  are chosen.
3. A procedure  $q_5(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} (p_n(y) + t_n(y)) \equiv p'_n(x) + t'_n(x)$  (err  $\frac{a_5}{n} + b_5\{\delta\}$ ) when two complex numbers  $x, \delta$  such that  $P(x)$ ,  $R(x)$ ,  $n > b_4$ , and  $0 < \|\delta\|^2 < c_5$ .

### Implementation

1. Let  $a_5 = a_1 + a_3$ .
2. Let  $b_5 = b_1 + b_3$ .
3. Let  $q_5(x, n, \delta)$  be the following procedure:
  - (a) Show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv p'_n(x)$  (err  $\frac{a_1}{n} + b_1\{\delta\}$ ) using procedure  $q_1$ .
  - (b) Show that  $\Delta_{y=x}^{+\delta} t_n(y) \equiv t'_n(x)$  (err  $\frac{a_3}{n} + b_3\{\delta\}$ ) using procedure  $q_3$ .
  - (c) Hence using **procedure IV:8**, show that  $\Delta_{y=x}^{+\delta} (p_n(y) + t_n(y))$ 
    - i.  $= \Delta_{y=x}^{+\delta} p_n(y) + \Delta_{y=x}^{+\delta} t_n(y)$
    - ii.  $\equiv p'_n(x) + \Delta_{y=x}^{+\delta} t_n(y)$  (err  $\frac{a_1}{n} + b_1\{\delta\}$ )
    - iii.  $\equiv p'_n(x) + t'_n(x)$  (err  $\frac{a_3}{n} + b_3\{\delta\}$ )
  - (d) **Hence show that**  $\Delta_{y=x}^{+\delta} (p_n(y) + t_n(y)) \equiv p'_n(x) + t'_n(x)$  (err  $\frac{a_5}{n} + b_5\{\delta\}$ ).
4. Let  $q_4(x, n)$  be the following procedure:
  - (a) Show that  $p'_n(x) \equiv 0$  (err  $a_0$ ) using procedure  $q_0$ .
  - (b) Show that  $t'_n(x) \equiv 0$  (err  $a_2$ ) using procedure  $q_2$ .
  - (c) Hence show that  $p'_n(x) + t'_n(x) \equiv 0$  (err  $a_0 + a_2$ ).
5. **Yield the tuple**  $\langle a_4, b_4, a_5, b_5, c_5, q_4, q_5 \rangle$ .

## Procedure IV:11(sat0308191134)

### Objective

Choose the following:

1. A procedure  $q_0(x, n)$  to show that  $p'_n(x) \equiv 0$  (err  $a_0$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$ , and  $n > b_0$  are chosen
2. A procedure  $q_1(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv p'_n(x)$  (err  $\frac{a_1}{n} + b_1\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > b_0$ , and  $0 < \|\delta\|^2 < c_1^2$  are chosen
3. A procedure  $q_2(x, n)$  to show that  $t'_n(x) \equiv 0$  (err  $a_2$ ) when a complex number  $x$  and a positive integer  $n$  such that  $R(x)$ , and  $n > b_2$  are chosen
4. A procedure  $q_3(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} t_n(y) \equiv t'_n(x)$  (err  $\frac{a_3}{n} + b_3\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $R(x)$ ,  $n > b_2$ , and  $0 < \|\delta\|^2 < c_3^2$  are chosen
5. A procedure  $q_4(x, n)$  to show that  $P(t_n(x))$  when a complex number  $x$  and a positive integer  $n$  such that  $R(x)$  and  $n > b_2$  are chosen

The objective of the following instructions is to construct the following:

1. Rational numbers  $a_5, b_5, a_6, b_6, c_6$ .
2. A procedure  $q_5(x, n)$  to show that  $p'_n(t_n(x))t'_n(x) \equiv 0$  (err  $a_5$ ) when a complex number  $x$  such that  $R(x)$ , and  $n > b_5$  are chosen.
3. A procedure  $q_6(x, n, \delta)$  to show that  $\Delta_{y=x}^{x+\delta} p_n(t_n(y)) \equiv p'_n(t_n(x))t'_n(x)$  (err  $\frac{a_6}{n} + b_6\{\delta\}$ ) when two complex numbers  $x, \delta$  such that  $R(x)$ ,  $n > b_5$ , and  $0 < \|\delta\|^2 < c_6^2$  are chosen.

### Implementation

1. Let  $a_5 = a_0 a_2$ .
2. Let  $b_5 = \max(b_0, b_2)$ .
3. Let  $a_6 = a_1 a_3 + a_1 a_2 + a_0 a_3$ .
4. Let  $b_6 = a_1 b_3 + b_1 a_3 + 2b_1 b_3 c_6 + b_1 a_2 + a_0 b_3$ .

5. Let  $c_6 = \min(c_3, \frac{c_1}{a_3 + 2b_3c_3 + a_2})$ .
6. Let  $q_5(x, n, \delta)$  be the following procedure:
  - (a) Show that  $P(t_n(x))$  using procedure  $q_4$ .
  - (b) If  $\Delta_{y=x}^{+\delta} t_n(y) = 0$ , then do the following:
    - i. Show that  $t_n(x + \delta) = t_n(x)$  given that  $t_n(x + \delta) - t_n(x) = 0\delta = 0$ .
    - ii. Hence using procedures  $q_0, q_3$ , show that  $\Delta_{y=x}^{+\delta} p_n(t_n(y))$ 
      - A.  $= \frac{p_n(t_n(x+\delta)) - p_n(t_n(x))}{\delta}$
      - B.  $= \frac{p_n(t_n(x)) - p_n(t_n(x))}{\delta}$
      - C.  $= 0$
      - D.  $= \Delta_{y=x}^{+\delta} t_n(y) p'_n(t_n(x))$
      - E.  $\equiv t'_n(x) p'_n(t_n(x))$  (err  $a_0(\frac{a_3}{n} + b_3\{\delta\})$ ).
  - (c) Otherwise do the following:
    - i. Using procedures  $q_3, q_4$ , show that  $\Delta_{y=x}^{+\delta} t_n(y)$ 
      - A.  $\equiv t'_n(x)$  (err  $\frac{a_3}{n} + b_3\{\delta\}$ )
      - B.  $\equiv 0$  (err  $a_2$ ).
    - ii. Show that  $\{\delta\} \leq 2c_6 \leq 2c_3$  given that  $\{\delta\}^2 \leq 2\|\delta\|^2 \leq 4c_6^2$ .
    - iii. Show that  $t_n(x + \delta) - t_n(x)$ 
      - A.  $= \Delta_{y=x}^{+\delta} t_n(y) \delta$
      - B.  $\equiv 0\delta$  (err  $(\frac{a_3}{n} + b_3\{\delta\} + a_2)c_6$ ) (err  $(a_3 + 2b_3c_3 + a_2)c_6$ ) (err  $c_1$ ).
    - iv. Hence using procedures  $q_0, q_1$ , show that  $\Delta_{z=t_n(x)}^{t_n(x+\delta)-t_n(x)} p_n(z)$ 
      - A.  $\equiv p'_n(t_n(x))$  (err  $\frac{a_1}{n} + b_1\{\delta\}$ )
      - B.  $\equiv 0$  (err  $a_0$ ).
    - v. Hence show that  $\Delta_{y=x}^{+\delta} p_n(t_n(y))$ 
      - A.  $= \Delta_{z=t_n(x)}^{t_n(x+\delta)-t_n(x)} p_n(z) \cdot \Delta_{y=x}^{+\delta} t_n(y)$
      - B.  $\equiv p'_n(t_n(x)) \Delta_{y=x}^{+\delta} t_n(y)$  (err  $(\frac{a_1}{n} + b_1\{\delta\})(\frac{a_3}{n} + b_3\{\delta\} + a_2)$ )
      - C.  $\equiv p'_n(t_n(x)) t'_n(x)$  (err  $a_0(\frac{a_3}{n} + b_3\{\delta\})$ ).
  - (d) **Hence show that**  $\Delta_{y=x}^{+\delta} p_n(t_n(y)) \equiv p'_n(t_n(x)) t'_n(x)$  (err  $\frac{a_6}{n} + b_6\{\delta\}$ ).
7. Let  $q_6(x, n)$  be the following procedure:

- (a) Show that  $P(t_n(x))$  using procedure  $q_4$ .
  - (b) Show that  $p'_n(t_n(x)) \equiv 0$  (err  $a_0$ ) using procedure  $q_0$ .
  - (c) Show that  $t'_n(x) \equiv 0$  (err  $a_2$ ) using procedure  $q_2$ .
  - (d) Hence show that  $p'_n(t_n(x)) t'_n(x) \equiv 0$  (err  $a_0 a_2$ ) (err  $a_5$ ).
8. **Yield the tuple**  $\langle a_5, b_5, a_6, b_6, c_6, q_5, q_6 \rangle$ .

## Procedure IV:12(tue2008191001)

### Objective

Choose the following:

1. A complex number  $B$
2. A procedure  $q_1(x, n)$  to show that  $p'_n(x) \equiv 0$  (err  $a_1$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > b_1$  are chosen.
3. A procedure  $q_2(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv p'(x)$  (err  $\frac{a_2}{n} + b_2\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > b_1$ , and  $0 < \|\delta\|^2 \leq c_2^2$  are chosen.

The objective of the following instructions is to construct the following:

1. Rational numbers  $a_3, b_3, a_4, b_4, c_4$ .
2. A procedure  $q_3(x, n)$  to show that  $Bp'_n(x) \equiv 0$  (err  $a_3$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > b_3$  are chosen.
3. A procedure  $q_4(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} (Bp_n(y)) \equiv Bp'_n(x)$  (err  $\frac{a_4}{n} + b_4\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > b_3$ , and  $0 < \|\delta\|^2 \leq c_4^2$  are chosen.

### Implementation

1. Let  $a_3 = \{B\}a_1$ .
2. Let  $b_3 = b_1$ .
3. Let  $a_4 = \{B\}a_2$ .
4. Let  $b_4 = \{B\}b_2$ .

5. Let  $c_4 = c_2$ .
6. Let  $q_3(x, n)$  be the following procedure:
  - (a) Show that  $p'_n(x) \equiv 0$  (err  $a_1$ ) using procedure  $q_1$ .
  - (b) Hence show that  $Bp'_n(x) \equiv 0B$  (err  $\{B\}a_1$ ) (err  $a_3$ ).
7. Let  $q_4(x, n, \delta)$  be the following procedure:
  - (a) Show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv p'(x)$  (err  $\frac{a_2}{n} + b_2\{\delta\}$ ) using procedure  $q_4$ .
  - (b) Hence show that  $B\Delta_{y=x}^{+\delta} p_n(y) \equiv Bp'(x)$ 
    - i. (err  $\{B\}(\frac{a_2}{n} + b_2\{\delta\})$ )
    - ii. (err  $\frac{a_4}{n} + b_4\{\delta\}$ ).
8. **Yield the tuple**  $\langle a_3, b_3, a_4, b_4, c_4, q_3, q_4 \rangle$ .

such that  $R(x)$ ,  $n > b_3$ , and  $0 < \|\delta\|^2 < c_5^2$  are chosen.

The objective of the following instructions is to construct the following:

1. Rational numbers  $a_6, b_6, a_7, b_7, c_7$ .
2. A procedure  $q_6(x, n)$  to show that  $p_n(x)t'_n(x) + p'_n(x)t_n(x) \equiv 0$  (err  $a_6$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$ ,  $R(x)$ , and  $n > b_6$  are chosen.
3. A procedure  $q_7(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta}(p_n(y)t_n(y)) \equiv p_n(x)t'_n(x) + p'_n(x)t_n(x)$  (err  $\frac{a_7}{n} + b_7\{\delta\}$ ) when two complex numbers  $x, \delta$  such that  $P(x)$ ,  $R(x)$ ,  $n > b_6$ , and  $0 < \|\delta\|^2 < c_7^2$  are chosen.

## Procedure IV:13(mon1908191207)

### Objective

Choose the following:

1. A procedure  $q_0(x, n)$  to show that  $p_n(x) \equiv 0$  (err  $a_0$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > b_0$  are chosen.
2. A procedure  $q_1(x, n)$  to show that  $p'_n(x) \equiv 0$  (err  $a_1$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > b_0$  are chosen.
3. A procedure  $q_2(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv p'_n(x)$  (err  $\frac{a_2}{n} + b_2\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > b_0$ , and  $0 < \|\delta\|^2 < c_2^2$  are chosen.
4. A procedure  $q_3(x, n)$  to show that  $t_n(x) \equiv 0$  (err  $a_3$ ) when a complex number  $x$  and a positive integer  $n$  such that  $R(x)$  and  $n > b_3$  are chosen.
5. A procedure  $q_4(x, n)$  to show that  $t'_n(x) \equiv 0$  (err  $a_4$ ) when a complex number  $x$  and a positive integer  $n$  such that  $R(x)$  and  $n > b_3$  are chosen.
6. A procedure  $q_5(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} t_n(y) \equiv t'_n(x)$  (err  $\frac{a_5}{n} + b_5\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$

### Implementation

1. Let  $a_6 = a_0a_4 + a_1a_3$ .
2. Let  $b_6 = \max(b_0, b_3)$ .
3. Let  $a_7 = 0$ .
4. Let  $b_7 = (a_5 + b_5c_7 + a_4)(a_2 + b_2c_7 + a_1)$ .
5. Let  $c_7 = \min(c_2, c_5)$ .
6. Let  $q_7(x, n, \delta)$  be the following procedure:
  - (a) Show that  $\{\delta\} \leq 2c_7$  given that  $\{\delta\}^2 \leq 2\|\delta\|^2 \leq 4c_7^2$ .
  - (b) Hence using procedures  $q_2, q_1$ , show that  $\Delta_{y=x}^{+\delta} p_n(y)$ 
    - i.  $\equiv p'_n(x)$  (err  $\frac{a_2}{n} + b_2\{\delta\}$ )
    - ii.  $\equiv 0$  (err  $a_1$ ).
  - (c) Hence using procedures  $q_5, q_4$ , show that  $\Delta_{y=x}^{+\delta} t_n(y)$ 
    - i.  $\equiv t'_n(x)$  (err  $\frac{a_5}{n} + b_5\{\delta\}$ )
    - ii.  $\equiv 0$  (err  $a_4$ ).
  - (d) Show that  $p_n(x) \equiv 0$  (err  $a_0$ ) using procedure  $q_0$ .
  - (e) Show that  $t_n(x) \equiv 0$  (err  $a_6$ ) using procedure  $q_3$ .
  - (f) Hence show that  $\Delta_{y=x}^{+\delta}(p_n(y)t_n(y))$ 
    - i.  $= p_n(x + \delta)\Delta_{y=x}^{+\delta} t_n(y) + t_n(x)\Delta_{y=x}^{+\delta} p_n(y)$

- ii.  $= (p_n(x) + \delta \Delta_{y=x}^{+\delta} p_n(y)) \Delta_{y=x}^{+\delta} t_n(y) + t_n(x) \Delta_{y=x}^{+\delta} p_n(y)$
- iii.  $= p_n(x) \Delta_{y=x}^{+\delta} t_n(y) + \delta \Delta_{y=x}^{+\delta} p_n(y) \Delta_{y=x}^{+\delta} t_n(y) + t_n(x) \Delta_{y=x}^{+\delta} p_n(y)$
- iv.  $\equiv p_n(x) \Delta_{y=x}^{+\delta} t_n(y) + 0 \delta \Delta_{y=x}^{+\delta} p_n(y) + t_n(x) \Delta_{y=x}^{+\delta} p_n(y)$  (err  $(\frac{a_5}{n} + b_5\{\delta\} + a_4)\{\delta\}(\frac{a_2}{n} + b_2\{\delta\} + a_1)$ )
- (g) **Hence show that**  $\Delta_{y=x}^{+\delta}(p_n(y)t_n(y)) \equiv p_n(x) \Delta_{y=x}^{+\delta} t_n(y) + t_n(x) \Delta_{y=x}^{+\delta} p_n(y)$  (err  $\frac{a_7}{n} + b_7\{\delta\}$ ).

7. Let  $q_6(x, n)$  be the following procedure:

- (a) Show that  $p'_n(x) \equiv 0$  (err  $a_1$ ) using procedure  $q_1$ .
- (b) Show that  $t'_n(x) \equiv 0$  (err  $a_4$ ) using procedure  $q_4$ .
- (c) Show that  $p_n(x) \equiv 0$  (err  $a_0$ ) using procedure  $q_0$ .
- (d) Show that  $t_n(x) \equiv 0$  (err  $a_3$ ) using procedure  $q_3$ .
- (e) **Hence show that**  $p_n(x)t'_n(x) + p'_n(x)t_n(x) \equiv 0$  (err  $a_0a_4 + a_1a_3$ ) (err  $a_6$ ).

8. **Yield the tuple**  $\langle a_6, b_6, a_7, b_7, c_7, q_6, q_7 \rangle$ .

## Procedure IV:14(fri2308191803)

### Objective

Choose the following:

- 1. A procedure  $q_0(x, n)$  to show that  $p'_n(x) \equiv q'_n(x)$  (err  $\frac{a_0}{n}$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > b_0$  are chosen.
- 2. A procedure  $q_1(x, n)$  to show that  $p'_n(x) \equiv 0$  (err  $a_1$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > b_1$  are chosen.
- 3. A procedure  $q_2(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv p'_n(x)$  (err  $\frac{a_2}{n} + b_2\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > b_1$ , and  $0 < \|\delta\|^2 \leq c_2^2$  are chosen.

The objective of the following instructions is to construct the following:

- 1. Rational numbers  $a_3, b_3, a_4, b_4, c_4$ .
- 2. A procedure  $q_3(x, n)$  to show that  $q'_n(x) \equiv 0$  (err  $a_3$ ) when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > b_3$  are chosen.
- 3. A procedure  $q_4(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv q'_n(x)$  (err  $\frac{a_4}{n} + b_4\{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > b_3$ , and  $0 < \|\delta\|^2 \leq c_4^2$  are chosen.

### Implementation

- 1. Let  $a_3 = a_0 + a_1$ .
- 2. Let  $b_3 = \max(b_0, b_1)$ .
- 3. Let  $a_4 = a_0 + a_2$ .
- 4. Let  $b_4 = b_2$ .
- 5. Let  $c_4 = c_2$ .
- 6. Let  $q_3(x, n)$  be the following procedure:
  - (a) Show that  $p'_n(x) \equiv q'_n(x)$  (err  $\frac{a_0}{n}$ ) using procedure  $q_0$ .
  - (b) Show that  $p'_n(x) \equiv 0$  (err  $a_1$ ) using procedure  $q_1$ .
  - (c) **Hence show that**  $q'_n(x) \equiv 0$  (err  $a_3$ ).
- 7. Let  $q_4(x, n, \delta)$  be the following procedure:
  - (a) Using procedures  $q_0, q_2$ , show that  $\Delta_{y=x}^{+\delta} p_n(y)$ 
    - i.  $\equiv p'_n(x)$  (err  $\frac{a_2}{n} + b_2\{\delta\}$ )
    - ii.  $\equiv q'_n(x)$  (err  $\frac{a_0}{n}$ ).
  - (b) Hence show that  $\Delta_{y=x}^{+\delta} p_n(y) \equiv q'_n(x)$ 
    - i. (err  $\frac{a_2}{n} + b_2\{\delta\} + \frac{a_0}{n}$ )
    - ii. (err  $\frac{a_4}{n} + b_4\{\delta\}$ ).
- 8. **Yield the tuple**  $\langle a_3, b_3, a_4, b_4, c_4, q_3, q_4 \rangle$ .

# Chapter 13

## Common Derivatives

### Procedure IV:15(tue2008191151)

#### Objective

Choose a complex number  $B$  and a rational number  $D > 0$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , a procedure  $p(x, n)$  to show that  $0 \equiv 0$  (err  $a$ ) when a complex number  $x$  and a positive integer  $n$  such that  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} B \equiv 0$  (err  $\frac{b}{n} + c\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

#### Implementation

1. Let  $a = b = c = d = 0$ .
2. Let  $p(x, n)$  be the following procedure:
  - (a) **Show that**  $0 \equiv 0$  (err  $a$ ).
3. Let  $q(x, n, \delta)$  be the following procedure:
  - (a) **Show that**  $\Delta_{y=x}^{+\delta} B \equiv 0$  (err  $\frac{b}{n} + c\{\delta\}$ ).
4. **Yield the tuple**  $\langle a, b, c, d, p, q \rangle$ .

### Procedure IV:16(tue2008191209)

#### Objective

Choose a positive integer  $N$  and positive rational numbers  $X, D$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , a procedure  $p(x, n)$  to show that  $Nx^{N-1} \equiv 0$  (err  $a$ ) when a complex number  $x$  and a positive integer

$n$  such that  $\|x\|^2 \leq X^2$  and  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} y^N \equiv Nx^{N-1}$  (err  $\frac{b}{n} + c\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

#### Implementation

1. Let  $a = NX^{N-1}$ .
2. Let  $b = d = 0$ .
3. Let  $c = \sum_r^{[0:N-1]} \binom{N}{r} X^r D^{N-r-2}$ .
4. Let  $p(x, n)$  be the following procedure:
  - (a) **Show that**  $Nx^{N-1} \equiv 0$  (err  $Nx^{N-1}$ )  
(err  $NX^{N-1}$ ) (err  $a$ ).
5. Let  $q(x, n, \delta)$  be the following procedure:
  - (a) Show that  $\Delta_{y=x}^{+\delta} y^N \equiv Nx^{N-1}$ 
    - i. (err  $\frac{(x+\delta)^N - x^N}{\delta} - Nx^{N-1}$ )
    - ii. (err  $\frac{1}{\delta} (\sum_r^{[0:N+1]} \binom{N}{r} x^r \delta^{N-r} - x^N) - Nx^{N-1}$ )
    - iii. (err  $\sum_r^{[0:N]} \binom{N}{r} x^r \delta^{N-r-1} - Nx^{N-1}$ )
    - iv. (err  $\delta (\sum_r^{[0:N-1]} \binom{N}{r} x^r \delta^{N-r-2})$ )
    - v. (err  $\frac{b}{n} + c\{\delta\}$ ).
6. **Yield the tuple**  $\langle a, b, c, d, p, q \rangle$ .

## Procedure IV:17(tue2008191254)

### Objective

Choose two rational numbers  $X > D > 0$ . The objective of the following instructions is to construct rational numbers  $a, b, c, d$ , a procedure  $p(x, n)$  to show that  $-\frac{1}{x^2} \equiv 0$  (err  $a$ ) when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \geq X^2$  and  $n > b$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} \frac{1}{y} \equiv -\frac{1}{x^2}$  (err  $\frac{c}{n} + d\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

### Implementation

1. Let  $a = \frac{1}{X^2}$ .
2. Let  $b = c = 0$ .
3. Let  $d = \frac{1}{X^2(X-D)}$ .
4. Let  $p(x, n)$  be the following procedure:
  - (a) **Show that**  $\frac{1}{x^2} \equiv 0$  (err  $\frac{1}{x^2}$ ) (err  $\frac{1}{X^2}$ ) (err  $a$ ).
5. Let  $q(x, n, \delta)$  be the following procedure:
  - (a) Show that  $\Delta_{y=x}^{+\delta} \frac{1}{y} \equiv -\frac{1}{x^2}$ 
    - i. (err  $\frac{1}{\delta}(\frac{1}{x+\delta} - \frac{1}{x}) + \frac{1}{x^2}$ )
    - ii. (err  $\frac{1}{\delta} \cdot \frac{-\delta}{x(x+\delta)} + \frac{1}{x^2}$ )
    - iii. (err  $\frac{1}{x^2} - \frac{1}{x(x+\delta)}$ )
    - iv. (err  $\frac{\delta}{x^2(x+\delta)}$ )
    - v. (err  $\frac{\{\delta\}}{X^2(X-D)}$ )
    - vi. (err  $\frac{c}{n} + d\{\delta\}$ ).
6. **Yield the tuple**  $\langle a, b, c, d, p, q \rangle$ .

## Procedure IV:18(tue2008191341)

### Objective

Choose a positive integer  $N$  and positive rational numbers  $X < Y$ . The objective of the following instructions is to construct positive rational numbers  $a, b, c, d, e$ , a procedure  $p(x, n)$  to show that  $-Nx^{-N-1} \equiv 0$  (err  $a$ ) when a complex number  $x$  and a positive integer  $n$  such that  $X^2 \leq \|x\|^2 \leq Y^2$  and  $n > b$  are chosen, and a procedure  $q(x, n, \delta)$  to

show that  $\Delta_{y=x}^{+\delta} y^{-N} \equiv -Nx^{-N-1}$  (err  $\frac{c}{n} + d\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq e^2$  is chosen.

### Implementation

1. Execute the following in post-order:
  - (a) Execute **procedure IV:11** on  $\langle q_2, q_3, q_4, q_5, q_6 \rangle$  and let  $\langle a, b, c, d, e, q_0, q_1 \rangle$  receive.
    - i. Execute **procedure IV:17** on  $\langle X^N, \frac{X^N}{2} \rangle$  and let  $\langle \dots, q_2, q_3 \rangle$  receive.
    - ii. Execute **procedure IV:16** on  $\langle N, Y, Y \rangle$  and let  $\langle \dots, q_4, q_5 \rangle$  receive.
    - iii. Let  $q_6(x, n)$  be the following procedure:
      - A. Show that  $\|x^N\|^2 = (\|x\|^2)^N \geq (X^2)^N = (X^N)^2$ .
2. Let  $p(x, n)$  be the following procedure:
  - (a) **Show that**  $-Nx^{-N-1} = -\frac{1}{(x^N)^2} \cdot Nx^{N-1} \equiv 0$  (err  $a$ ) **using procedure**  $q_0$ .
3. Let  $q(x, n, \delta)$  be the following procedure:
  - (a) Using procedure  $q_1$ , show that  $\Delta_{y=x}^{+\delta} y^{-N}$ 
    - i.  $= \frac{1}{\delta}(((x+\delta)^N)^{-1} - (x^N)^{-1})$
    - ii.  $\equiv -\frac{1}{(x^N)^2} \cdot Nx^{N-1}$  (err  $\frac{c}{n} + d\{\delta\}$ )
    - iii.  $= -Nx^{-N-1}$
  - (b) **Hence show that**  $\Delta_{y=x}^{+\delta} y^{-N} \equiv -Nx^{-N-1}$  (err  $\frac{c}{n} + d\{\delta\}$ ).
4. **Yield the tuple**  $\langle a, b, c, d, e, p, q \rangle$ .

## Procedure IV:19(3.18)

### Objective

Choose a rational number  $D \geq 0$ . The objective of the following instructions is to construct two rational numbers  $a, c$  and a procedure,  $p(n, \delta)$ , to show that  $\Delta_{x=0}^{+\delta} \exp_n(x) \equiv 1$  (err  $a\delta$ ) (err  $a\{\delta\}$ ) when a complex number  $\delta$  and a positive integer  $n$  such that  $0 < \|\delta\|^2 \leq D^2$  and  $n > c$  are chosen.

## Implementation

1. Execute **procedure III:34** on  $\langle D \rangle$  and let  $\langle a, c, q \rangle$  receive.
2. Let  $p(\delta, n)$  be the following procedure:
  - (a) Now using **procedure II:29**, and procedure  $q$ , show that  $\exp_n(\delta) - 1 \equiv \delta$ 
    - i.  $(\text{err } \exp_n(\delta) - 1 - \delta)$
    - ii.  $(\text{err } (1 + \frac{\delta}{n})^n - 1 - \delta)$
    - iii.  $(\text{err } \frac{\delta}{n} \sum_r^{[0:n]} (1 + \frac{\delta}{n})^r - n \frac{\delta}{n})$
    - iv.  $(\text{err } \frac{\delta}{n} \sum_r^{[0:n]} ((1 + \frac{\delta}{n})^r - 1))$
    - v.  $(\text{err } \frac{\delta}{n} \sum_r^{[0:n]} \frac{\delta}{n} \sum_k^{[0:r]} (1 + \frac{\delta}{n})^k)$
    - vi.  $(\text{err } \frac{\delta^2}{n^2} \sum_r^{[0:n]} \sum_k^{[0:r]} (1 + \frac{\delta}{n})^k)$
    - vii.  $(\text{err } \frac{\delta^2}{n^2} \sum_r^{[0:n]} \sum_k^{[0:r]} a)$
    - viii.  $(\text{err } \delta^2 a)$ .
  - (b) **Therefore show that**  $\Delta_{x=0}^{+\delta} \exp_n(x) \equiv 1 (\text{err } a\delta) (\text{err } a\{\delta\})$ .
3. **Yield the tuple**  $\langle a, c, p \rangle$ .

## Procedure IV:20(3.19)

### Objective

Choose two rational numbers  $X \geq 0$ ,  $D \geq 0$ . The objective of the following instructions is to construct rational numbers  $l, a, b, d$ , a procedure  $t(x, n)$  to show that  $\exp_n(x) \equiv 0 (\text{err } l)$  when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$  and  $n > d$  are chosen, and a procedure,  $q(x, n, \delta)$ , to show that  $\Delta_{y=x}^{+\delta} \exp_n(y) \equiv \exp_n(x) (\text{err } \frac{a}{n} + b\{\delta\})$  when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

### Implementation

1. Execute **procedure III:36** on  $\langle \max(X, D) \rangle$  and let  $\langle e, f, u \rangle$  receive.
2. Execute **procedure IV:19** on  $\langle X \rangle$  and let  $\langle h, j, r \rangle$  receive.
3. Execute **procedure III:34** on  $\langle X \rangle$  and let  $\langle l, m, t \rangle$  receive.

4. Let  $a = eX$ .
5. Let  $b = lh$ .
6. Let  $d = \max(f, j, m)$ .
7. Let  $p(x, n, \delta)$  be the following procedure:
  - (a) Using procedures  $u, r, t$ , show that  $\Delta_{y=x}^{+\delta} \exp_n(y)$ 
    - i.  $= \frac{\exp_n(x+\delta) - \exp_n(x)}{\delta}$
    - ii.  $= \frac{\exp_n(x) \exp_n(\delta) - \exp_n(x)}{\delta}$ 
      - A.  $(\text{err } \frac{ex\delta}{n\delta})$
      - B.  $(\text{err } \frac{eX}{n})$
    - iii.  $= \exp_n(x) \Delta_{y=0}^{\delta} \exp_n(y)$
    - iv.  $\equiv \exp_n(x) \cdot 1 (\text{err } lh\{\delta\})$ .
  - (b) **Hence show that**  $\Delta_{y=x}^{+\delta} \exp_n(y) \equiv \exp_n(x) (\text{err } \frac{ex}{n} + lh\{\delta\}) (\text{err } \frac{a}{n} + b\{\delta\})$ .
8. **Yield the tuple**  $\langle a, b, d, p \rangle$ .

## Procedure IV:21(3.27)

### Objective

Choose non-negative rational numbers  $X, D$ . The objective of the following instructions is to construct rational numbers  $l, d, a, b$ , a procedure  $q(x, n)$  to show that  $\cos_n(x) \equiv 0 (\text{err } l)$  when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$  and  $n > d$  are chosen, and a procedure  $p(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} \sin_n(y) \equiv \cos_n(x) (\text{err } \frac{a}{n} + b\{\delta\})$  when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

### Implementation

- 1) Execute the following in post-order:
  - a) Execute **procedure IV:12** on  $\langle \frac{1}{2i}, q_2, q_3 \rangle$  and let  $\langle l, d, a, b, D, q_0, q_1 \rangle$  receive.
  - i) Execute **procedure IV:10** on  $\langle q_4, q_5, q_6, q_7 \rangle$  and let  $\langle q_2, q_3 \rangle$  receive.
- (1) Execute **procedure IV:11** on  $\langle q_8, q_9, q_{10}, q_{11}, q_{12} \rangle$  and let  $\langle q_4, q_5 \rangle$  receive.
  - (a) Execute **procedure IV:20** on  $\langle X, D \rangle$  and let  $\langle q_8, q_9 \rangle$  receive.



- (b) Execute **procedure IV:12** on  $\langle i, q_{13}, q_{14} \rangle$  and let  $\langle q_{10}, q_{11} \rangle$  receive.
- (i) Execute **procedure IV:16** on  $\langle 1, X, D \rangle$  and let  $\langle q_{13}, q_{14} \rangle$  receive.
- (c) Let  $q_{12}(x, n)$  be the following procedure:
- (i) Show that  $\|ix\|^2 = \|x\|^2 \leq X^2$ .
- (2) Execute **procedure IV:12** on  $\langle -1, q_{15}, q_{16} \rangle$  and let  $\langle q_6, q_7 \rangle$ .
- (a) Execute **procedure IV:11** on  $\langle q_{17}, q_{18}, q_{19}, q_{20}, q_{21} \rangle$  and let  $\langle q_{15}, q_{16} \rangle$  receive.
- (i) Execute **procedure IV:20** on  $\langle X, D \rangle$  and let  $\langle q_{17}, q_{18} \rangle$  receive.
- (ii) Execute **procedure IV:12** on  $\langle -i, q_{22}, q_{23} \rangle$  and let  $\langle q_{19}, q_{20} \rangle$  receive.
- (1) Execute **procedure IV:16** on  $\langle 1, X, D \rangle$  and let  $\langle q_{22}, q_{23} \rangle$  receive.
- (iii) Let  $q_{21}(x, n)$  be the following procedure:
- (1) Show that  $\|-ix\|^2 = \|x\|^2 \leq X^2$ .
- 2) Let  $p(x, n)$  be the following procedure:
1. Using procedure  $q_0$ , show that  $\cos_n(x)$
- (a)  $= \frac{1}{2}(\exp_n(ix) + \exp_n(-ix))$
- (b)  $= \frac{1}{2i}(\exp_n(ix^1) \cdot i \cdot 1x^0 + (-1)\exp_n(-ix^1) \cdot (-i) \cdot 1 \cdot x^0)$
- (c)  $\equiv 0$  (err  $l$ ).
- 3) Let  $q(x, n, \delta)$  be the following procedure:
1. Using procedure  $q_1$ , show that  $\Delta_{y=x}^{+\delta} \sin_n(y)$
- (a)  $= \Delta_{y=x}^{+\delta} \left( \frac{\exp_n(ix) - \exp_n(-ix)}{2i} \right)$
- (b)  $= \Delta_{y=x}^{+\delta} \left( \frac{1}{2i}(\exp_n(ix^1) + (-1)\exp_n((-i)x^1)) \right)$
- (c)  $\equiv \frac{1}{2i}(\exp_n(ix^1) \cdot i \cdot 1x^0 + (-1)\exp_n(-ix^1) \cdot (-i) \cdot 1 \cdot x^0)$  (err  $\frac{a}{n} + b\{\delta\}$ )
- (d)  $= \frac{\exp_n(ix) + \exp_n(-ix)}{2}$
- (e)  $= \cos_n(x)$ .
2. Hence show that  $\Delta_{y=x}^{+\delta} \sin_n(y) \equiv \cos_n(x)$  (err  $\frac{a}{n} + b\{\delta\}$ ).
- 4) Yield the tuple  $\langle l, d, a, b, D, p, q \rangle$ .

## Procedure IV:22(3.28)

### Objective

Choose non-negative rational numbers  $X, D$ . The objective of the following instructions is to construct rational numbers  $l, d, a, b$ , a procedure  $q(x, n)$  to show that  $-\sin_n(x) \equiv 0$  (err  $l$ ) when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$  and  $n > d$  are chosen, and a procedure  $p(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} \cos_n(y) \equiv -\sin_n(x)$  (err  $\frac{a}{n} + b\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

### Implementation

Implementation is analogous to that of **procedure IV:21**.

## Procedure IV:23(wed2108191034)

### Objective

Choose non-negative rational numbers  $X, D$  such that  $X + D < 1$ . The objective of the following instructions is to construct rational numbers  $l, d, a, b$ , a procedure  $p(x, n)$  to show that  $(1 + x)_{n-1}^{-1} \equiv 0$  (err  $l$ ) when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$  and  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} \ln_n(1 + y) \equiv (1 + x)_{n-1}^{-1}$  (err  $\frac{a}{n} + b\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

### Implementation

1. Execute **procedure III:55** on  $\langle 1, X \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Let  $l = a_1$ .
3. Let  $d = 1$ .
4. Let  $a = 0$ .
5. Let  $b = \frac{1}{D^2((X+D)^{-1}-1)}$ .
6. Let  $p(x, n)$  be the following procedure:
  - (a) Using procedure  $p_1$ , show that  $(1 + x)_{n-1}^{-1}$ 
    - i.  $= ((1 + x)_{n-1}^{-1})^1$

- ii.  $\equiv 0$  (err  $a_1$ ).
7. Let  $q(x, n, \delta)$  be the following procedure:
- (a) Using **procedure II:30**, show that  $\Delta_{y=x}^{+\delta} \ln_n(1+y) \equiv (1+x)_{n-1}^{-1}$
- i. (err  $\Delta_{y=x}^{+\delta} \ln_n(1+y) - (1+x)_{n-1}^{-1}$ )
- ii. (err  $\frac{1}{\delta} (\sum_r^{[1:n]} \frac{(-1)^{r-1}}{r} (x + \delta)^r - \sum_r^{[1:n]} \frac{(-1)^{r-1}}{r} x^r) - \sum_r^{[0:n-1]} \frac{(-1)^r}{r} x^r$ )
- iii. (err  $\frac{1}{\delta} \sum_r^{[1:n]} \frac{(-1)^{r-1}}{r} (\sum_k^{[0:r+1]} \binom{r}{k} x^k \delta^{r-k} - x^r) - \sum_r^{[0:n-1]} (-1)^r x^r$ )
- iv. (err  $\sum_r^{[1:n]} \frac{(-1)^{r-1}}{r} \sum_k^{[0:r]} \binom{r}{k} x^k \delta^{r-1-k} - \sum_r^{[0:n-1]} (-1)^r x^r$ )
- v. (err  $\sum_r^{[1:n]} \frac{(-1)^{r-1}}{r} (\sum_k^{[0:r]} \binom{r}{k} x^k \delta^{r-1-k} - r x^{r-1})$ )
- vi. (err  $\delta (\sum_r^{[1:n]} \frac{(-1)^{r-1}}{r} \sum_k^{[0:r-1]} \binom{r}{k} x^k \delta^{r-2-k})$ )
- vii. (err  $\delta (\sum_r^{[1:n]} \sum_k^{[0:r-1]} \binom{r}{k} X^k D^{r-2-k})$ )
- viii. (err  $\delta (\frac{1}{D^2} \sum_r^{[1:n]} \sum_k^{[0:r-1]} \binom{r}{k} X^k D^{r-k})$ )
- ix. (err  $\delta (\frac{1}{D^2} \sum_r^{[1:n]} (X+D)^r)$ )
- x. (err  $\delta (\frac{1}{D^2((X+D)^{-1}-1)})$ )
- xi. (err  $\frac{a}{n} + b\{\delta\}$ ).
8. **Yield the tuple**  $\langle l, d, a, b, p, q \rangle$ .

## Procedure IV:24(wed2108191140)

### Objective

Choose non-negative rational numbers  $X, D$  such that  $X + D < 1$ . The objective of the following instructions is to construct rational numbers  $l, d, a, b$ , a procedure  $p(x, n)$  to show that  $\frac{1}{1+x} \equiv 0$  (err  $l$ ) when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$  and  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} \ln_n(1+y) \equiv \frac{1}{1+x}$  (err  $\frac{a}{n} + b\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

### Implementation

1. Execute **procedure III:53** on  $\langle X, 1 \rangle$  and let  $\langle a_1, b_1, p_1 \rangle$  receive.

2. Execute **procedure IV:23** on  $\langle X, D \rangle$  and let  $\langle a_2, b_2, c_2, b, p_2, q_2 \rangle$  receive.
3. Let  $l = \frac{1}{1-X}$ .
4. Let  $d = \max(1, b_2)$ .
5. Let  $a = 2a_1l + c_2$ .
6. Let  $p(x, n)$  be the following procedure:
- (a) Show that  $|\operatorname{re}(x)| \leq X$  given that  $\operatorname{re}(x)^2 \leq \|x\|^2 \leq X^2$ .
- (b) Hence show that  $\|1+x\|^2$
- i.  $\geq \operatorname{re}(1+x)^2$
- ii.  $= (1 + \operatorname{re}(x))^2$
- iii.  $\leq (1-X)^2$ .
- (c) Hence show that  $\frac{1}{1+x} \equiv 0$  (err  $\frac{1}{1+x}$ ) (err  $l$ ).
7. Let  $q(x, n, \delta)$  be the following procedure:
- (a) Show that  $\|\frac{1}{1+x}\|^2 \leq l^2$  using procedure  $p$ .
- (b) Show that  $\|(n-1)(-x)^{n-1}\|^2 \leq (a_1b_1^{n-1})^2 \leq a_1^2$  using procedure  $p_1$ .
- (c) Hence show that  $\|(-x)^{n-1}\|^2 \leq (\frac{a_1}{n-1})^2 \leq (\frac{2a_1}{n})^2$ .
- (d) Now using procedure  $q_2$ , show that  $\Delta_{y=x}^{+\delta} \ln_n(1+y)$
- i.  $\equiv (1+x)_{n-1}^{-1}$  (err  $\frac{c_2}{n} + b\{\delta\}$ )
- ii.  $= \frac{1-(-x)^{n-1}}{1-(-x)}$
- iii.  $\equiv \frac{1}{1+x}$  (err  $\frac{(-x)^{n-1}}{1+x}$ ) (err  $\frac{2a_1l}{n}$ ).
- (e) **Hence show that**  $\Delta_{y=x}^{+\delta} \ln_n(1+y) \equiv \frac{1}{1+x}$  (err  $\frac{c_2}{n} + b\{\delta\} + \frac{2a_1l}{n}$ ) (err  $\frac{a}{n} + b\{\delta\}$ ).
8. **Yield the tuple**  $\langle l, d, a, b, p, q \rangle$ .

## Procedure IV:25(sun0812191401)

### Objective

Choose a rational number  $X$  such that  $0 < X \leq 1$ . The objective of the following instructions is to construct a rational number  $f$  such that  $0 \leq f < 1$ , and a procedure  $p(x)$  to show that  $\operatorname{re}(\frac{x-1}{x+1}) \geq -f$  and  $\|\frac{x-1}{x+1}\|^2 \leq 1$  when a complex number  $x$  such that  $\operatorname{re}(x) \geq 0$  and  $\|x\|^2 \geq X^2$  is chosen.

### Implementation

1. Let  $f = \frac{1-X^2}{1+X^2}$ .
2. Verify that  $0 \leq f < 1$ .
3. Let  $p(x)$  be the following procedure:
  - (a) Show that  $\|\frac{x-1}{x+1}\|^2 \leq 1$ 
    - i. given that  $\|x-1\|^2 \leq \|x+1\|^2$
    - ii. given that  $(\operatorname{re}(x)-1)^2 \leq (\operatorname{re}(x)+1)^2$
    - iii. given that  $0 \leq \operatorname{re}(x)$ .
  - (b) Show that  $\operatorname{re}(\frac{x-1}{x+1})$ 
    - i.  $= \operatorname{re}(\frac{(x-1)(\overline{x+1})}{(x+1)(\overline{x+1})})$
    - ii.  $= \operatorname{re}(\frac{\|x\|^2 + x\overline{x} - 1}{\|x\|^2 + x\overline{x} + 1})$
    - iii.  $= \frac{\|x\|^2 - 1}{\|x\|^2 + 2\operatorname{re}(x) + 1}$
    - iv.  $\geq \frac{X^2 - 1}{\|x\|^2 + 2\operatorname{re}(x) + 1}$
    - v.  $\geq \frac{X^2 - 1}{X^2 + 1} = -f$ .

### Declaration IV:2

The notation  $\ln_n(x)$ , where  $x$  is a complex number, will be used as a shorthand for  $\ln_n(1 + \frac{x-1}{x+1}) - \ln_n(1 - \frac{x-1}{x+1})$  when  $\operatorname{re}(x) \geq 0$ ,  $\ln_n(\frac{x}{i}) + \frac{\tau_n}{4}i$  when  $\operatorname{im}(x) \geq 0$ , and  $\ln_n(xi) - \frac{\tau_n}{4}i$  if otherwise.

### Procedure IV:26(thu2208191250)

#### Objective

Choose a rational number  $X$  such that  $1 \geq X > 0$ . The objective of the following instructions is to construct a positive rational number  $a$ , and a procedure  $p(x, k)$  to show that  $\|\ln_k(x)\|^2 \leq a^2$  when a positive integer  $k$  and a complex number  $x$  such that  $\|x\|^2 \geq X^2$  and  $\operatorname{re}(x) \geq 0$  are chosen.

#### Implementation

1. Execute **procedure IV:25** on  $\langle X \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Verify that  $0 < a_1 < 1$ .

3. Execute **procedure III:70** on  $\langle a_1 \rangle$  and let  $\langle a_2, p_2 \rangle$  receive.
4. Let  $a = 2a_2$ .
5. Let  $p(x, k)$  be the following procedure:
  - (a) Show that  $\|\frac{x-1}{x+1}\|^2 \leq a_1^2$  using procedure  $p_1$ .
  - (b) Show that  $\|\ln_k(1 + \frac{x-1}{x+1})\|^2 \leq a_2^2$  using procedure  $p_2$ .
  - (c) Show that  $\|\ln_k(1 - \frac{x-1}{x+1})\|^2 \leq a_2^2$  using procedure  $p_2$ .
  - (d) Hence show that  $\|\ln_k(x)\|^2$ 
    - i.  $= \|\ln_k(1 + \frac{x-1}{x+1}) - \ln_k(1 - \frac{x-1}{x+1})\|^2$
    - ii.  $\leq a^2$ .
6. Yield the tuple  $\langle a, p \rangle$ .

### Procedure IV:27(sun0812191440)

#### Objective

Choose a rational number  $X$  such that  $0 < X \leq 1$ . The objective of the following instructions is to construct positive rational numbers  $a, b$ , and a procedure  $p(x, k)$  to show that  $\|\ln_k(x)\|^2 \leq a^2$  when a positive integer  $k$  and a complex number  $x$  such that  $\|x\|^2 \geq X^2$  and  $k > b$  are chosen.

#### Implementation

1. Execute **procedure IV:26** on  $\langle X \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Execute **procedure III:74** and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Verify that  $a_2 > 0$ .
4. Let  $a = a_1 + \frac{a_2}{4}$ .
5. Let  $p(x, k)$  be the following procedure:
  - (a) Show that  $\frac{1}{4}\tau_k \leq \frac{a_2}{4}$  using procedure  $p_2$ .
  - (b) If  $\operatorname{re}(x) \geq 0$ , then do the following:
    - i. **Show that  $\|\ln_k(x)\|^2 \leq a_1^2 \leq a^2$  using procedure  $p_1$ .**
  - (c) Otherwise if  $\operatorname{im}(x) \geq 0$ , then do the following:
    - i. Show that  $\|\frac{x}{i}\|^2 = \|x\|^2 \geq X^2$ .

- ii. Show that  $\operatorname{re}(\frac{x}{i}) = \operatorname{re}(\operatorname{im}(x) - \operatorname{re}(x)i) = \operatorname{im}(x) \geq 0$ .
  - iii. Hence show that  $\|\ln_k(\frac{x}{i})\|^2 \leq a_1^2$  using procedure  $p_1$ .
  - iv. **Hence show that**  $\|\ln_k(x)\|^2 = \|\ln_k(\frac{x}{i}) + \frac{\tau_k}{4}i\|^2 \leq (a_1 + \frac{a_2}{4})^2 = a^2$ .
  - (d) Otherwise do the following:
    - i. Show that  $\|xi\|^2 = \|x\|^2 \geq X^2$ .
    - ii. Show that  $\operatorname{re}(xi) = \operatorname{re}(-\operatorname{im}(x) + \operatorname{re}(x)i) = -\operatorname{im}(x) > 0$ .
    - iii. Hence show that  $\|\ln_k(xi)\|^2 \leq a_1^2$  using procedure  $p_1$ .
    - iv. **Hence show that**  $\|\ln_k(x)\|^2 = \|\ln_k(xi) - \frac{\tau_k}{4}i\|^2 \leq (a_1 + \frac{a_2}{4})^2 = a^2$ .
6. **Yield the tuple**  $\langle a, b, p \rangle$ .

## Procedure IV:28(wed2108191401)

### Objective

Choose a rational number  $X$  such that  $1 \geq X > 0$ . The objective of the following instructions is to construct positive rational numbers  $a, c, d, e$  and a procedure  $p(x, n, k)$  to show that  $\exp_n(\ln_k(x)) \equiv x \pmod{\frac{an}{k} + \frac{c}{n}}$  when a complex number  $x$  and integers  $k, n$  such that  $\operatorname{re}(x) \geq 0$ ,  $\|x\|^2 \geq X^2$ ,  $n > e$ , and  $k > d$  are chosen.

### Implementation

1. Execute **procedure IV:25** on  $\langle X \rangle$  and let  $\langle f, p_0 \rangle$  receive.
2. Execute **procedure III:70** on  $\langle f \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
3. Execute **procedure III:37** on  $\langle a_1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
4. Execute **procedure III:35** on  $\langle a_1 \rangle$  and let  $\langle a_3, b_3, p_3 \rangle$  receive.
5. Execute **procedure III:71** on  $\langle f \rangle$  and let  $\langle a_4, c_4, d, e_4, p_4 \rangle$  receive.
6. Let  $a = \frac{a_4}{a_3} + \frac{a_4(1+f)}{a_3(1-f)}$ .
7. Let  $c = a_2 + \frac{c_4}{a_3} + \frac{c_4(1+f)}{a_3(1-f)}$ .

8. Let  $e = \max(b_2, b_3, e_4)$ .
  9. Let  $p(x, n, k)$  be the following procedure:
    - (a) Show that  $\operatorname{re}(\frac{x-1}{x+1})^2 \leq \|\frac{x-1}{x+1}\|^2 \leq f^2$  using procedure  $p_0$ .
    - (b) Hence show that  $|\operatorname{re}(\frac{x-1}{x+1})| \leq f$ .
    - (c) Show that  $\|\ln_k(1 + \frac{x-1}{x+1})\|^2 \leq a_1^2$  using procedure  $p_1$ .
    - (d) Show that  $\|\ln_k(1 - \frac{x-1}{x+1})\|^2 \leq a_1^2$  using procedure  $p_1$ .
    - (e) Hence using procedures  $p_0, p_2, p_3, p_4$ , show that  $\exp_n(\ln_k(x))$ 
      - i.  $= \exp_n(\ln_k(1 + \frac{x-1}{x+1}) - \ln_k(1 - \frac{x-1}{x+1}))$
      - ii.  $\equiv \frac{\exp_n(\ln_k(1 + \frac{x-1}{x+1}))}{\exp_n(\ln_k(1 - \frac{x-1}{x+1}))} \pmod{\frac{a_2}{n}}$
      - iii.  $\equiv \frac{1 + \frac{x-1}{x+1}}{\exp_n(\ln_k(1 - \frac{x-1}{x+1}))} \pmod{\frac{1}{a_3}(\frac{a_4n}{k} + \frac{c_4}{n})}$
      - iv.  $\equiv \frac{1 + \frac{x-1}{x+1}}{1 - \frac{x-1}{x+1}} \pmod{(\frac{1+f}{a_3(1-f)})(\frac{a_4n}{k} + \frac{c_4}{n})}$
      - v.  $= x$ .
    - (f) Hence show that  $\exp_n(\ln_k(x)) \equiv x$ 
      - i.  $(\operatorname{err} \frac{a_2}{n} + \frac{1}{a_3}(\frac{a_4n}{k} + \frac{c_4}{n}) + (\frac{1+f}{a_3(1-f)})(\frac{a_4n}{k} + \frac{c_4}{n}))$
      - ii.  $(\operatorname{err} \frac{an}{k} + \frac{c}{n})$ .
10. **Yield the tuple**  $\langle a, c, d, e, p \rangle$ .

## Procedure IV:29(sun0812191512)

### Objective

Choose a rational number  $X$  such that  $0 < X \leq 1$ . The objective of the following instructions is to construct positive rational numbers  $a, c, d, e$  and a procedure  $p(x, n, k)$  to show that  $\exp_n(\ln_k(x)) \equiv x \pmod{\frac{an}{k} + \frac{c}{n}}$  when a complex number  $x$  and integers  $k, n$  such that  $\|x\|^2 \geq X^2$ ,  $n > e$ , and  $k > d$  are chosen.

## Implementation

1. Execute **procedure IV:28** on  $\langle X \rangle$  and let  $\langle a_1, b_1, c_1, d_1, p_1 \rangle$  receive.
2. Execute **procedure IV:27** on  $\langle X \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:82** on  $\langle a_2, 1 \rangle$  and let  $\langle a_3, b_3, c_3, d_3, p_3 \rangle$  receive.
4. Let  $a = a_2 + a_3$ .
5. Let  $c = b_1 + b_3$ .
6. Let  $d = \max(c_1, b_2, d_3)$ .
7. Let  $e = \max(c_3, d_1)$ .
8. Let  $p(x, n, k)$  be the following procedure:
  - (a) If  $\text{re}(x) \geq 0$ , then do the following:
    - i. **Show that**  $\exp_n(\ln_k(x)) \equiv x \left( \text{err } \frac{a_2 n}{k} + \frac{b_1}{n} \right) \left( \text{err } \frac{a_3 n}{k} + \frac{c}{n} \right)$  **using procedure**  $p_1$ .
  - (b) Otherwise if  $\text{im}(x) \geq 0$ , then do the following:
    - i. Show that  $\left\| \frac{x}{i} \right\|^2 = \|x\|^2 \geq X^2$ .
    - ii. Show that  $\text{re}(\frac{x}{i}) = \text{re}(\text{im}(x) - \text{re}(x)i) = \text{im}(x) \geq 0$ .
    - iii. Hence show that  $\exp_n(\ln_k(\frac{x}{i})) \equiv \frac{x}{i} \left( \text{err } \frac{a_2 n}{k} + \frac{b_1}{n} \right)$  using procedure  $p_1$ .
    - iv. Hence show that  $\|\ln_k(\frac{x}{i})\|^2 \leq a_2^2$  using procedure  $p_2$ .
    - v. Hence using procedure  $p_3$ , show that  $\exp_n(\ln_k(x))$ 
      - A.  $= \exp_n(\ln_k(\frac{x}{i}) + \frac{\tau_k}{4}i)$
      - B.  $\equiv i^1 \exp_n(\ln_k(\frac{x}{i})) \left( \text{err } \frac{a_3 n}{k} + \frac{b_3}{n} \right)$
      - C.  $\equiv i \cdot \frac{x}{i} \left( \text{err } \frac{a_2 n}{k} + \frac{b_1}{n} \right)$
      - D.  $= x$ .
    - vi. **Hence show that**  $\exp_n(\ln_k(x)) \equiv x \left( \text{err } \frac{a_3 n}{k} + \frac{c}{n} \right)$ .
  - (c) Otherwise do the following:
    - i. Show that  $\|xi\|^2 = \|x\|^2 \geq X^2$ .
    - ii. Show that  $\text{re}(xi) = \text{re}(-\text{im}(x) + \text{re}(x)i) = -\text{im}(x) > 0$ .
    - iii. Hence show that  $\exp_n(\ln_k(xi)) \equiv xi \left( \text{err } \frac{a_2 n}{k} + \frac{b_1}{n} \right)$  using procedure  $p_1$ .

iv. Hence show that  $\|\ln_k(xi)\|^2 \leq a_2^2$  using procedure  $p_2$ .

v. Hence using procedure  $p_3$ , show that  $\exp_n(\ln_k(x))$

$$A. = \exp_n(\ln_k(xi) - \frac{\tau_k}{4}i)$$

$$B. \equiv i^{-1} \exp_n(\ln_k(xi)) \left( \text{err } \frac{a_3 n}{k} + \frac{b_3}{n} \right)$$

$$C. \equiv i^{-1}xi \left( \text{err } \frac{a_2 n}{k} + \frac{b_1}{n} \right)$$

$$D. = x.$$

vi. **Hence show that**  $\exp_n(\ln_k(x)) \equiv x \left( \text{err } \frac{a_3 n}{k} + \frac{c}{n} \right)$ .

9. **Yield the tuple**  $\langle a, c, d, e, p \rangle$ .

## Procedure IV:30(wed2108191324)

### Objective

Choose two rational numbers  $X, \epsilon$  such that  $0 < \epsilon < 1$  and  $X \geq 0$ . The objective of the following instructions is to construct a rational number  $f$  such that  $0 < f < 1$ , and a procedure  $p(x)$  to show that  $\left\| \frac{x-1}{x+1} \right\|^2 \leq f^2$  when a complex number  $x$  such that  $\text{re}(x) \geq \epsilon$  and  $\|x\|^2 \leq X^2$  is chosen.

### Implementation

1. Let  $f = 1 - \frac{2\epsilon}{X^2+1+2\epsilon}$ .

2. Show that  $0 < f < 1$ .

3. Let  $p(x)$  be the following procedure:

(a) Show that  $X^2 + 1 + 2\epsilon \leq \frac{4\epsilon}{1-f^2}$

i. given that  $\frac{4\epsilon}{X^2+1+2\epsilon} \geq 1 - f^2$

ii. given that  $f^2 = 1 - \frac{4\epsilon}{X^2+1+2\epsilon} + \left( \frac{2\epsilon}{X^2+1+2\epsilon} \right)^2 \geq 1 - \frac{4\epsilon}{X^2+1+2\epsilon}$ .

(b) Hence show that  $\|x\|^2$

i.  $\leq X^2$

ii.  $\leq \frac{4\epsilon}{1-f^2} - (1 + 2\epsilon)$

iii.  $= \frac{4\epsilon-1+f^2-2\epsilon+2\epsilon f^2}{1-f^2}$

iv.  $= \frac{f^2-1+2\epsilon(1+f^2)}{1-f^2}$

v.  $\leq \frac{f^2-1+2\text{re}(x)(1+f^2)}{1-f^2}$ .

- (c) **Hence show that**  $\|\frac{x-1}{x+1}\|^2 \leq f^2$
- given that  $(\text{re}(x) - 1)^2 + \text{im}(x)^2 \leq f^2((\text{re}(x) + 1)^2 + \text{im}(x)^2)$
  - given that  $(1 - f^2)(\text{re}(x)^2 + \text{im}(x)^2) \leq f^2 - 1 + 2\text{re}(x)(1 + f^2)$ .

4. **Yield the tuple**  $\langle f, p \rangle$ .

## Procedure IV:31(wed2108191603)

### Objective

Choose two rational numbers  $X, \epsilon$  such that  $0 < \epsilon < 1$  and  $X \geq 0$ . The objective of the following instructions is to construct rational numbers  $c, d, a, b, e$ , a procedure  $p(x, n)$  to show that  $\|\frac{1}{x}\|^2 \leq c^2$  when a complex number  $x$  and a positive integer  $n$  such that  $\text{re}(x) \geq \epsilon$ ,  $\|x\|^2 \leq X^2$ , and  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} \ln_n(y) \equiv \frac{1}{x} (\text{err } \frac{a}{n} + b\{\delta\})$  when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq e^2$  is chosen.

### Implementation

- Execute the following in post-order:
  - Execute **procedure IV:10** on  $\langle q_3, q_4, q_5, q_6 \rangle$  and let  $\langle c, d, a, b, e, q_1, q_2 \rangle$  receive.
  - Execute **procedure IV:11** on  $\langle q_7, q_8, q_9, q_{10}, q_{11} \rangle$  and let  $\langle q_3, q_4 \rangle$  receive.
- Execute **procedure IV:24** on  $\langle e_1, \frac{1-e_1}{2} \rangle$  and let  $\langle q_7, q_8 \rangle$  receive.
  - Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle e_1, q_{12} \rangle$  receive.
- Execute **procedure IV:10** on  $\langle q_{13}, q_{14}, q_{15}, q_{16} \rangle$  and let  $\langle q_9, q_{10} \rangle$  receive.
  - Execute **procedure IV:15** on  $\langle 1, 1 \rangle$  and let  $\langle q_{13}, q_{14} \rangle$  receive.
  - Execute **procedure IV:12** on  $\langle -2, q_{17}, q_{18} \rangle$  and let  $\langle q_{15}, q_{16} \rangle$  receive.
- Execute **procedure IV:11** on  $\langle q_{19}, q_{20}, q_{21}, q_{22}, q_{23} \rangle$  and let  $\langle q_{17}, q_{18} \rangle$  receive.

- Execute **procedure IV:18** on  $\langle 1, 1 + \epsilon, X + 1 \rangle$  and let  $\langle q_{19}, q_{20} \rangle$  receive.
- Execute **procedure IV:10** on  $\langle q_{24}, q_{25}, q_{26}, q_{27} \rangle$  and let  $\langle q_{21}, q_{22} \rangle$  receive.
  - Execute **procedure IV:15** on  $\langle 1, 1 \rangle$  and let  $\langle q_{24}, q_{25} \rangle$  receive.
  - Execute **procedure IV:16** on  $\langle X, 1 \rangle$  and let  $\langle q_{26}, q_{27} \rangle$  receive.
- Let  $q_{24}(x, n)$  be the following procedure:
  - Show that  $(1 + \epsilon)^2 \leq (1 + \text{re}(x))^2 = \text{re}(x+1)^2 \leq \|x+1\|^2 \leq (X+1)^2$ .
- Let  $q_{11}(x, n)$  be the following procedure:
  - Show that  $\|1 - 2(x+1)^{-1}\|^2 = \|\frac{x-1}{x+1}\|^2 \leq e_1^2$  using procedure  $q_{12}$ .
- Execute **procedure IV:12** on  $\langle -1, q_{28}, q_{29} \rangle$  and let  $\langle q_5, q_6 \rangle$  receive.
- Execute **procedure IV:11** on  $\langle q_{30}, q_{31}, q_{32}, q_{33}, q_{34} \rangle$  and let  $\langle q_{28}, q_{29} \rangle$  receive.
  - Execute **procedure IV:24** on  $\langle e_1, \frac{1-e_1}{2} \rangle$  and let  $\langle q_{30}, q_{31} \rangle$  receive.
  - Execute **procedure IV:12** on  $\langle -1, q_9, q_{10} \rangle$  and let  $\langle q_{32}, q_{33} \rangle$  receive.
  - Let  $q_{34}(x, n)$  be the following procedure:
    - Show that  $\|-(1+(-2)(x+1)^{-1})\|^2 = \|\frac{x-1}{x+1}\|^2 \leq e_1^2$  using procedure  $q_{12}$ .
- Let  $p(x, n)$  be the following procedure:
  - Show that**  $\frac{1}{x} \equiv 0 \text{ (err } \frac{1}{x}) \text{ (err } \frac{1}{\text{re}(x)}) \text{ (err } \frac{1}{\epsilon}) \text{ (err } c)$ .
- Let  $q(x, n, \delta)$  be the following procedure:
  - Using procedure  $q_2$ , show that  $\Delta_{y=x}^{+\delta} \ln_n(y)$ 
    - $= \Delta_{y=x}^{+\delta} (\ln_n(1 + \frac{y-1}{y+1}) - \ln_n(1 - \frac{y-1}{y+1}))$
    - $= \Delta_{y=x}^{+\delta} (\ln_n(1 + (1 + (-2)(y+1)^{-1})) + (-1) \ln_n(1 + (-1)(1 + (-2)(y+1)^{-1})))$
    - $\equiv ((1 + (1 + (-2)(x+1)^{-1}))^{-1} (0 + (-2)(-1)(x+1)^{-2} (1+0) + (-1)((1 + (-1)(1 + (-2)(x+1)^{-1}))^{-1} \cdot (0 + (-1)(0 +$

$$(-2)(-1)(x+1)^{-2}(1+0)))))) (\text{err } \frac{a}{n} + b\{\delta\})$$

$$4. = \frac{1}{x}$$

$$b) \text{ Hence show that } \Delta_{y=x}^{+\delta} \ln_n(y) \equiv \frac{1}{x} (\text{err } \frac{a}{n} + b\{\delta\}).$$

4) Yield the tuple  $\langle c, d, a, b, e, p, q \rangle$ .

## Procedure IV:32(thu2208191330)

### Objective

Choose a complex number  $A$  and non-negative rational numbers  $X, D$  such that  $X + D < 1$ . The objective of the following instructions is to construct rational numbers  $l, d, a, b$ , a procedure  $p(x, n)$  to show that  $\|A(1+x)^{A-1}\|^2 \leq l^2$  when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$  and  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta}(1+y)_n^A \equiv A(1+x)_{n-1}^{A-1} (\text{err } \frac{a}{n} + b\{\delta\})$  when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.

### Implementation

1. Execute **procedure III:52** on  $\langle \{A+1\}, X+D \rangle$  and let  $\langle a_1, b_1, p_1 \rangle$  receive.
2. Execute **procedure III:55** on  $\langle \{A-1\}, X \rangle$  and let  $\langle a_2, p_2 \rangle$  receive.
3. Let  $l = \{A\}a_2$ .
4. Let  $d = 1$ .
5. Let  $a = 0$ .
6. Let  $b = \frac{a_1 b_1}{D^2(1-b_1)}$ .
7. Let  $p(x, n)$  be the following procedure:
  - (a) Using procedure  $p_2$ , show that  $A(1+x)_{n-1}^{A-1} \equiv 0$ 
    - i.  $(\text{err } A(1+x)_{n-1}^{A-1})$
    - ii.  $(\text{err } \{A\}a_2)$
    - iii.  $(\text{err } l)$ .
8. Let  $q(x, n, \delta)$  be the following procedure:
  - (a) For  $r \in [1 : n]$ , do the following:
    - i. Show that  $\|A+1\|^2 \leq \{A+1\}^2$ .

$$\text{ii. Hence show that } \|(A_r)(X+D)^r\|^2 \leq (a_1 b_1^r)^2 \text{ using procedure } p_1.$$

$$\text{iii. Hence show that } \|(A_r) \sum_k^{[0:r-1]} \binom{r}{k} x^k \delta^{r-2-k}\|^2$$

$$A. \leq \|(A_r)\|^2 (\sum_k^{[0:r-1]} \binom{r}{k} X^k D^{r-2-k})^2$$

$$B. = \|(A_r)\|^2 (\frac{1}{D^2} \sum_k^{[0:r-1]} \binom{r}{k} X^k D^{r-k})^2$$

$$C. \leq \|\frac{1}{D^2} (A_r)(X+D)^r\|^2$$

$$D. \leq (\frac{a_1 b_1^r}{D^2})^2.$$

$$(b) \text{ Now show that } \Delta_{y=x}^{+\delta}(1+y)_n^A \equiv A(1+x)_{n-1}^{A-1}$$

$$\text{i. } (\text{err } \frac{1}{\delta} ((1+x+\delta)_n^A - (1+x)_n^A) - A(1+x)_{n-1}^{A-1})$$

$$\text{ii. } (\text{err } \frac{1}{\delta} (\sum_r^{[0:n]} \binom{A}{r} (x+\delta)^r - \sum_r^{[0:n]} \binom{A}{r} x^r) - A \sum_r^{[0:n-1]} \binom{A-1}{r} x^r)$$

$$\text{iii. } (\text{err } \frac{1}{\delta} (\sum_r^{[0:n]} \binom{A}{r} (\sum_k^{[0:r+1]} \binom{r}{k} x^k \delta^{r-k} - x^r)) - A \sum_r^{[0:n-1]} \binom{A-1}{r} x^r)$$

$$\text{iv. } (\text{err } \sum_r^{[0:n]} \binom{A}{r} \sum_k^{[0:r]} \binom{r}{k} x^k \delta^{r-1-k} - \sum_r^{[0:n-1]} (r+1) \binom{A}{r+1} x^r)$$

$$\text{v. } (\text{err } \sum_r^{[1:n]} \binom{A}{r} \sum_k^{[0:r]} \binom{r}{k} x^k \delta^{r-1-k} - \sum_r^{[1:n]} r \binom{A}{r} x^{r-1})$$

$$\text{vi. } (\text{err } \delta (\sum_r^{[1:n]} \binom{A}{r} \sum_k^{[0:r-1]} \binom{r}{k} x^k \delta^{r-2-k}))$$

$$\text{vii. } (\text{err } \delta (\sum_r^{[1:n]} \frac{a_1 b_1^r}{D^2}))$$

$$\text{viii. } (\text{err } \delta (\frac{a_1 b_1}{D^2(1-b_1)}))$$

$$\text{ix. } (\text{err } \frac{a}{n} + b\{\delta\}).$$

9. Yield the tuple  $\langle l, d, a, b, p, q \rangle$ .

## Procedure IV:33(thu2208191432)

### Objective

Choose a complex number  $A$  and non-negative rational numbers  $X, D$  such that  $X + D < 1$ . The objective of the following instructions is to construct rational numbers  $l, d, a, b$ , a procedure  $p(x, n)$  to show that  $\|A(1+x)^{A-1}\|^2 \leq l^2$  when a complex number  $x$  and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$  and  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta}(1+y)_n^A \equiv A(1+x)_{n-1}^{A-1} (\text{err } \frac{a}{n} + b\{\delta\})$  when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq D^2$  is chosen.



## Implementation

1. Execute **procedure III:52** on  $\langle \{A\}, X \rangle$  and let  $\langle a_1, b_1, p_1 \rangle$  receive.
2. Execute **procedure III:55** on  $\langle \{A-1\}, X \rangle$  and let  $\langle a_2, p_2 \rangle$  receive.
3. Execute **procedure III:53** on  $\langle b_1, 1 \rangle$  and let  $\langle a_3, b_3, p_3 \rangle$  receive.
4. Execute **procedure IV:32** on  $\langle X, D \rangle$  and let  $\langle a_4, b_4, c_4, d_4, p_4, q_4 \rangle$  receive.
5. Let  $l = \{a\}a_2$ .
6. Let  $d = \max(1, b_4)$ .
7. Let  $a = c_4 + 2\{A\}a_1a_3$ .
8. Let  $b = d_4$ .
9. Let  $p(x, n)$  be the following procedure:
  - (a) Using procedure  $p_2$ , show that  $A(1+x)_n^{A-1} \equiv 0$ 
    - i. (err  $A(1+x)_n^{A-1}$ )
    - ii. (err  $\{A\}a_2$ )
    - iii. (err  $l$ ).
10. Let  $q(x, n, \delta)$  be the following procedure:
  - (a) Show that  $\|(A-1)x^{n-1}\|^2 \leq (a_1b_1)^{n-1}$  using procedure  $p_1$ .
  - (b) Show that  $\|(n-1)b_1^{n-1}\|^2 \leq (a_3b_3)^{n-1}$  using procedure  $p_3$ .
  - (c) Hence show that  $\|b_1^{n-1}\|^2 \leq (\frac{a_3}{n-1})^2 \leq (\frac{2a_3}{n})^2$ .
  - (d) Now using procedure  $q_4$ , show that  $\Delta_{y=x}^{+\delta}(1+y)_n^A$ 
    - i.  $\equiv A(1+x)_{n-1}^{A-1}$  (err  $\frac{c_4}{n} + d_4\{\delta\}$ )
    - ii.  $\equiv A(1+x)_n^{A-1}$ 
      - A. (err  $A(\frac{A-1}{n-1})x^{n-1}$ )
      - B. (err  $\{A\}a_1b_1^{n-1}$ )
      - C. (err  $\frac{2\{A\}a_1a_3}{n}$ ).
  - (e) Hence show that  $\Delta_{y=x}^{+\delta}(1+y)_n^A \equiv A(1+x)_n^{A-1}$ 
    - i. (err  $\frac{c_4}{n} + d_4\{\delta\} + \frac{2\{A\}a_1a_3}{n}$ )
    - ii. (err  $\frac{a}{n} + b\{\delta\}$ ).

## 11. Yield the tuple $\langle l, d, a, b, p, q \rangle$ .

### Declaration IV:3

The notation  $x_n^a$ , where  $x, a$  are complex numbers and  $n$  is a positive integer, will be used as a shorthand for  $(1 + \frac{x-1}{x+1})_n^a (1 - \frac{x-1}{x+1})_n^{-a}$ .

### Procedure IV:34(sat2408190819)

#### Objective

Choose three non-negative rational numbers  $A, X, \epsilon$  such that  $0 < \epsilon < 1$ . The objective of the following instructions is to construct positive rational numbers  $B, C, D$ , and a procedure  $p(x, a, n)$  to show that  $x_n^a \equiv x^a$  (err  $BC^n$ ) when a complex number  $x$ , and integers  $a, n$  such that  $\|x\|^2 \leq X^2$ ,  $\|a\|^2 \leq A^2$ ,  $\text{re}(x) \geq \epsilon$ , and  $n > D$  are chosen.

#### Implementation

1. Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Show that  $0 < a_1 < 1$ .
3. Execute **procedure III:57** on  $\langle A, a_1 \rangle$  and let  $\langle a_2, b_2, c_2, p_2 \rangle$  receive.
4. Execute **procedure III:55** on  $\langle A, a_1 \rangle$  and let  $\langle a_3, p_3 \rangle$  receive.
5. Let  $B = a_3a_2$ .
6. Let  $C = b_2$ .
7. Let  $D = \max(c_2, A)$ .
8. Let  $p(x, a, n)$  be the following procedure:
  - (a) Show that  $\|\frac{x-1}{x+1}\|^2 \leq a_1^2$  using procedure  $p_1$ .
  - (b) If  $a \geq 0$ , then do the following:
    - i. Using **procedure III:49** and procedures  $p_2, p_3$ , show that  $x_n^a$ 
      - A.  $= (1 + \frac{x-1}{x+1})_n^a (1 - \frac{x-1}{x+1})_n^{0-a}$
      - B.  $\equiv (1 + \frac{x-1}{x+1})_n^a \frac{(1 - \frac{x-1}{x+1})_n^0}{(1 - \frac{x-1}{x+1})_n^a}$  (err  $a_3a_2b_2^n$ )
      - C.  $= \frac{(1 + \frac{x-1}{x+1})_n^a}{(1 - \frac{x-1}{x+1})_n^a}$
      - D.  $= x^a$



(c) Otherwise do the following:

- i. Using **procedure III:49** and procedures  $p_2, p_3$ , show that  $x_n^a$

$$A. = (1 + \frac{x-1}{x+1})_n^{0-(-a)} (1 - \frac{x-1}{x+1})_n^{-a}$$

$$B. \equiv \frac{(1 + \frac{x-1}{x+1})_n^0}{(1 + \frac{x-1}{x+1})_n^{-a}} (1 - \frac{x-1}{x+1})_n^{-a} \text{ (err } a_3 a_2 b_2^n \text{)}$$

$$C. = \frac{(1 - \frac{x-1}{x+1})_n^{-a}}{(1 + \frac{x-1}{x+1})_n^{-a}}$$

$$D. = (\frac{1}{x})^{-a}$$

$$E. = x^a.$$

- (d) **Hence show that**  $x_n^a \equiv x^a \text{ (err } a_3 a_2 b_2^n \text{)}$   
(err  $BC^n$ ).

9. **Yield the tuple**  $\langle B, C, D, p \rangle$ .

## Procedure IV:35(sat2408191109)

### Objective

Choose three non-negative rational numbers  $A, X, \epsilon$  such that  $0 < \epsilon < 1$ . The objective of the following instructions is to construct positive rational numbers  $B, C$ , and a procedure  $p(x, a, b, n)$  to show that  $x_n^{a+b} \equiv x_n^a x_n^b \text{ (err } BC^n \text{)}$  when complex numbers  $x, a, b$ , and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$ ,  $\|a\|^2 \leq A^2$ ,  $\|b\|^2 \leq A^2$ , and  $\text{re}(x) \geq \epsilon$  are chosen.

### Implementation

1. Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Execute **procedure III:54** on  $\langle A, a_1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:55** on  $\langle 2A, a_1 \rangle$  and let  $\langle a_3, p_3 \rangle$  receive.
4. Let  $B = a_2 a_3 (1 + a_3)$ .
5. Let  $C = b_2$ .
6. Let  $p(x, a, b, n)$  be the following procedure:
  - (a) Using procedures  $p_1, p_2, p_3$ , show that  $x_n^{a+b}$ 
    - i.  $= (1 + \frac{x-1}{x+1})_n^{a+b} (1 - \frac{x-1}{x+1})_n^{-(a+b)}$
    - ii.  $\equiv (1 + \frac{x-1}{x+1})_n^a (1 + \frac{x-1}{x+1})_n^b (1 - \frac{x-1}{x+1})_n^{(-a)+(-b)} \text{ (err } a_2 b_2^n a_3 \text{)}$

$$\text{iii.} \equiv (1 + \frac{x-1}{x+1})_n^a (1 + \frac{x-1}{x+1})_n^b (1 - \frac{x-1}{x+1})_n^{-a} (1 - \frac{x-1}{x+1})_n^{-b} \text{ (err } a_3^2 a_2 b_2^n \text{)}$$

$$\text{iv.} = x_n^a x_n^b.$$

- (b) **Hence show that**  $x_n^{a+b} \equiv x_n^a x_n^b \text{ (err } BC^n \text{)}$ .

7. **Yield the tuple**  $\langle B, C, p \rangle$ .

## Procedure IV:36(sat2408191137)

### Objective

Choose three non-negative rational numbers  $A, X, \epsilon$  such that  $0 < \epsilon < 1$ . The objective of the following instructions is to construct a positive rational number  $D$ , and a procedure  $p(x, n, a, k)$  to show that  $(x_n^a)^k \equiv 0 \text{ (err } D \text{)}$  when complex numbers  $x, a, k$ , and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$ ,  $\|ka\|^2 \leq A^2$ , and  $\text{re}(x) \geq \epsilon$  are chosen.

### Implementation

1. Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Execute **procedure III:55** on  $\langle A, a_1 \rangle$  and let  $\langle a_2, p_2 \rangle$  receive.
3. Let  $D = a_2^2$ .
4. Let  $p(x, n, a, k)$  be the following procedure:
  - (a) Using procedures  $p_1, p_2$ , show that  $(x_n^a)^k$ 
    - i.  $= ((1 + \frac{x-1}{x+1})_n^a (1 - \frac{x-1}{x+1})_n^{-a})^k$
    - ii.  $= ((1 + \frac{x-1}{x+1})_n^a)^k ((1 - \frac{x-1}{x+1})_n^{-a})^k$
    - iii.  $\equiv 0 ((1 - \frac{x-1}{x+1})_n^{-a})^k \text{ (err } a_2^2 \text{)}$
    - iv.  $= 0$ .
  - (b) **Hence show that**  $(x_n^a)^k \equiv 0 \text{ (err } D \text{)}$ .
5. **Yield the tuple**  $\langle D, p \rangle$ .

## Procedure IV:37(thu2908190744)

### Objective

Choose three non-negative rational numbers  $A, X, \epsilon$  such that  $0 < \epsilon < 1$ . The objective of the following instructions is to construct a positive rational number  $D$ , and a procedure  $p(x, n, a)$  to show that  $\|x_n^a\|^2 \geq D^2$  when complex numbers  $x, a$ , and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$ ,  $\|a\|^2 \leq A^2$ , and  $\text{re}(x) \geq \epsilon$  are chosen.

### Implementation

1. Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Execute **procedure III:56** on  $\langle A, a_1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Let  $D = a_2^2$ .
4. Let  $p(x, n, a)$  be the following procedure:
  - (a) Show that  $\|\frac{x-1}{x+1}\|^2 \leq a_1^2$  using procedure  $p_1$ .
  - (b) Show that  $\|(1 + \frac{x-1}{x+1})_n^a\|^2 \geq a_2^2$  using procedure  $p_2$ .
  - (c) Show that  $\|(1 - \frac{x-1}{x+1})_n^{-a}\|^2 \geq a_2^2$  using procedure  $p_2$ .
  - (d) Hence using **declaration IV:3**, show that  $\|x_n^a\|^2$ 
    - i.  $= \|(1 + \frac{x-1}{x+1})_n^a\|^2 \|(1 - \frac{x-1}{x+1})_n^{-a}\|^2$
    - ii.  $\geq a_2^2 a_2^2$
    - iii.  $= D^2$ .
5. Yield the tuple  $\langle D, p \rangle$ .

## Procedure IV:38(thu2908190802)

### Objective

Choose three non-negative rational numbers  $A, X, \epsilon$  such that  $0 < \epsilon < 1$ . The objective of the following instructions is to construct positive rational numbers  $B, C, D$ , and a procedure  $p(x, a, b, n)$  to show that  $x_n^{a-b} \equiv \frac{x_n^a}{x_n^b} \pmod{BC^n}$  when complex numbers  $x, a, b$ , and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$ ,  $\|a\|^2 \leq A^2$ ,  $\|b\|^2 \leq A^2$ ,  $\text{re}(x) \geq \epsilon$ , and  $n > D$  are chosen.

### Implementation

1. Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Execute **procedure III:57** on  $\langle A, a_1 \rangle$  and let  $\langle a_2, b_2, c_2, p_2 \rangle$  receive.
3. Execute **procedure III:56** on  $\langle A, a_1 \rangle$  and let  $\langle a_3, b_3, p_3 \rangle$  receive.
4. Execute **procedure III:55** on  $\langle 2A, a_1 \rangle$  and let  $\langle a_4, p_4 \rangle$  receive.
5. Let  $B = a_2 a_4 (1 + \frac{1}{a_3})$ .
6. Let  $C = b_2$ .
7. Let  $D = \max(c_2, b_3)$ .
8. Let  $p(x, a, b, n)$  be the following procedure:
  - (a) Using procedures  $p_1, p_2, p_3, p_4$ , show that  $x_n^{a-b}$ 
    - i.  $= (1 + \frac{x-1}{x+1})_n^{a-b} (1 - \frac{x-1}{x+1})_n^{(-a)-(-b)}$
    - ii.  $\equiv (1 + \frac{x-1}{x+1})_n^{a-b} \frac{(1 - \frac{x-1}{x+1})_n^{-a}}{(1 - \frac{x-1}{x+1})_n^{-b}} \pmod{a_4 a_2 b_2^n}$
    - iii.  $\equiv \frac{(1 + \frac{x-1}{x+1})_n^a (1 - \frac{x-1}{x+1})_n^{-a}}{(1 + \frac{x-1}{x+1})_n^b (1 - \frac{x-1}{x+1})_n^{-b}} \pmod{a_2 b_2^n \frac{a_4}{a_3}}$
    - iv.  $= \frac{x_n^a}{x_n^b}$ .
  - (b) Hence show that  $x_n^{a-b} \equiv \frac{x_n^a}{x_n^b} \pmod{a_4 a_2 b_2^n + a_2 b_2^n \frac{a_4}{a_3}} \pmod{BC^n}$ .
9. Yield the tuple  $\langle B, C, D, p \rangle$ .

## Procedure IV:39(sat2408191538)

### Objective

Choose three non-negative rational numbers  $A, X, \epsilon$  such that  $0 < \epsilon < 1$ . The objective of the following instructions is to construct a positive rational number  $B, C$ , and a procedure  $p(x, n, a, k)$  to show that  $(x_n^a)^k \equiv x_n^{ak} \pmod{BC^n}$  when complex numbers  $x, a, k$ , and a positive integer  $n$  such that  $\|x\|^2 \leq X^2$ ,  $\|ka\|^2 \leq A^2$ , and  $\text{re}(x) \geq \epsilon$  are chosen.

## Implementation

1. Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
2. Execute **procedure III:58** on  $\langle A, a_1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.
3. Execute **procedure III:55** on  $\langle A, a_1 \rangle$  and let  $\langle a_3, p_3 \rangle$  receive.
4. Let  $B = 2a_2a_3$ .
5. Let  $C = b_2$ .
6. Let  $p(x, n, a, k)$  be the following procedure:
  - (a) Using procedures  $p_1, p_2, p_3$ , show that  $(x_n^a)^k$ 
    - i.  $= ((1 + \frac{x-1}{x+1})_n^a (1 - \frac{x-1}{x+1})_n^{-a})^k$
    - ii.  $= ((1 + \frac{x-1}{x+1})_n^a ((1 - \frac{x-1}{x+1})_n^{-a})^k$
    - iii.  $\equiv (1 + \frac{x-1}{x+1})_n^{ak} ((1 - \frac{x-1}{x+1})_n^{-a})^k$  (err  $a_2b_2^n a_3$ )
    - iv.  $\equiv ((1 + \frac{x-1}{x+1})_n^{ak})^1 (1 - \frac{x-1}{x+1})_n^{-ak}$  (err  $a_3a_2b_2^n$ )
    - v.  $= x_n^{ak}$ .
  - (b) **Hence show that**  $(x_n^a)^k \equiv x_n^{ak}$  (err  $BC^n$ ).
7. Yield the tuple  $\langle B, C, p \rangle$ .

## Procedure IV:40(thu2208191610)

### Objective

Choose a complex number  $f$  and two rational numbers  $X, \epsilon$  such that  $0 < \epsilon < 1$  and  $X \geq 0$ . Let  $g(f, x, n)$  be a shorthand for  $[2f(1 + \frac{x-1}{x+1})_n^f \cdot (1 - \frac{x-1}{x+1})_n^{-f-1}(x+1)^{-2}] + [2f(1 + \frac{x-1}{x+1})_n^{f-1} \cdot (1 - \frac{x-1}{x+1})_n^{-f}(x+1)^{-2}]$ . The objective of the following instructions is to construct rational numbers  $c, d, a, b, e$ , a procedure  $p(x, n)$  to show that  $\|g(f, x, n)\|^2 \leq c^2$  when a complex number  $x$  and a positive integer  $n$  such that  $\text{re}(x) \geq \epsilon$ ,  $\|x\|^2$ , and  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} x_n^f \equiv g(f, x, n)$  (err  $\frac{a}{n} + b\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq e^2$  is chosen.

## Implementation

- 1) Execute the following in post-order:
  - a) Execute **procedure IV:13** on  $\langle q_7, q_3, q_4, q_8, q_5, q_6 \rangle$  and let  $\langle c, d, a, b, e, p, q \rangle$  receive.

- i) Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle e_1, q_9 \rangle$  receive.
- ii) Execute **procedure III:55** on  $\langle \{f\}, e_1 \rangle$  and let  $\langle e_2, q_{10} \rangle$  receive.
- iii) Let  $q_7(x, n)$  be the following procedure:
  - (1) Show that  $\|\frac{x-1}{x+1}\|^2 \leq e_1^2$  using procedure  $q_9$ .
  - (2) Using procedure  $q_{10}$ , show that  $\|(1 + (1 + (-2)(x+1)^{-1}))_n^f\|^2$ 
    - (a)  $= \|(1 + \frac{x-1}{x+1})_n^f\|^2$
    - (b)  $\leq e_2^2$ .
- iv) Let  $q_8(x, n)$  be the following procedure:
  - (1) Show that  $\|\frac{x-1}{x+1}\|^2 \leq e_1^2$  using procedure  $q_9$ .
  - (2) Using procedure  $q_{10}$ , show that  $\|(1 - (1 + (-2)(x+1)^{-1}))_n^f\|^2$ 
    - (a)  $= \|(1 - \frac{x-1}{x+1})_n^f\|^2$
    - (b)  $\leq e_2^2$ .
- v) Execute **procedure IV:11** on  $\langle q_{11}, q_{12}, q_{13}, q_{14}, q_{15} \rangle$  and let  $\langle q_3, q_4 \rangle$  receive.
  - (1) Execute **procedure IV:33** on  $\langle f, e_1, \frac{1-e_1}{2} \rangle$  and let  $\langle q_{11}, q_{12} \rangle$  receive.
  - (2) Execute **procedure IV:10** on  $\langle q_{16}, q_{17}, q_{18}, q_{19} \rangle$  and let  $\langle q_{13}, q_{14} \rangle$  receive.
    - (a) Execute **procedure IV:15** on  $\langle 1, 1 \rangle$  and let  $\langle q_{16}, q_{17} \rangle$  receive.
    - (b) Execute **procedure IV:12** on  $\langle -2, q_{20}, q_{21} \rangle$  and let  $\langle q_{18}, q_{19} \rangle$  receive.
      - (i) Execute **procedure IV:11** on  $\langle q_{22}, q_{23}, q_{24}, q_{25}, q_{26} \rangle$  and let  $\langle q_{20}, q_{21} \rangle$  receive.
        - (1) Execute **procedure IV:18** on  $\langle 1, 1 + \epsilon, 1 + X \rangle$  and let  $\langle q_{22}, q_{23} \rangle$  receive.
        - (2) Execute **procedure IV:10** on  $\langle q_{27}, q_{28}, q_{29}, q_{30} \rangle$  and let  $\langle q_{24}, q_{25} \rangle$  receive.
          - (a) Execute **procedure IV:15** on  $\langle 1, 1 \rangle$  and let  $\langle q_{27}, q_{28} \rangle$  receive.
          - (b) Execute **procedure IV:16** on  $\langle 1, X, 1 \rangle$  and let  $\langle q_{29}, q_{30} \rangle$  receive.

- (3) Let  $q_{26}(x, n)$  be the following procedure:
- (a) Show that  $(1 + \epsilon)^2$ 
    - (i)  $\leq (1 + \operatorname{re}(x))^2$
    - (ii)  $\leq \operatorname{re}(x + 1)^2$
    - (iii)  $\leq \|x + 1\|^2$
    - (iv)  $\leq (X + 1)^2$ .
- (3) Let  $q_{15}(x, n)$  be the following procedure:
- (a) **Hence show that**  $\|1 + (-2)(x + 1)^{-1}\|^2 = \|\frac{x-1}{x+1}\|^2 \leq e_1^2$  **using procedure**  $q_9$ .
- vi) Execute **procedure IV:11** on  $\langle q_{31}, q_{32}, q_{33}, q_{34}, q_{35} \rangle$  and let  $\langle q_5, q_6 \rangle$  receive.
- (1) Execute **procedure IV:33** on  $\langle -f, e_1, \frac{1-\epsilon}{2} \rangle$  and let  $\langle q_{31}, q_{32} \rangle$  receive.
  - (2) Execute **procedure IV:12** on  $\langle -1, q_{13}, q_{14} \rangle$  and let  $\langle q_{33}, q_{34} \rangle$  receive.
  - (3) Let  $q_{35}(x, n)$  be the following procedure:
    - (a) **Hence show that**  $\|-(1 + (-2)(x + 1)^{-1})\|^2 = \|\frac{x-1}{x+1}\|^2 \leq e_1^2$  **using procedure**  $q_9$ .

## Procedure IV:41(thu2208191859)

### Objective

Choose a complex number  $f$  and two rational numbers  $X, \epsilon$  such that  $0 < \epsilon < 1$  and  $X \geq 0$ . The objective of the following instructions is to construct rational numbers  $c, d, a, b, e$ , a procedure  $p(x, n)$  to show that  $\|fx_n^{f-1}\|^2 \leq c^2$  when a complex number  $x$  and a positive integer  $n$  such that  $\operatorname{re}(x) \geq \epsilon$ ,  $\|x\|^2 \leq X^2$ , and  $n > d$  are chosen, and a procedure  $q(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} x_n^f \equiv fx_n^{f-1}$  (err  $\frac{a}{n} + b\{\delta\}$ ) when in addition a complex number  $\delta$  such that  $0 < \|\delta\|^2 \leq e^2$  is chosen.

### Implementation

- 1) Execute **procedure IV:30** on  $\langle X, \epsilon \rangle$  and let  $\langle a_1, p_1 \rangle$  receive.
- 2) Execute **procedure III:54** on  $\langle \{f\} + 1, a_1 \rangle$  and let  $\langle a_2, b_2, p_2 \rangle$  receive.

- 3) Execute **procedure III:55** on  $\langle \{f\} + 1, a_1 \rangle$  and let  $\langle a_3, p_3 \rangle$  receive.
- 4) Execute **procedure III:53** on  $\langle b_2, 1 \rangle$  and let  $\langle a_4, b_4, p_4 \rangle$  receive.
- 5) Execute **procedure IV:40** on  $\langle f, X, \epsilon \rangle$  and let  $\langle p_5, p_6 \rangle$  receive.
- 6) Let  $t$  be **subprocedure IV:42:0**.
- 7) **Execute procedure IV:14** on  $\langle t, p_5, p_6 \rangle$  and let  $\langle c, d, a, b, e, p, q \rangle$  receive.

### Subprocedure IV:42:0

**Objective** Choose a complex number  $f$  and two rational numbers  $X, \epsilon$  such that  $0 < \epsilon < 1$  and  $X \geq 0$ . Let  $g(f, x, n)$  be a shorthand for  $[2f(1 + \frac{x-1}{x+1})_n^f \cdot (1 - \frac{x-1}{x+1})_n^{-f-1}(x+1)^{-2}] + [2f(1 + \frac{x-1}{x+1})_n^{f-1} \cdot (1 - \frac{x-1}{x+1})_n^{-f}(x+1)^{-2}]$ . The objective of the following instructions is to construct a rational number  $h$ , and a procedure  $t(x, n)$  to show that  $g(f, x, n) \equiv fx_n^{f-1}$  (err  $\frac{h}{n}$ ) when a complex number  $x$  and a positive integer  $n$  such that  $\operatorname{re}(x) \geq \epsilon$ ,  $\|x\|^2 \leq X^2$ , and  $n > d$  are chosen.

### Implementation

- 1. Let  $h = \{f\}a_2a_3a_4((\frac{2}{1+\epsilon})^2 + 1)$ .
- 2. Let  $t(x, n)$  be the following procedure:
  - (a) Show that  $\|\frac{1}{(x+1)^2}\|^2$ 
    - i.  $= \frac{1}{\|1+x\|^4}$
    - ii.  $= \frac{1}{(\operatorname{re}(1+x)^2 + \operatorname{im}(x)^2)^2}$
    - iii.  $\leq (\frac{1}{1+\epsilon})^4$ .
  - (b) Using procedures  $p_1, p_2, p_3, p_4$ , show that
 
$$[2f(1 + \frac{x-1}{x+1})_n^f \cdot (1 - \frac{x-1}{x+1})_n^{-f-1}(x+1)^{-2}] + [2f(1 + \frac{x-1}{x+1})_n^{f-1} \cdot (1 - \frac{x-1}{x+1})_n^{-f}(x+1)^{-2}]$$
    - i.  $\equiv [2f(1 + \frac{x-1}{x+1})_n^{f-1}(1 + \frac{x-1}{x+1})_n^1 \cdot (1 - \frac{x-1}{x+1})_n^{-f-1}(x+1)^{-2}] + [2f(1 + \frac{x-1}{x+1})_n^{f-1} \cdot (1 - \frac{x-1}{x+1})_n^{-f}(x+1)^{-2}]$
    - A. (err  $2\{f\}a_2b_2^n a_3(\frac{1}{1+\epsilon})^2$ )
    - B. (err  $\frac{2\{f\}a_2a_3a_4}{n}(\frac{1}{1+\epsilon})^2$ )

$$\text{ii.} \equiv [2f(1 + \frac{x-1}{x+1})_n^{f-1}(1 + \frac{x-1}{x+1})_n^1 \cdot (1 - \frac{x-1}{x+1})_n^{-f-1}(x+1)^{-2}] + [2f(1 + \frac{x-1}{x+1})_n^{f-1} \cdot (1 - \frac{x-1}{x+1})_n^{-f-1}(1 - \frac{x-1}{x+1})_n^1(x+1)^{-2}]$$

$$\text{A. } (\text{err } 2\{f\}a_3a_2b_2^n(\frac{1}{1+\epsilon})^2)$$

$$\text{B. } (\text{err } \frac{2\{f\}a_3a_2a_4}{n}(\frac{1}{1+\epsilon})^2)$$

$$\text{iii.} = 2f(1 + \frac{x-1}{x+1})_n^{f-1}(1 - \frac{x-1}{x+1})_n^{-f-1}[(1 + \frac{x-1}{x+1})^1 \cdot (x+1)^{-2} + (1 - \frac{x-1}{x+1})^1(x+1)^{-2}]$$

$$\text{iv.} = 4f(1 + \frac{x-1}{x+1})_n^{f-1}(1 - \frac{x-1}{x+1})_n^{-f-1}[(x+1)^{-2}]$$

$$\text{v.} = f(1 + \frac{x-1}{x+1})_n^{f-1}(1 - \frac{x-1}{x+1})_n^{-f-1}(1 - \frac{x-1}{x+1})_n^2$$

$$\text{vi.} \equiv f(1 + \frac{x-1}{x+1})_n^{f-1}(1 - \frac{x-1}{x+1})_n^{-(f-1)}$$

$$\text{A. } (\text{err } \{f\}a_3a_2b_2^n)$$


$$\text{B. } (\text{err } \frac{\{f\}a_3a_2a_4}{n})$$

$$\text{vii.} = fx_n^{f-1}$$

$$\text{(c) Hence show that } [2f(1 + \frac{x-1}{x+1})_n^f \cdot (1 - \frac{x-1}{x+1})_n^{-f-1}(x+1)^{-2}] + [2f(1 + \frac{x-1}{x+1})_n^{f-1} \cdot (1 - \frac{x-1}{x+1})_n^{-f}(x+1)^{-2}] \equiv fx_n^{f-1} (\text{err } \frac{h}{n}).$$

3. Yield the tuple  $\langle h, t \rangle$ .

Figure IV:0



A plot of the lists of complex numbers  $(\frac{[0:40]}{10})_{15}^{\frac{1}{2}}$  and  $\frac{1}{2}(\frac{[0:40]}{10})_{15}^{\frac{1}{2}-1}$ . Note that the steepness of  $(\frac{[0:40]}{10})_{15}^{\frac{1}{2}}$  is approximately given by the y-coordinates of  $\frac{1}{2}(\frac{[0:40]}{10})_{15}^{\frac{1}{2}-1}$ . That is, where the graph of  $(\frac{[0:40]}{10})_{15}^{\frac{1}{2}}$  is rapidly increasing, the graph of  $\frac{1}{2}(\frac{[0:40]}{10})_{15}^{\frac{1}{2}-1}$  has a relatively large positive value, and where the graph of  $(\frac{[0:40]}{10})_{15}^{\frac{1}{2}}$  flattens out, the graph of  $\frac{1}{2}(\frac{[0:40]}{10})_{15}^{\frac{1}{2}-1}$  has a relatively small positive value.

# Chapter 14

## Integral Arithmetic

### Declaration IV:4

The notation  $\int_r^R f(r, \delta_r)$ , where:

1.  $f(r, \delta_r)$  is a procedure to construct a complex number when complex numbers  $r, \delta_r$  such that  $P(r, \delta_r)$  are chosen
2.  $R$  is a non-empty list of complex numbers such that  $P(R_t, R_{t+1} - R_t)$  for  $t \in [0 : |R| - 1]$

will be used as a shorthand for  $\sum_t^{[0:|R|-1]} f(R_t, R_{t+1} - R_t)$ .

### Procedure IV:42(3.86)

#### Objective

Choose the following:

1. A procedure  $f(r, \delta)$  to construct a complex number when complex numbers  $r, \delta_r$  such that  $P(r, \delta_r)$  are chosen.
2. A procedure  $g(r, \delta)$  to construct a complex number when complex numbers  $r, \delta_r$  such that  $Q(r, \delta_r)$  are chosen.
3. A non-empty list of complex numbers  $R$  such that  $P(R_t, R_{t+1} - R_t)$  and  $Q(R_t, R_{t+1} - R_t)$  for  $t \in [0 : |R| - 1]$ .

The objective of the following instructions is to show that  $\int_r^R (f(r, \delta_r) + g(r, \delta_r)) = \int_r^R f(r, \delta_r) + \int_r^R g(r, \delta_r)$ .

### Implementation

1. Show that  $\int_r^R (f(r, \delta_r) + g(r, \delta_r))$ 
  - (a)  $= \sum_t^{[0:|R|-1]} (f(R_t, R_{t+1} - R_t) + g(R_t, R_{t+1} - R_t))$
  - (b)  $= \sum_t^{[0:|R|-1]} f(R_t, R_{t+1} - R_t) + \sum_t^{[0:|R|-1]} g(R_t, R_{t+1} - R_t)$
  - (c)  $= \int_r^R f(r, \delta_r) + \int_r^R g(r, \delta_r)$

### Procedure IV:43(3.87)

#### Objective

Choose the following:

1. A complex number  $a$ .
2. A procedure  $f(r, \delta)$  to construct a complex number when complex numbers  $r, \delta_r$  such that  $P(r, \delta_r)$  are chosen.
3. A non-empty list of complex numbers  $R$  such that  $P(R_t, R_{t+1} - R_t)$  for  $t \in [0 : |R| - 1]$ .

The objective of the following instructions is to show that  $\int_r^R a f(r, \delta_r) = a \int_r^R f(r, \delta_r)$ .

### Implementation

1. Show that  $\int_r^R a f(r, \delta_r)$ 
  - (a)  $= \sum_t^{[0:|R|-1]} a f(R_t, R_{t+1} - R_t)$
  - (b)  $= a \sum_t^{[0:|R|-1]} f(R_t, R_{t+1} - R_t)$

$$(c) = a \int_r^R f(r, \delta_r)$$

#### Procedure IV:44(3.88)

##### Objective

Choose the following:

1. A procedure  $f(r)$  to construct a complex number when a complex number  $r$  such that  $P(r)$  is chosen.
2. A non-empty list of complex numbers  $R$  such that  $P(R_t)$  for  $t \in [0 : |R| - 1]$ .
3. A non-empty list of complex numbers  $S$  such that  $P(S_t)$  for  $t \in [0 : |R| - 1]$  and  $R_{|R|-1} = S_0$ .

The objective of the following instructions is to show that  $\int_r^{R \cap S} f(r) \delta_r = \int_r^R f(r) \delta_r + \int_r^S f(r) \delta_r$ .

##### Implementation

1. Let  $T = R \cap S$ .
2. Show that  $\int_r^T f(r) \delta_r$ 
  - (a)  $= \sum_t^{[0:|T|-1]} f(T_t)(T_{t+1} - T_t)$
  - (b)  $= \sum_t^{[0:|R|-1]} f(T_t)(T_{t+1} - T_t) + \sum_t^{[|R|-1:|R|]} f(T_t)(T_{t+1} - T_t) + \sum_t^{[|R|:|T|-1]} f(T_t)(T_{t+1} - T_t)$
  - (c)  $= \sum_t^{[0:|R|-1]} f(R_t)(R_{t+1} - R_t) + f(T_{|R|-1})(T_{|R|-1} - R_{|R|-1}) + \sum_t^{[|R|:|T|-1]} f(S_{t-|R|})(S_{t+1-|R|} - S_{t-|R|})$
  - (d)  $= \sum_t^{[0:|R|-1]} f(R_t)(R_{t+1} - R_t) + f(T_{|R|-1})(S_0 - R_{|R|-1}) + \sum_t^{[0:|S|-1]} f(S_t)(S_{t+1} - S_t)$
  - (e)  $= \int_r^R f(r) \delta_r + \int_r^S f(r) \delta_r$ .

#### Procedure IV:45(3.34)

##### Objective

Choose the following:

1. A procedure  $f(r)$  to construct a complex number when a complex number  $r$  such that  $P(r)$  is chosen.

2. A non-empty list of complex numbers  $R$  such that  $P(R_t)$  for  $t \in [0 : |R| - 1]$ .

The objective of the following instructions is to show that  $\int_r^R \delta_r \Delta_{z=r}^{+\delta_r} f(z) = f(R_{|R|-1}) - f(R_0)$ .

##### Implementation

1. Show that  $\int_r^R \delta_r \Delta_{z=r}^{+\delta_r} f(z)$ 
  - (a)  $= \int_r^R \delta_r \left( \frac{f(r+\delta_r) - f(r)}{\delta_r} \right)$
  - (b)  $= \int_r^R (f(r + \delta_r) - f(r))$
  - (c)  $= \sum_k^{[0:|R|-1]} (f(R_{k+1}) - f(R_k))$
  - (d)  $= f(R_{|R|-1}) - f(R_0)$ .

##### Declaration IV:5

The notation  $\Delta X$ , where  $X$  is a list, will be used as a shorthand for  $\langle X_1 - X_0, X_2 - X_1, \dots, X_{|X|-1} - X_{|X|-2} \rangle$ .

#### Procedure IV:46(fri3008190328)

##### Objective

Choose the following:

1. A non-negative rational number  $A$ .
2. A procedure  $q_1(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} f_n(y) \equiv f'_n(x)$  (err  $\frac{a_1}{n} + b_1 \{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > c_1$ , and  $0 < \|\delta\|^2 < d_1^2$  are chosen.

The objective of the following instructions is to construct the following:

1. Non-negative rational numbers  $a, b, c, d$ .
2. A procedure  $p(R, n)$  to show that  $\int_r^R f'_n(r) \delta_r \equiv f_n(R_{|R|-1}) - f_n(R_0)$  (err  $\frac{a}{n} + b \max(\{\Delta R\})$ ) when an integer  $n$  and a non-empty list of complex numbers  $R$  such that  $P(R_t)$  and  $0 < \|R_{t+1} - R_t\|^2 < d^2$  for  $t \in [0 : |R| - 1]$ ,  $\int_r^R \{\delta_r\} \leq A$ , and  $n > c$  are chosen.

## Implementation

1. Let  $a = a_1 A$ .
2. Let  $b = b_1 A$ .
3. Let  $c = c_1$ .
4. Let  $d = d_1$ .
5. Let  $p(R, n)$  be the following procedure:

- (a) Using procedure  $q_1$ , show that  $\int_r^R f'_n(r) \delta_r$ 
  - i.  $= \sum_k^{[0:|R|-1]} f'_n(R_k)(R_{k+1} - R_k)$
  - ii.  $\equiv \sum_k^{[0:|R|-1]} \Delta_{y=R_k}^{R_{k+1}-R_k} f_n(y)(R_{k+1} - R_k)$ 
    - A. (err  $\sum_k^{[0:|R|-1]} (\frac{a_1}{n} + b_1 \{R_{k+1} - R_k\}) \{R_{k+1} - R_k\}$ )
    - B. (err  $(\frac{a_1}{n} + b_1 \max(\{\Delta R\})) \sum_k^{[0:|R|-1]} \{R_{k+1} - R_k\}$ )
    - C. (err  $(\frac{a_1}{n} + b_1 \max(\{\Delta R\})) \int_r^R \{\delta_r\}$ )
    - D. (err  $(\frac{a_1}{n} + b_1 \max(\{\Delta R\})) A$ )
    - E. (err  $\frac{a}{n} + b \max(\{\Delta R\})$ )
  - iii.  $= \sum_k^{[0:|R|-1]} \frac{f_n(R_k + (R_{k+1} - R_k)) - f_n(R_k)}{R_{k+1} - R_k} (R_{k+1} - R_k)$
  - iv.  $= \sum_k^{[0:|R|-1]} (f_n(R_{k+1}) - f_n(R_k))$
  - v.  $= f_n(R_{|R|-1}) - f_n(R_0)$ .

(b) **Therefore show that**  $\int_r^R f'_n(r) \delta_r \equiv f_n(R_{|R|-1}) - f_n(R_0)$  (err  $\frac{a}{n} + b \max(\{\Delta R\})$ ).

6. **Yield the tuple**  $\langle a, b, c, d, p \rangle$ .

## Procedure IV:47(fri3008190457)

### Objective

Choose the following:

1. A non-negative rational number  $A$ .
2. A procedure  $q_1(x, n, \delta)$  to show that  $\Delta_{y=x}^{+\delta} g_n(y) \equiv g'_n(x)$  (err  $\frac{a_1}{n} + b_1 \{\delta\}$ ) when two complex numbers  $x, \delta$  and a positive integer  $n$  such that  $P(x)$ ,  $n > c_1$ , and  $0 < \|\delta\|^2 < d_1^2$  are chosen.

3. A procedure  $q_2(x, n)$  to show that  $f_n(x) \equiv 0$  (err  $a_2$ ) when a complex number  $x$  and a positive integer  $n$  such that  $Q(x)$  and  $n > b_2$  are chosen.
4. A procedure  $q_3(x, n)$  to show that  $Q(g_n(x))$  when a complex number  $x$  and a positive integer  $n$  such that  $P(x)$  and  $n > c_1$  are chosen

The objective of the following instructions is to construct the following:

1. Non-negative rational numbers  $a, b, c, d$ .
2. A procedure  $p(R, n)$  to show that  $\int_r^{g(R)} f_n(r) \delta_r \equiv \int_r^R f_n(g_n(r)) g'_n(r) \delta_r$  (err  $\frac{a}{n} + b \max(\{\Delta R\})$ ) when an integer  $n$  and a non-empty list of complex numbers  $R$  such that  $P(R_t)$  and  $0 < \|R_{t+1} - R_t\|^2 < d^2$  for  $t \in [0 : |R| - 1]$ ,  $\int_r^R \{\delta_r\} \leq A$ , and  $n > c$  are chosen.

## Implementation

1. Let  $a = a_1 a_2 A$ .
2. Let  $b = b_1 a_2 A$ .
3. Let  $c = \max(c_1, b_2)$ .
4. Let  $d = d_1$ .
5. Let  $p(R, n)$  be the following procedure:
  - (a) Using procedures  $q_1, q_2, q_3$ , show that  $\int_r^{g_n(R)} f_n(r) \delta_r$ 
    - i.  $= \sum_k^{[0:|R|-1]} f_n(g_n(R_k))(g_n(R_{k+1}) - g_n(R_k))$
    - ii.  $= \sum_k^{[0:|R|-1]} f_n(g_n(R_k)) \Delta_{y=R_k}^{R_{k+1}-R_k} g_n(y)(R_{k+1} - R_k)$
    - iii.  $\equiv \sum_k^{[0:|R|-1]} f_n(g_n(R_k)) g'_n(R_k)(R_{k+1} - R_k)$ 
      - A. (err  $\sum_k^{[0:|R|-1]} a_2 (\frac{a_1}{n} + b_1 \{R_{k+1} - R_k\}) \{R_{k+1} - R_k\}$ )
      - B. (err  $a_2 (\frac{a_1}{n} + b_1 \max(\{\Delta R\})) \sum_k^{[0:|R|-1]} \{R_{k+1} - R_k\}$ )
      - C. (err  $a_2 (\frac{a_1}{n} + b_1 \max(\{\Delta R\})) \int_r^R \{\delta_r\}$ )
      - D. (err  $a_2 (\frac{a_1}{n} + b_1 \max(\{\Delta R\})) A$ )
      - E. (err  $\frac{a}{n} + b \max(\{\Delta R\})$ )



$$\text{iv.} = \int_r^R f_n(g_n(r))g'_n(r)\delta_r.$$

$$\begin{aligned} \text{(b) Hence show that } \int_r^{g_n(R)} f_n(r)\delta_r &\equiv \\ \int_r^R f_n(g_n(r))g'_n(r)\delta_r &(\text{err } \frac{a}{n} + b \max(\{\Delta R\})). \end{aligned}$$

6. Yield the tuple  $\langle a, b, c, d, p \rangle$ .

## Procedure IV:48(fri3008190709)

### Objective

Choose three rational numbers  $A, X, \epsilon$  such that  $0 < \epsilon < 1$  and  $X \geq 0$ . The objective of the following instructions is to construct the following:

1. Non-negative rational numbers  $a, b, c, d$ .
2. A procedure  $p(R, n)$  to show that  $\int_r^R \frac{\delta_r}{r} \equiv \ln_n(R_{|R|-1}) (\text{err } \frac{a}{n} + b \max(\{\Delta R\}))$  when an integer  $n$  and a non-empty list of complex numbers  $R$  such that  $\text{re}(R_t) \geq \epsilon$ ,  $\|R_t\|^2 \leq X^2$  and  $0 < \|R_{t+1} - R_t\|^2 < d^2$  for  $t \in [0 : |R| - 1]$ ,  $R_0 = 1$ ,  $\int_r^R \{\delta_r\} \leq A$ , and  $n > c$  are chosen.

### Implementation

1. Execute **procedure IV:31** on  $\langle X, \epsilon \rangle$  and let  $\langle \dots, q \rangle$  receive.
2. Hence execute **procedure IV:46** on  $\langle A, q \rangle$  and let  $\langle a, b, c, d, t \rangle$  receive.
3. Let  $p(R, n)$  be the following procedure:
  - (a) Using procedure  $t$ , show that  $\int_r^R \frac{\delta_r}{r}$ 
    - i.  $\equiv \ln_n(R_{|R|-1}) - \ln_n(R_0) (\text{err } \frac{a}{n} + b \max(\{\Delta R\}))$
    - ii.  $= \ln_n(R_{|R|-1}) - \ln_n(1)$
    - iii.  $= \ln_n(R_{|R|-1})$ .
  - (b) **Hence show that**  $\int_r^R \frac{\delta_r}{r} \equiv \ln_n(R_{|R|-1}) (\text{err } \frac{a}{n} + b \max(\{\Delta R\}))$ .
4. Yield the tuple  $\langle a, b, c, d, p \rangle$ .

## Part V

# Matrix Arithmetic

# Chapter 15

## Matrix Arithmetic

### Declaration V:0

The phrase "matrix" will be used as a shorthand for a list of equally lengthed lists of polynomials. In particular, the phrase " $m \times n$  matrix" will be used as a shorthand for a length- $m$  list of length- $n$  lists of polynomials.

### Declaration V:1

The notation  $A_{I,J}$ , where  $A$  is a matrix and  $I, J$  are lists of indices, will be used as a shorthand for  $\langle (A_j)_J \text{ for } j \in I \rangle$ .

### Declaration V:2

The phrase " $A = B$ ", where  $A, B$  are  $m \times n$  matrices, will be used as a shorthand for " $A_{i,j} = B_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ ".

### Procedure V:0(4.73)

#### Objective

Choose an  $m \times n$  matrix  $A$ . The objective of the following instructions is to show that  $A = A$ .

#### Implementation

1. Verify that  $A_{i,j} = A_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .
2. Hence verify that  $A = A$ .

### Procedure V:1(4.74)

#### Objective

Choose two  $m \times n$  matrices  $A, B$  such that  $A = B$ . The objective of the following instructions is to show that  $B = A$ .

#### Implementation

1. Verify that  $A_{i,j} = B_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .
2. Hence verify that  $B_{i,j} = A_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .
3. Hence verify that  $B = A$ .

### Procedure V:2(4.75)

#### Objective

Choose three  $m \times n$  matrices  $A, B, C$  such that  $A = B$  and  $B = C$ . The objective of the following instructions is to show that  $A = C$ .

#### Implementation

1. Verify that  $A_{i,j} = B_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .
2. Verify that  $B_{i,j} = C_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .

3. Hence verify that  $A_{i,j} = C_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .

4. **Hence verify that  $A = C$ .**

#### Declaration V:3

The notation  $A + B$ , where  $A, B$  are  $m \times n$  matrices, will be used as a shorthand for the list  $\langle\langle A_{i,j} + B_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$ .

#### Procedure V:3(4.76)

##### Objective

Choose four  $m \times n$  matrices  $A, B, C, D$  such that  $A = C$  and  $B = D$ . The objective of the following instructions is to show that  $A + B = C + D$ .

##### Implementation

1. Verify that  $A_{i,j} = C_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .
2. Verify that  $B_{i,j} = D_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .
3. Hence verify that  $A + B$ 
  - (a)  $= \langle\langle A_{i,j} + B_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (b)  $= \langle\langle C_{i,j} + D_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (c)  $= C + D$ .

#### Procedure V:4(4.77)

##### Objective

Choose three  $m \times n$  matrices  $A, B, C$ . The objective of the following instructions is to show that  $(A + B) + C = A + (B + C)$ .

##### Implementation

1. Verify that  $(A + B) + C$ 
  - (a)  $= \langle\langle (A + B)_{i,j} + C_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (b)  $= \langle\langle (A_{i,j} + B_{i,j}) + C_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$

(c)  $= \langle\langle A_{i,j} + (B_{i,j} + C_{i,j}) \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$

(d)  $= \langle\langle A_{i,j} + (B + C)_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$

(e)  $= A + (B + C)$ .

#### Procedure V:5(4.78)

##### Objective

Choose two  $m \times n$  matrices  $A, B$ . The objective of the following instructions is to show that  $A + B = B + A$ .

##### Implementation

1.  $A + B$ 
  - (a)  $= \langle\langle A_{i,j} + B_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (b)  $= \langle\langle B_{i,j} + A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (c)  $= B + A$ .

#### Declaration V:4

The notation  $0_{m \times n}$  will contextually be used as a shorthand for the list  $\langle\langle 0 \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$  where the natural numbers  $m, n$  are determined by the context.

#### Procedure V:6(4.79)

##### Objective

Choose an  $m \times n$  matrix  $A$ . The objective of the following instructions is to show that  $0 + A = A$ .

##### Implementation

1. Verify that  $0 + A$ 
  - (a)  $= 0_{m \times n} + A$
  - (b)  $= \langle\langle 0_{i,j} + A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (c)  $= \langle\langle 0 + A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (d)  $= \langle\langle A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (e)  $= A$ .

**Declaration V:5**

The notation  $-A$ , where  $A$  is an  $m \times n$  matrix, will be used as a shorthand for the list  $\langle \langle -A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$ .

**Procedure V:7(4.80)****Objective**

Choose two  $m \times n$  matrices  $A, B$  such that  $A = B$ . The objective of the following instructions is to show that  $-A = -B$ .

**Implementation**

1. Verify that  $A_{i,j} = B_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .
2. Hence verify that  $-A$ 
  - (a)  $= \langle \langle -A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (b)  $= \langle \langle -B_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (c)  $= -B$ .

**Procedure V:8(4.81)****Objective**

Choose a  $m \times n$  matrix  $A$ . The objective of the following instructions is to show that  $-A + A = 0$ .

**Implementation**

1. Verify that  $-A + A$ 
  - (a)  $\langle \langle (-A)_{i,j} + A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (b)  $\langle \langle -(A_{i,j}) + A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (c)  $\langle \langle 0 \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (d)  $= 0$ .

**Declaration V:6**

The notation  $AB$ , where  $A$  is an  $m \times n$  matrix and  $B$  is an  $n \times k$  matrix, will be used as a shorthand for the list  $\langle \langle \sum_r^{[0:n]} A_{i,r} B_{r,j} \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle$ .

**Procedure V:9(4.82)****Objective**

Choose two  $m \times n$  matrices  $A, C$  and two  $n \times k$  matrices  $B, D$  such that  $A = C$  and  $B = D$ . The objective of the following instructions is to show that  $AB = CD$ .

**Implementation**

1. Verify that  $A_{i,j} = C_{i,j}$  for  $j \in [0 : n]$ , for  $i \in [0 : m]$ .
2. Verify that  $B_{i,j} = D_{i,j}$  for  $j \in [0 : k]$ , for  $i \in [0 : n]$ .
3. Hence verify that  $AB$ 
  - (a)  $= \langle \langle \sum_r^{[0:n]} A_{i,r} B_{r,j} \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle$
  - (b)  $= \langle \langle \sum_r^{[0:n]} C_{i,r} D_{r,j} \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle$
  - (c)  $= CD$ .

**Procedure V:10(4.02)****Objective**

Choose an  $m \times n$  matrix,  $A$ , an  $n \times p$  matrix,  $B$ , and a  $p \times q$  matrix,  $C$ . The objective of the following instructions is to show that  $(AB)C = A(BC)$ .

**Implementation**

1. Verify that  $(AB)C$ 
  - (a)  $= \langle \langle \sum_r^{[0:p]} (AB)_{i,r} C_{r,j} \text{ for } j \in [0 : q] \rangle \text{ for } i \in [0 : m] \rangle$
  - (b)  $= \langle \langle \sum_r^{[0:p]} (\sum_l^{[0:n]} A_{i,l} B_{l,r}) C_{r,j} \text{ for } j \in [0 : q] \rangle \text{ for } i \in [0 : m] \rangle$

- (c) =  $\langle \langle \sum_r^{[0:p]} \sum_l^{[0:n]} A_{i,l} B_{l,r} C_{r,j} \text{ for } j \in [0 : q] \rangle \text{ for } i \in [0 : m] \rangle$
- (d) =  $\langle \langle \sum_l^{[0:n]} \sum_r^{[0:p]} A_{i,l} B_{l,r} C_{r,j} \text{ for } j \in [0 : q] \rangle \text{ for } i \in [0 : m] \rangle$
- (e) =  $\langle \langle \sum_l^{[0:n]} A_{i,l} \sum_r^{[0:p]} B_{l,r} C_{r,j} \text{ for } j \in [0 : q] \rangle \text{ for } i \in [0 : m] \rangle$
- (f) =  $\langle \langle \sum_l^{[0:n]} A_{i,l} (BC)_{l,j} \text{ for } j \in [0 : q] \rangle \text{ for } i \in [0 : m] \rangle$
- (g) =  $A(BC)$ .

#### Declaration V:7

The notation  $\textcolor{red}{a}_{m \times m}$ , where  $a \neq 0$  is a polynomial, will contextually be used as a shorthand for the list  $\langle \langle a[i = j] \text{ for } j \in [0 : m] \rangle \text{ for } i \in [0 : m] \rangle$ .

#### Procedure V:11(4.84)

##### Objective

Choose an  $m \times n$  matrix,  $A$ . The objective of the following instructions is to show that  $1A = A$ .

##### Implementation

1. Verify that  $1A$ 
  - (a) =  $1_m A$
  - (b) =  $\langle \langle \sum_r^{[0:m]} 1_{i,r} A_{r,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (c) =  $\langle \langle \sum_r^{[0:m]} [i = r] A_{r,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (d) =  $\langle \langle A_{i,j} \text{ for } j \in [0 : n] \rangle \text{ for } i \in [0 : m] \rangle$
  - (e) =  $A$ .

#### Procedure V:12(4.85)

##### Objective

Choose an  $m \times n$  matrix  $A$ , and two  $n \times k$  matrices  $B, C$ . The objective of the following instructions is to show that  $A(B + C) = AB + AC$ .

##### Implementation

1.  $A(B + C)$ 
  - (a) =  $\langle \langle \sum_r^{[0:n]} A_{i,r} (B + C)_{r,j} \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle$
  - (b) =  $\langle \langle \sum_r^{[0:n]} A_{i,r} (B_{r,j} + C_{r,j}) \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle$
  - (c) =  $\langle \langle \sum_r^{[0:n]} (A_{i,r} B_{r,j} + A_{i,r} C_{r,j}) \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle$
  - (d) =  $\langle \langle \sum_r^{[0:n]} A_{i,r} B_{r,j} + \sum_r^{[0:n]} A_{i,r} C_{r,j} \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle$
  - (e) =  $\langle \langle \sum_r^{[0:n]} A_{i,r} B_{r,j} \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle + \langle \langle \sum_r^{[0:n]} \sum_r^{[0:n]} A_{i,r} C_{r,j} \text{ for } j \in [0 : k] \rangle \text{ for } i \in [0 : m] \rangle$
  - (f) =  $AB + AC$ .

#### Declaration V:8

The phrase "row  $i$  of  $A$ " and the notation  $\textcolor{red}{A}_{i,*}$ , where  $A$  is an  $m \times n$  matrix and  $0 \leq i < m$ , will be used as a shorthand for  $A_{i,[0:n]}$ .

#### Declaration V:9

The phrase "column  $i$  of  $A$ " and the notation  $\textcolor{red}{A}_{*,i}$ , where  $A$  is an  $m \times n$  matrix and  $0 \leq i < n$ , will be used as a shorthand for  $A_{[0:m],i}$ .

#### Procedure V:13(4.00)

##### Objective

Choose an  $m \times 2$  matrix,  $A$ . Let  $\deg(0) = \infty$ . Let  $k = \min(\deg(A_{0,0}), \deg(A_{0,1}))$  and  $q = \deg(A_{0,0})$ . The objective of the following instructions is to make  $A_{0,1} = 0$ ,  $\deg(A_{0,0}) \leq k$ , and either leave  $A_{*,0}$  unchanged or make  $\deg(A_{0,0}) < q$  by a sequence of operations whereby, in each step a polynomial times either of the columns is added to the other.

## Implementation

1. Let  $A$  be our working matrix.
2. While  $A_{0,1} \neq 0$ , do the following:
  - (a) If  $\deg(A_{0,0}) \leq \deg(A_{0,1})$ , then:
    - i. Subtract  $\frac{(A_{0,1})^{\deg(A_{0,1})}}{(A_{0,0})^{\deg(A_{0,0})}} \lambda^{\deg(A_{0,1}) - \deg(A_{0,0})}$  times  $A_{0,0}$  from  $A_{0,1}$ .
    - ii. Now verify that either  $A_{0,1}$ 's degree has decreased or  $A_{0,1} = 0$ .
  - (b) Otherwise, do the following:
    - i. Let  $p = \frac{(A_{0,0})^{\deg(A_{0,0})}}{(A_{0,1})^{\deg(A_{0,1})}} \lambda^{\deg(A_{0,0}) - \deg(A_{0,1})}$ .
    - ii. If  $A_{0,0} = pA_{0,1}$ , then do the following:
      - A. Add  $1 - p$  times  $A_{0,1}$  to  $A_{0,0}$ .
      - B. Verify that now  $A_{0,0} = A_{0,1}$ .
    - iii. Otherwise, do the following:
      - A. Verify that  $A_{0,0} \neq pA_{0,1}$ .
      - B. Add  $-p$  times  $A_{0,1}$  to  $A_{0,0}$ .
    - iv. Therefore verify that  $A_{0,0} \neq 0$ .
    - v. Also verify that  $A_{0,0}$ 's degree has decreased.
3. **Verify that**  $A_{0,1} = 0$ .
4. Verify that the changes to  $A_{0,0}$ , if any, have decreased its degree.
5. If both operations are well-defined, then do the following:
  - (a) Verify that all changes to  $A_{0,1}$  but the last have decreased its degree.
  - (b) Verify that  $\deg(A_{0,0}) \leq$  the degree of the penultimate value of  $A_{0,1}$ .
6. **Therefore verify that**  $\deg(A_{0,0}) \leq k$ .
7. If  $A_{*,0}$  was changed, then do the following:
  - (a) Verify that  $A_{0,0}$  was also changed.
  - (b) **Therefore verify that**  $\deg(A_{0,0}) < q$ .
8. **Yield the tuple**  $\langle A \rangle$ .

## Declaration V:10

The phrase "**matrix diagonal**" will be used as a shorthand for matrix positions such that the row index equals the column index.

## Declaration V:11

The phrase "**diagonal matrix**" will be used to refer to matrices with 0s in all off-diagonal positions.

## Procedure V:14(4.01)

### Objective

Choose a  $m \times n$  matrix,  $A$ . The objective of the following instructions is to transform  $A$  into an  $m \times n$  diagonal matrix by a sequence of operations whereby either a polynomial times any of the columns is added to a different column, or a polynomial times any of the rows is added to a different row.

### Implementation

1. If  $m = 0$  or  $n = 0$ , then do the following:
  - (a) **Verify that  $A$  is an  $m \times n$  diagonal matrix.**
  - (b) **Yield the tuple**  $\langle A \rangle$ .
2. Otherwise do the following:
3. Verify that  $m > 0$  and  $n > 0$ .
4. Let  $A$  be our working matrix.
5. Now do the following:
  - (a) While  $A_{0,[1:n]} \neq 0$ , do the following:
    - i. Select the  $m \times 2$  matrix whose top-right entry coincides with the last non-zero entry of the first row
    - ii. Apply **procedure V:13** on this submatrix.
    - iii. Verify that the top-left and top-right entries of the submatrix are now non-zero and zero respectively.
    - iv. If  $A_{*,0}$  was modified by (5aii), then do the following:
      - A. Verify that  $\deg(A_{0,0})$  decreased.

- B. Go back to (5).
- (b) Now do the same operations as in (a), but this time with the operations themselves reflected across the matrix's diagonal.
6. Verify that  $A_{0,[1:n]} = 0$ .
7. Also verify that  $A_{[1:m],0} = 0$ .
8. Apply **procedure V:14** on the submatrix  $A_{[1:m],[1:n]}$ .
9. Verify that (8)'s execution leaves the first row and column unchanged.
10. Also verify that  $A_{[1:m],[1:n]}$  is now a  $(m-1) \times (n-1)$  diagonal matrix.
11. **Therefore verify that  $A$  is now an  $m \times n$  diagonal matrix.**
12. **Yield the tuple  $\langle A \rangle$ .**

#### Declaration V:12

The phrase "tilt matrix" will be used to refer to square matrices with only 1s on the diagonal, a single polynomial off the diagonal, and 0s everywhere else.

#### Procedure V:15(4.03)

##### Objective

Choose a procedure,  $A$ , and two non-negative integers  $m, n$ . The objective of the following instructions is, once  $A$  has been executed, to construct a list of  $m \times m$  tilts,  $M$ , and a list of  $n \times n$  tilts,  $N$  such that  $M_{|M|-1-i}$  equals  $1_m$  after applying the  $i^{th}$  row operation carried out by  $A$  also on it, and  $N_i$  equals  $1_n$  after applying the  $i^{th}$  row operation carried out by  $A$  also on it.

##### Implementation

1. Make an empty list,  $N$ .
2. Augment procedure  $A$  so that each time a polynomial  $x$  times a column  $i$  is added onto column  $j$ , an  $n \times n$  matrix that only has 1s on its diagonal, and such that the only non-zero entry off its diagonal is  $x$  at position  $(i, j)$ , is appended onto  $N$ .

3. Make an empty list,  $M$ .
4. Also augment procedure  $A$  so that each time a polynomial  $x$  times a row  $i$  is added onto row  $j$ , an  $n \times n$  matrix that only has 1s on its diagonal, and such that the only non-zero entry off its diagonal is  $x$  at position  $(j, i)$ , is prepended onto  $M$ .
5. Now run procedure  $A$ .
6. **Yield the tuple  $\langle M, N \rangle$ .**

#### Procedure V:16(4.04)

##### Objective

Choose a  $m \times n$  matrix,  $A$ . The objective of the following instructions is to show that  $1_m A = A = A 1_n$ .

##### Implementation

1. For  $0 \leq r < m$ , do the following:
  - (a) For  $0 \leq t < n$ , do the following:
    - i. Verify that  $(1_m A)_{r,t} = \sum_u^{[0:m]} (1_m)_{r,u} A_{u,t} = (1_m)_{r,r} A_{r,t} = 1 * A_{r,t} = A_{r,t}$ .
2. **Therefore verify that  $1_m A = A$ .**
3. For  $0 \leq r < m$ , do the following:
  - (a) For  $0 \leq t < n$ , do the following:
    - i. Verify that  $(A 1_n)_{r,t} = \sum_u^{[0:m]} A_{r,u} (1_n)_{u,t} = A_{r,t} (1_n)_{t,t} = A_{r,t} * 1 = A_{r,t}$ .
4. **Therefore verify that  $A 1_n = A$ .**

#### Declaration V:13

The notation  $A^{-1}$ , where  $A$  is a list of  $m \times m$  tilts, will be used to refer to the result yielded by executing the following instructions:

1. Let  $A^{-1}$  be  $\langle \rangle$ .
2. For  $i$  in  $[0 : |A|]$ , do the following:
  - (a) Let  $(j, k)$  be the position of the off diagonal entry of  $A_i$ .
  - (b) Let  $B$  equal  $A_i$  but with entry  $(j, k)$  negated.
  - (c) Now prepend  $B$  onto  $A^{-1}$ .



### 3. Yield $\langle A^{-1} \rangle$ .

#### Procedure V:17(4.05)

##### Objective

Choose a list of  $m \times m$  tilts,  $A$ . The objective of the following instructions is to show that  $A_* A^{-1}_* = 1_m$ .

##### Implementation

1. Verify that  $|A| = |A^{-1}|$ .
2. For  $i$  in  $[0 : |A|]$ , do the following:
  - (a) Let  $(j, k)$  be the position of the off diagonal entry of  $A_i$ .
  - (b) Let  $B = A^{-1}_{|A|-1-i}$ .
  - (c) For  $r$  in  $[0 : m]$  and  $r \neq j$ , do the following:
    - i. For  $t$  in  $[0 : m]$ , do the following:
      - A. Verify that  $(A_i B)_{r,t} = \sum_u^{[0:m]} (A_i)_{r,u} B_{u,t} = (A_i)_{r,r} B_{r,t} = 1 * B_{r,t} = [r = t]$ .
  - (d) For  $t$  in  $[0 : m]$  and  $t \neq k$ , do the following:
    - i. Verify that  $(A_i B)_{j,t} = \sum_u^{[0:m]} (A_i)_{j,u} B_{u,t} = (A_i)_{j,t} B_{t,t} = (A_i)_{j,t} * 1 = [j = t]$ .
  - (e) Verify that  $(A_i B)_{j,k} = \sum_u^{[0:m]} (A_i)_{j,u} B_{u,k} = (A_i)_{j,j} B_{j,k} + (A_i)_{j,k} B_{k,k} = 1 * B_{j,k} + (A_i)_{j,k} * 1 = B_{j,k} + (A_i)_{j,k} = 0$ .
  - (f) Therefore verify that  $A_i B = 1_m$ .
3. Therefore using **procedure V:10** and **procedure V:16**, verify that  $A_* A^{-1}_* = 1_m$ .
  - (a)  $= A_0 \cdots A_{|A|-2} A_{|A|-1} A^{-1}_0 A^{-1}_1 \cdots A^{-1}_{|A|-1}$
  - (b)  $= A_0 \cdots A_{|A|-3} A_{|A|-2} 1_m A^{-1}_1 A^{-1}_2 \cdots A^{-1}_{|A|-1}$
  - (c)  $= A_0 \cdots A_{|A|-3} A_{|A|-2} A^{-1}_1 A^{-1}_2 \cdots A^{-1}_{|A|-1}$
  - (d)  $\vdots$
  - (e)  $= A_0 1_m A^{-1}_{|A|-1}$
  - (f)  $= A_0 A^{-1}_{|A|-1}$
  - (g)  $= 1_m$ .

#### Procedure V:18(4.06)

##### Objective

Choose a list of  $m \times m$  tilts,  $A$ . The objective of the following instructions is to show that  $(A^{-1})^{-1} = A$  and  $A^{-1}_* A_* = 1_m$ .

##### Implementation

1. Verify that  $(A^{-1})^{-1} = A$ .
2. Therefore using **procedure V:17**, verify that  $A^{-1}_* A_* = A^{-1}_* (A^{-1})^{-1}_* = 1_m$ .

#### Procedure V:19(4.07)

##### Objective

Choose a  $2 \times 2$  diagonal matrix,  $A$ . The objective of the following instructions is to construct polynomials  $u, v$  and transform  $A$  into a  $2 \times 2$  diagonal matrix,  $A'$ , such that  $A'_{1,1} = u A'_{0,0}$  and  $A_{0,0} = v A'_{0,0}$  by a sequence of operations whereby either a polynomial times any of the columns is added to a different column, or a polynomial times any of the rows is added to a different row.

##### Implementation

1. Add row 1 to row 0.
2. Now verify that  $A_{0,1} = A_{1,1}$ .
3. Set  $A' = A$  and let  $A'$  be our working matrix.
4. Let  $\langle M, N \rangle$  receive the results of executing **procedure V:15** on the pair  $\langle 2, 2 \rangle$  and the following procedure:
  - (a) Execute **procedure V:13** on  $A'$ .
5. Using (4), verify that  $M$  is empty.
6. Using (4) and (5), verify that  $AN_* = M_* AN_* = A'$ .
7. Using (6), verify that  $A = A 1_n = AN_* N^{-1}_* = A' N^{-1}_*$ .
8. Using (4), verify that  $A'_{0,1} = 0$ .
9. Using (4) and (7), verify that  $A_{0,0} = A'_{0,0} N^{-1}_*{}_{0,0} + A'_{0,1} N^{-1}_*{}_{1,0} = A'_{0,0} N^{-1}_*{}_{0,0}$ .

10. Using (4) and (7), verify that  $A_{1,1} = A_{0,1} = A'_{0,0}N^{-1}_{*0,1} + A'_{0,1}N^{-1}_{*1,1} = A'_{0,0}N^{-1}_{*0,1}$ .
11. Using (2), verify that  $A_{1,0} = 0$ .
12. Using (6) and (11), verify that  $A'_{1,0} = A_{1,0}N_{*0,0} + A_{1,1}N_{*1,0} = A_{1,1}N_{*1,0} = A'_{0,0}N^{-1}_{*0,1}N_{*1,0}$ .
13. **Using (6) and (11), verify that  $A'_{1,1} = A_{1,0}N_{*0,1} + A_{1,1}N_{*1,1} = A_{1,1}N_{*1,1} = A'_{0,0}N^{-1}_{*0,1}N_{*1,1}$ .**
14. Subtract  $N^{-1}_{*0,1}N_{*1,0}$  times row 0 from row 1.
15. Now using (14) and (12), verify that  $A'_{1,0} = 0$ .
16. **Therefore verify that  $A'$  is a  $2 \times 2$  diagonal matrix.**
17. **Let  $A = A'$ .**
18. **Yield  $\langle N^{-1}_{*0,1}N_{*1,1}, N^{-1}_{*0,0} \rangle$ .**

## Procedure V:20(4.08)

### Objective

Choose a  $m \times n$  matrix,  $A$  such that  $\min(m, n) > 0$ . The objective of the following instructions is to define a list of polynomials  $u$  and transform  $A$  into an  $m \times n$  diagonal matrix such that  $A_{k,k} = u_k A_{0,0}$  for  $k$  in  $[0 : \min(m, n)]$  by a sequence of operations whereby either a polynomial times any of the columns is added to a different column, or a polynomial times any of the rows is added to a different row.

### Implementation

1. Let  $u = \langle 1 \rangle$ .
2. Execute **procedure V:14** on  $A$ .
3. Verify that  $A$  is an  $m \times n$  diagonal matrix.
4. For  $j$  in  $[1 : \min(m, n)]$ , do the following:
  - (a) Using (h), verify that  $A_{k,k} = u_k A_{0,0}$  for  $k$  in  $[0 : j]$ .
  - (b) Set  $A' = A$ .
  - (c) Execute **procedure V:19** on  $A'_{\langle 0,j \rangle, \langle 0,j \rangle}$  and let  $\langle u_j, v \rangle$  receive.

- (d) Using (c), verify that  $A$  and  $A'$  are the same modulo positions  $\langle 0, 0 \rangle$  and  $\langle j, j \rangle$ .
  - (e) Therefore verify that  $A'$  is an  $m \times n$  diagonal matrix.
  - (f) Also, using (c), verify that  $A'_{j,j} = u_j A'_{0,0}$ .
  - (g) Also, for  $k$  in  $[1 : j]$ , do the following:
    - i. Using (a), (c), and (d), verify that  $A'_{k,k} = A_{k,k} = u_k A_{0,0} = u_k A'_{0,0} v$ .
    - ii. Set  $u_k = u_k v$ .
    - iii. Hence verify that  $A'_{k,k} = u_k A'_{0,0}$ .
  - (h) Therefore verify that  $A_{k,k} = u_k A_{0,0}$  for  $k$  in  $[0 : j + 1]$ .
  - (i) Now let  $A = A'$ .
5. **Hence using (4h), verify that  $A_{k,k} = u_k A_{0,0}$  for  $k$  in  $[0 : \min(m, n)]$ .**
  6. **Also, using (4e), verify that  $A$  is an  $m \times n$  diagonal matrix.**
  7. **Yield  $\langle u \rangle$ .**

## Procedure V:21(4.09)

### Objective

Choose a  $m \times n$  matrix,  $A$ , and a  $n \times k$  matrix,  $B$ . Choose integers  $0 \leq a < m$ ,  $0 \leq b < n$ , and  $0 \leq c < k$ . The objective of the following instructions is to show that

1.  $(AB)_{[0:a],[0:c]} = A_{[0:a],[0:b]}B_{[0:b],[0:c]} + A_{[0:a],[b:n]}B_{[b:n],[0:c]}$
2.  $(AB)_{[0:a],[c:k]} = A_{[0:a],[0:b]}B_{[0:b],[c:k]} + A_{[0:a],[b:n]}B_{[b:n],[c:k]}$
3.  $(AB)_{[a:m],[0:c]} = A_{[a:m],[0:b]}B_{[0:b],[0:c]} + A_{[a:m],[b:n]}B_{[b:n],[0:c]}$
4.  $(AB)_{[a:m],[c:k]} = A_{[a:m],[0:b]}B_{[0:b],[c:k]} + A_{[a:m],[b:n]}B_{[b:n],[c:k]}$

### Implementation

1. For each  $0 \leq i < a$ , do the following:
  - (a) For each  $0 \leq j < c$ , do the following:

- i. Verify that  $(AB)_{i,j} = \sum_p^{[0:n]} A_{i,p} B_{p,j} = \sum_p^{[0:b]} A_{i,p} B_{p,j} + \sum_p^{[b:n]} A_{i,p} B_{p,j} = \sum_p^{[0:b]} (A_{[0:a],[0:b]} )_{i,p} (B_{[0:b],[0:c]} )_{p,j} + \sum_p^{[0:n-b]} (A_{[0:a],[b:n]} )_{i,p} (B_{[b:n],[0:c]} )_{p,j} = (A_{[0:a],[0:b]} B_{[0:b],[0:c]} )_{i,j} + (A_{[0:a],[b:n]} B_{[b:n],[0:c]} )_{i,j}$
2. **Therefore verify that**  $(AB)_{[0:a],[0:c]} = A_{[0:a],[0:b]} B_{[0:b],[0:c]} + A_{[0:a],[b:n]} B_{[b:n],[0:c]}$
3. **Using computations analogous to (1) and (2), show items (2), (3), and (4) of the objective.**

#### Declaration V:14

The phrase "number of rows of  $A$ " and the notation  $\text{rows}(A)$ , where  $A$  is an  $m \times n$  matrix, will be used as a shorthand for  $m$ .

#### Declaration V:15

The phrase "number of columns of  $A$ " and the notation  $\text{cols}(A)$ , where  $A$  is an  $m \times n$  matrix, will be used as a shorthand for  $n$ .

#### Declaration V:16

The notation  $\text{diag}(C)$ , where  $C$  is a list of rational square matrices, will be used to refer to the result yielded by executing the following instructions:

1. Let  $E$  be a  $0 \times 0$  matrices.
2. Now for  $i$  in  $[0 : |C|]$ :
  - (a) Add  $\text{cols}(C_i)$  columns filled with zeros to the right end of  $E$ .
  - (b) Add  $\text{rows}(C_i)$  rows filled with zeros to the bottom end of  $E$ .
  - (c) Set the bottom-right corner of  $E$  equal to  $C_i$ .
3. **Yield the tuple**  $\langle E \rangle$ .

#### Procedure V:22(4.10)

##### Objective

Choose a  $m \times n$  matrix,  $A$ . Let  $A_{-1,-1} = 1$ . The objective of the following instructions is to construct

the list of polynomials  $v$  and transform  $A$  into an  $m \times n$  diagonal matrix such that  $A_{k,k} = v_k A_{k-1,k-1}$  for  $k$  in  $[0 : \min(m, n)]$  by a sequence of operations whereby either a polynomial times any of the columns is added to a different column, or a polynomial times any of the rows is added to a different row.

##### Implementation

1. If  $\min(m, n) = 0$ , then do the following:
  - (a) **Verify that  $A$  is an  $m \times n$  diagonal matrix.**
  - (b) **Yield**  $\langle \rangle$ .
2. Otherwise do the following:
  - (a) Apply **procedure V:20** on  $A$ , and let  $\langle u \rangle$  receive.
  - (b) Verify that  $A$  is an  $m \times n$  diagonal matrix.
  - (c) Verify that  $A_{k,k} = u_k A_{0,0}$  for  $k$  in  $[0 : \min(m, n)]$ .
  - (d) Let  $B, C$  be an  $(m-1) \times (n-1)$  diagonal matrix with  $u_{1:|u|}$  on the diagonal.
  - (e) Let  $\langle M, N \rangle$  receive the results of executing **procedure V:15** on the pair  $\langle m-1, n-1 \rangle$  and the following procedure:
    - i. Execute **procedure V:22** on  $C$  and let  $\langle w \rangle$  receive.
  - (f) Therefore verify that  $C$  is an  $(m-1) \times (n-1)$  diagonal matrix.
  - (g) Also verify that  $C = M_* B N_*$ .
  - (h) Let  $C_{-1,-1} = 1$ .
  - (i) Now using (ei), verify that  $C_{k,k} = w_k C_{k-1,k-1}$  for  $k$  in  $[0 : \min(m, n) - 1]$ .
  - (j) Therefore using (c), verify that  $A_{0,0} C = M_* (A_{0,0} B) N_* = M_* A_{[1:m],[1:n]} N_*$ .
  - (k) Premultiply  $A$  by  $\text{diag}(1, M_k)$  for  $k$  in  $[|M| : 0]$ .
  - (l) Postmultiply  $A$  by  $\text{diag}(1, N_k)$  for  $k$  in  $[0 : |N|]$ .
  - (m) Now verify that  $A_{[1:m],[1:n]} = A_{0,0} C$ .
  - (n) Now let  $u = \langle A_{0,0} \rangle \frown w$ .

- (o) **Therefore verify that**  $A_{k,k} = u_k A_{k-1,k-1}$   
**for**  $k$  **in**  $[0 : \min(m, n)]$ .
- (p) **Yield the tuple**  $\langle u \rangle$ .

# Chapter 16

## Compound Matrices

### Declaration V:17

The notation  $\text{det}(A)$ , where  $A$  is a  $m \times m$  matrix, will be used to refer to the result yielded by executing the following instructions:

1. If  $m = 0$ , then do the following:
  - (a) **Yield the tuple**  $\langle 1 \rangle$ .
2. Otherwise, do the following:
  - (a) Let  $h_r = A_{[0:r] \cap [r+1:m], [1:m]}$  for  $r$  in  $[0 : m]$ .
  - (b) **Yield the tuple**  $\langle \sum_r^{[0:m]} (-1)^r A_{r,0} \text{det}(h_r) \rangle$ .

### Procedure V:23(4.11)

#### Objective

Choose a polynomial  $p$ . Choose two  $1 \times m$  matrices,  $B$  and  $C$ . Choose an integer  $0 \leq i < m$ . Choose a  $m \times m$  matrix,  $A$ , such that its  $i^{\text{th}}$  row is  $B + pC$ . Let  $A'$  be  $A$  but with the  $i^{\text{th}}$  row replaced by  $B$  and let  $A''$  be  $A$  but with the  $i^{\text{th}}$  row replaced by  $C$ . The objective of the following instructions is to show that  $\text{det}(A) = \text{det}(A') + p \text{det}(A'')$ .

#### Implementation

1. If  $m = 1$ , then do the following:
  - (a) Verify that  $i = 0$ .
  - (b) **Therefore verify that**  $\text{det}(A) = A_{0,0} = B_{0,0} + pC_{0,0} = \text{det}(A') + p \text{det}(A'')$ .
2. Otherwise, do the following:

- (a) For  $r$  in  $[0 : i]$ , do the following:
  - i. Verify that  $(A_{[0:r] \cap [r+1:m], [1:m]})_{i-1,*} = B + pC$ .
  - ii. Verify that  $A'_{[0:r] \cap [r+1:m], [1:m]}$  is  $A_{[0:r] \cap [r+1:m], [1:m]}$  with row  $i-1$  replaced by  $B$ .
  - iii. Verify that  $A''_{[0:r] \cap [r+1:m], [1:m]}$  is  $A_{[0:r] \cap [r+1:m], [1:m]}$  with row  $i-1$  replaced by  $C$ .
  - iv. Execute **procedure V:23** on  $\langle p, B, C, i-1, A_{[0:r] \cap [r+1:m], [1:m]} \rangle$ .
  - v. Therefore verify that  $\text{det}(A_{[0:r] \cap [r+1:m], [1:m]}) = \text{det}(A'_{[0:r] \cap [r+1:m], [1:m]}) + p \text{det}(A''_{[0:r] \cap [r+1:m], [1:m]})$ .
- (b) For  $r$  in  $[i+1 : m]$ , do the following:
  - i. Verify that  $(A_{[0:r] \cap [r+1:m], [1:m]})_{i,*} = B + pC$ .
  - ii. Verify that  $A'_{[0:r] \cap [r+1:m], [1:m]}$  is  $A_{[0:r] \cap [r+1:m], [1:m]}$  with row  $i$  replaced by  $B$ .
  - iii. Verify that  $A''_{[0:r] \cap [r+1:m], [1:m]}$  is  $A_{[0:r] \cap [r+1:m], [1:m]}$  with row  $i$  replaced by  $C$ .
  - iv. Execute **procedure V:23** on  $\langle p, B, C, i, A_{[0:r] \cap [r+1:m], [1:m]} \rangle$ .
  - v. Therefore verify that  $\text{det}(A_{[0:r] \cap [r+1:m], [1:m]}) = \text{det}(A'_{[0:r] \cap [r+1:m], [1:m]}) + p \text{det}(A''_{[0:r] \cap [r+1:m], [1:m]})$ .
- (c) Therefore using (av) and (bv), verify that
  - i.  $= \sum_r^{[0:m]} (-1)^r A_{r,0} \text{det}(A_{[0:r] \cap [r+1:m], [1:m]})$

- ii.  $= \sum_r^{[0:i]} (-1)^r A_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]}) + (-1)^i A_{i,0} \det(A_{[0:i] \cap [i+1:m], [1:m]}) + \sum_r^{[i+1:m]} (-1)^r A_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]})$
  - iii.  $= \sum_r^{[0:i]} (-1)^r A_{r,0} (\det(A'_{[0:r] \cap [r+1:m], [1:m]}) + p \det(A''_{[0:r] \cap [r+1:m], [1:m]})) + (-1)^i (A'_{i,0} + p A''_{i,0}) \det(A_{[0:i] \cap [i+1:m], [1:m]}) + \sum_r^{[i+1:m]} (-1)^r A_{r,0} (\det(A'_{[0:r] \cap [r+1:m], [1:m]}) + p \det(A''_{[0:r] \cap [r+1:m], [1:m]}))$
  - iv.  $= \sum_r^{[0:i]} (-1)^r A_{r,0} \det(A'_{[0:r] \cap [r+1:m], [1:m]}) + (-1)^i A'_{i,0} \det(A_{[0:i] \cap [i+1:m], [1:m]}) + \sum_r^{[i+1:m]} (-1)^r A_{r,0} \det(A'_{[0:r] \cap [r+1:m], [1:m]}) + \sum_r^{[0:i]} (-1)^r A_{r,0} p \det(A''_{[0:r] \cap [r+1:m], [1:m]}) + (-1)^i p A''_{i,0} \det(A_{[0:i] \cap [i+1:m], [1:m]}) + \sum_r^{[i+1:m]} (-1)^r A_{r,0} p \det(A''_{[0:r] \cap [r+1:m], [1:m]})$
  - v.  $= \sum_r^{[0:m]} (-1)^r A'_{r,0} \det(A'_{[0:r] \cap [r+1:m], [1:m]}) + p \sum_r^{[0:m]} (-1)^r A''_{r,0} \det(A''_{[0:r] \cap [r+1:m], [1:m]})$
  - vi.  $= \det(A') + p \det(A'')$
  - (e)  $= \sum_r^{[0:m]} (-1)^r (A')_{r,0} \det(A'_{[0:r] \cap [r+1:m], [1:m]}) + p \sum_r^{[0:m]} (-1)^r (A'')_{r,0} \det(A''_{[0:r] \cap [r+1:m], [1:m]})$
  - (f)  $= \det(A') + p \det(A'')$
2. Otherwise, do the following:
- (a) For  $r$  in  $[0 : m]$ , do the following:
    - i. Execute **procedure V:24** on  $\langle p, B_{[0:r] \cap [r+1:m], 0}, C_{[0:r] \cap [r+1:m], 0}, i - 1, A_{[0:r] \cap [r+1:m], [1:m]} \rangle$ .
    - ii. Therefore verify that  $\det(A_{[0:r] \cap [r+1:m], [1:m]}) = \det(A'_{[0:r] \cap [r+1:m], [1:m]}) + p \det(A''_{[0:r] \cap [r+1:m], [1:m]})$ .
  - (b) Therefore using (a), verify that  $\det(A)$ 
    - i.  $= \sum_r^{[0:m]} (-1)^r A_{r,0} \cdot \det(A_{[0:r] \cap [r+1:m], [1:m]})$
    - ii.  $= \sum_r^{[0:m]} (-1)^r A_{r,0} (\det(A'_{[0:r] \cap [r+1:m], [1:m]}) + p \det(A''_{[0:r] \cap [r+1:m], [1:m]}))$
    - iii.  $= \sum_r^{[0:m]} (-1)^r A'_{r,0} \det(A'_{[0:r] \cap [r+1:m], [1:m]}) + \sum_r^{[0:m]} (-1)^r A''_{r,0} p \det(A''_{[0:r] \cap [r+1:m], [1:m]})$
    - iv.  $= \det(A') + p \det(A'')$

## Procedure V:24(4.12)

### Objective

Choose a polynomial  $p$ . Choose two  $m \times 1$  matrices,  $B$  and  $C$ . Choose an integer  $0 \leq i < m$ . Choose a  $m \times m$  matrix,  $A$ , such that its  $i^{th}$  column is  $B + pC$ . Let  $A'$  be  $A$  but with the  $i^{th}$  column replaced by  $B$  and let  $A''$  be  $A$  but with the  $i^{th}$  column replaced by  $C$ . The objective of the following instructions is to show that  $\det(A) = \det(A') + p \det(A'')$ .

### Implementation

1. If  $i = 0$ , then verify that  $\det(A)$ 
  - (a)  $= \sum_r^{[0:m]} (-1)^r A_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]})$
  - (b)  $= \sum_r^{[0:m]} (-1)^r (B + pC)_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]})$
  - (c)  $= \sum_r^{[0:m]} (-1)^r (B)_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]}) + \sum_r^{[0:m]} (-1)^r (pC)_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]})$
  - (d)  $= \sum_r^{[0:m]} (-1)^r (B)_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]}) + p \sum_r^{[0:m]} (-1)^r (C)_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]})$

## Procedure V:25(4.13)

### Objective

Choose a  $m \times m$  matrix,  $A$ . Choose an integer  $0 < i < m$ . Let  $A'$  be  $A$  with rows  $i - 1$  and  $i$  swapped. The objective of the following instructions is to show that  $\det(A') = -\det(A)$ .

### Implementation

1. If  $m = 2$ , then do the following:
  - (a) Verify that  $i = 1$ .
  - (b) Therefore verify that  $\det(A') = A'_{0,0} A'_{1,1} - A'_{1,0} A'_{0,1} = A_{1,0} A_{0,1} - A_{0,0} A_{1,1} = -\det(A)$ .
2. Otherwise do the following:
  - (a) For  $r$  in  $[0 : i - 1]$ , do the following:
    - i. Verify that  $A_{[0:r] \cap [r+1:m], [1:m]}$  is the same as  $A'_{[0:r] \cap [r+1:m], [1:m]}$  but with rows  $i - 2$  and  $i - 1$  swapped.
    - ii. Execute **procedure V:25** on  $\langle A_{[0:r] \cap [r+1:m], [1:m]}, i - 1 \rangle$ .

- iii. Hence verify that  $\det(A'_{[0:r] \cap [r+1:m], [1:m]}) = -\det(A_{[0:r] \cap [r+1:m], [1:m]})$ .
- (b) For  $r$  in  $[i+1 : m]$ , do the following:
- Verify that  $A_{[0:r] \cap [r+1:m], [1:m]}$  is the same as  $A'_{[0:r] \cap [r+1:m], [1:m]}$  but with rows  $i-1$  and  $i$  swapped.
  - Execute **procedure V:25** on  $\langle A_{[0:r] \cap [r+1:m], [1:m]}, i \rangle$ .
  - Hence verify that  $\det(A'_{[0:r] \cap [r+1:m], [1:m]}) = -\det(A_{[0:r] \cap [r+1:m], [1:m]})$ .
- (c) Verify that  $\det(A)$
- $= \sum_r^{[0:m]} (-1)^r A_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]})$
  - $= \sum_r^{[0:i-1]} (-1)^r A_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]}) + (-1)^{i-1} A_{i-1,0} \det(A_{[0:i-1] \cap [i:m], [1:m]}) + (-1)^i A_{i,0} \det(A_{[0:i] \cap [i+1:m], [1:m]}) + \sum_r^{[i+1:m]} (-1)^r A_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]})$
  - $= -\sum_r^{[0:i-1]} (-1)^r A'_{r,0} \det(A'_{[0:r] \cap [r+1:m], [1:m]}) - (-1)^i A'_{i,0} \det(A'_{[0:i] \cap [i+1:m], [1:m]}) - (-1)^{i-1} A'_{i-1,0} \det(A'_{[0:i-1] \cap [i:m], [1:m]}) - \sum_r^{[i+1:m]} (-1)^r A'_{r,0} \det(A'_{[0:r] \cap [r+1:m], [1:m]})$
  - $= -\sum_r^{[0:m]} (-1)^r A'_{r,0} \det(A'_{[0:r] \cap [r+1:m], [1:m]})$
  - $= -\det(A')$ .

## Procedure V:26(4.14)

### Objective

Choose a  $m \times m$  matrix,  $A$ . Choose an integer  $0 < i < m$ . Let  $A'$  be  $A$  with columns  $i-1$  and  $i$  swapped. The objective of the following instructions is to show that  $\det(A') = -\det(A)$ .

### Implementation

- If  $i = 1$ , then verify that  $\det(A)$ 
  - $= \sum_r^{[0:m]} (-1)^r A_{r,0} \det(A_{[0:r] \cap [r+1:m], [1:m]})$
  - $= \sum_r^{[0:m]} (-1)^r A_{r,0} \sum_t^{[r+1:m]} (-1)^{t-1} A_{t,1} * \det(A_{[0:r] \cap [r+1:t] \cap [t+1:m], [2:m]}) + \sum_t^{[0:m]} (-1)^t A_{t,0} \sum_r^{[0:t]} (-1)^r A_{r,1} * \det(A_{[0:r] \cap [r+1:t] \cap [t+1:m], [2:m+1]})$

- $= \sum_t^{[0:m]} (-1)^{t-1} A_{t,1} \sum_r^{[0:t]} (-1)^r A_{r,0} * \det(A_{[0:r] \cap [r+1:t] \cap [t+1:m], [2:m+1]}) + \sum_r^{[0:m]} (-1)^r A_{r,1} \sum_t^{[r+1:m]} (-1)^t A_{t,0} * \det(A_{[0:r] \cap [r+1:t] \cap [t+1:m], [2:m+1]})$
  - $= \sum_t^{[0:m]} (-1)^{t-1} A'_{t,0} \sum_r^{[0:t]} (-1)^r A'_{r,1} * \det(A'_{[0:r] \cap [r+1:t] \cap [t+1:m], [2:m+1]}) + \sum_r^{[0:m]} (-1)^r A'_{r,0} \sum_t^{[r+1:m]} (-1)^t A'_{t,1} * \det(A'_{[0:r] \cap [r+1:t] \cap [t+1:m], [2:m]})$
  - $= -(\sum_r^{[0:m]} (-1)^r A'_{r,0} \sum_t^{[r+1:m]} (-1)^{t-1} A'_{t,1} * \det(A'_{[0:r] \cap [r+1:t] \cap [t+1:m], [2:m]}) + \sum_t^{[0:m]} (-1)^t A'_{t,0} \sum_r^{[0:t]} (-1)^r A'_{r,1} * \det(A'_{[0:r] \cap [r+1:t] \cap [t+1:m], [2:m+1]})$
  - $= -\det(A')$ .
2. Otherwise do the following:
- Verify that  $i > 1$ .
  - For  $r$  in  $[0 : m]$ , do the following:
    - Execute **procedure V:26** on  $\langle i-1, A_{[0:r] \cap [r+1:m], [1:m]} \rangle$ .
    - Therefore verify that  $\det(A_{[0:r] \cap [r+1:m], [1:m]}) = -\det(A'_{[0:r] \cap [r+1:m], [1:m]})$ .
  - Therefore using (bii), verify that  $\det(A) = \sum_r^{[0:m]} (-1)^r A_{r,0} \cdot \det(A_{[0:r] \cap [r+1:m], [1:m]}) = \sum_r^{[0:m]} (-1)^r A'_{r,0} \cdot (-\det(A'_{[0:r] \cap [r+1:m], [1:m]})) = -\det(A')$ .

## Procedure V:27(4.15)

### Objective

Choose integers  $0 < i < m$ . Choose a  $m \times m$  matrix,  $A$ , such that columns  $i-1$  and  $i$  are the same. The objective of the following instructions is to show that  $\det(A) = 0$ .

### Implementation

- Let  $A'$  be  $A$  with columns  $i-1$  and  $i$  swapped.
- Execute **procedure V:26** on  $\langle A, i \rangle$ .
- Also, verify that  $A' = A$ .
- Therefore verify that  $\det(A) = \det(A') = -\det(A)$ .

5. **Therefore verify that**  $\det(A) = 0$ .

### Procedure V:28(4.16)

#### Objective

Choose integers  $0 < i < m$ . Choose a  $m \times m$  matrix,  $A$ , such that rows  $i-1$  and  $i$  are the same. The objective of the following instructions is to show that  $\det(A) = 0$ .

#### Implementation

Instructions are analogous to those of **procedure V:27**.

### Procedure V:29(4.17)

#### Objective

Choose integers  $0 \leq i < m$ . Choose an integer  $-i \leq j < m-i$ . Choose a  $m \times m$  matrix,  $A$ . Let  $A'$  be  $A$  but with column  $i$  moved  $j$  places. The objective of the following instructions is to show that  $\det(A') = (-1)^j \det(A)$ .

#### Implementation

1. Let  $B = \langle A \rangle$ .
2. For  $k$  in  $[i : i+j]$ , do the following:
  - (a) Let  $B_{|B|}$  be the result of swapping columns  $k$  and  $k+1$  of  $B_{|B|-1}$ .
  - (b) Using **procedure V:26**, verify that  $\det(B_{|B|-1}) = -\det(B_{|B|-2})$ .
3. Verify that  $A' = B_{|B|-1}$ .
4. **Therefore verify that**  $\det(A') = \det(B_{|B|-1}) = (-1)^1 \det(B_{|B|-2}) = \dots = (-1)^j \det(B_0) = (-1)^j \det(A)$ .

### Procedure V:30(4.18)

#### Objective

Choose integers  $0 \leq i < m$ . Choose an integer  $-i \leq j < m-i$ . Choose a  $m \times m$  matrix,  $A$ . Let

$A'$  be  $A$  but with row  $i$  moved  $j$  places. The objective of the following instructions is to show that  $\det(A') = (-1)^j \det(A)$ .

#### Implementation

Instructions are analogous to those of **procedure V:29**.

#### Declaration V:18

The notation  $C_k(A)$ , where  $A$  is a  $m \times n$  matrix and  $k$  is an integer such that  $0 \leq k \leq \min(m, n)$ , will be used to refer to the  $\binom{m}{k} \times \binom{n}{k}$  matrix with the following specification:

1. The rows are labeled by the colexicographically sorted list of increasing length- $k$  sequences whose elements are picked from  $[0 : m]$ .
2. The columns are labeled by the colexicographically sorted list of increasing length- $k$  sequences whose elements are picked from  $[0 : n]$ .
3. For each row label  $I$ : For each column label  $J$ : The entry at position  $(I, J)$  is  $\det(A_{I,J})$ .

#### Declaration V:19

The notation  $A_{I,J}$  will be used to refer to the entry of  $A$  with row label  $I$  and column label  $J$ .

### Procedure V:31(4.19)

#### Objective

Choose two integers  $0 \leq k \leq m$ . The objective of the following instructions is to show that  $C_k(1_m) = 1_{\binom{m}{k}}$ .

#### Implementation

1. For each row label  $I$  of  $C_k(1_m)$ , for each column label  $J$  of  $C_k(1_m)$ , do the following:
  - (a) If  $I = J$ , then do the following:



- i. Verify that  $((1_m)_{I,J})_{i,j} = ((1_m)_{J,J})_{i,j} = (1_m)_{J_i,J_j} = [J_i = J_j] = [i = j]$  for  $0 \leq i < k$ , for  $0 \leq j < k$ .
  - ii. Therefore verify that  $(C_k(1_m))_{\underline{I},\underline{J}} = 1_k$ .
  - iii. **Therefore verify that**  $(C_k(1_m))_{\underline{I},\underline{J}} = \det((1_m)_{I,J}) = \det(1_k) = 1$ .
- (b) Otherwise, do the following:
- i. Verify that  $I \neq J$ .
  - ii. Let  $i$  be the index of an element of  $I$  that is not an element of  $J$ .
  - iii. Now verify that  $(1_m)_{I_i,j} = [I_i = j] = 0$ , for each  $j$  in  $J$ .
  - iv. Therefore verify that  $((1_m)_{I,J})_{i,*} = 0_{1 \times k}$ .
  - v. **Therefore verify that**  $(C_k(1_m))_{\underline{I},\underline{J}} = \det((1_m)_{I,J}) = 0$ .
2. **Therefore verify that**  $C_k(1_m) = 1_{\binom{m}{k}}$ .

## Procedure V:32(4.20)

### Objective

Choose an integer  $0 \leq k \leq \min(m, n)$ . Choose a  $m \times m$  tilt,  $A$ , such that the off diagonal entry is the polynomial  $p$  at  $(i, j)$ . Also choose a  $m \times n$  matrix,  $B$ . The objective of the following instructions is to construct a  $\binom{m}{k} \times \binom{m}{k}$  matrix  $D$  such that  $C_k(AB) = DC_k(B)$ .

### Implementation

1. Let  $D = C_k(1_m) = 1_{\binom{m}{k}}$ .
2. Verify that  $AB$  equals  $B$ , but with its row  $i$  having  $p$  times  $B$ 's row  $j$  added to it.
3. Go through the row labels,  $I$ , of  $C_k(AB)$  and do the following:
  - (a) If  $i \notin I$ , then do the following:
    - i. Verify that  $(AB)_{I,*} = B_{I,*}$ .
    - ii. Therefore for each column label  $J$ , verify that  $C_k(AB)_{\underline{I},\underline{J}} = \det((AB)_{I,J}) = \det(B_{I,J}) = C_k(B)_{\underline{I},\underline{J}}$ .
    - iii. **Therefore verify that**  $(C_k(AB))_{\underline{I},*} = (C_k(B))_{\underline{I},*}$ .

- (b) Otherwise, if  $i \in I$ , then:
  - i. Let  $I'$  be  $I$  but with an in-place replacement of  $i$  by  $j$ .
  - ii. For each column label  $J$ : Using **procedure V:24**, verify that  $C_k(AB)_{\underline{I},\underline{J}} = \det((AB)_{I,J}) = \det(B_{I,J}) + p * \det(B_{I',J})$ .
  - iii. If  $j \in I$ , then do the following:
    - A. Verify that the sequence  $I'$  contains two  $j$ 's.
    - B. For each column label  $J$ : Using **procedure V:28** verify that  $\det(B_{I',J}) = 0$ .
    - C. Therefore for each column label  $J$ : verify that  $C_k(AB)_{\underline{I},\underline{J}} = \det(B_{I,J}) = C_k(B)_{\underline{I},\underline{J}}$ .
    - D. **Therefore verify that**  $C_k(AB)_{\underline{I},*} = C_k(B)_{\underline{I},*}$ .
  - iv. Otherwise if  $j \notin I$ , do the following:
    - A. Let  $l$  be the signed number of places that the  $j$  introduced above needs to be moved in order to make  $I'$  an increasing sequence.
    - B. Let  $I''$  be obtained from  $I'$  by moving the integer  $j$  in  $I'$  by  $l$  places.
    - C. For each column label  $J$ : Using **procedure V:30**, verify that  $\det(B_{I',J}) = (-1)^l \det(B_{I'',J})$ .
    - D. Therefore for each column label  $J$ : Verify that  $C_k(AB)_{\underline{I},\underline{J}} = \det(B_{I,J}) + p * \det(B_{I',J}) = \det(B_{I,J}) + (-1)^l p * \det(B_{I'',J})$ .
    - E. Verify that  $I''$  is a row label of  $C_k(B)$ .
    - F. Therefore for each column label  $J$ : Verify that  $C_k(AB)_{\underline{I},\underline{J}} = \det(B_{I,J}) + (-1)^l p * \det(B_{I'',J}) = C_k(B)_{\underline{I},\underline{J}} + (-1)^l p * C_k(B)_{\underline{I'',J}}$ .
    - G. **Therefore verify that**  $(C_k(AB))_{\underline{I},*} = (C_k(B))_{\underline{I},*} + (-1)^l p (C_k(B))_{\underline{I'',*}}$ .
    - H. **Set**  $D_{\underline{I},\underline{I''}}$  **to**  $(-1)^l p$ .
- (c) **Therefore verify that**  $C_k(AB)_{\underline{I},*} = D_{\underline{I},*} C_k(B)$ .
- 4. **Therefore verify that**  $C_k(AB) = DC_k(B)$ .

## 5. Yield $\langle D \rangle$ .

### Procedure V:33(4.21)

#### Objective

Choose an  $m \times n$  diagonal matrix,  $A$ . Also choose an  $n \times n$  matrix,  $B$ . Also choose an integer  $0 \leq k \leq \min(m, n)$ . The objective of the following instructions is to construct an  $\binom{m}{k} \times \binom{n}{k}$  diagonal matrix  $D$  such that  $C_k(AB) = DC_k(B)$ .

#### Implementation

1. Let  $D = C_k(0_{m \times n}) = 0_{\binom{m}{k} \times \binom{n}{k}}$ .
2. Verify that  $AB$  equals  $B_{[0:\min(m,n)],*}$  with each row  $i$  multiplied by  $A_{i,i}$ .
3. Go through the row labels,  $I$ , of  $C_k(AB)$  and do the following:
  - (a) If  $I_k < \min(m, n)$ , then do the following:
    - i. Verify that every element of  $I$  is less than  $\min(m, n)$ .
    - ii. Let  $A_0 = A$ .
    - iii. For  $i$  in  $[0 : k]$ : Let  $A_{i+1}$  equal  $A_i$  but with position  $(I_i, I_i)$  set to 1.
    - iv. For each column label  $J$ : Repeatedly using **procedure V:24**, verify that  $C_k(AB)_{I,J}$ 
      - A.  $= \det((AB)_{I,J})$
      - B.  $= \det((A_0B)_{I,J})$
      - C.  $= A_{I_0,I_0} \det((A_1B)_{I,J})$
      - D.  $= A_{I_0,I_0} A_{I_1,I_1} \det((A_2B)_{I,J})$
      - E.  $\vdots$
      - F.  $= A_{I_0,I_0} A_{I_1,I_1} \cdots A_{I_{k-1},I_{k-1}} \det((A_kB)_{I,J})$
      - G.  $= A_{I_0,I_0} A_{I_1,I_1} \cdots A_{I_{k-1},I_{k-1}} \det(B_{I,J})$
      - H.  $= A_{I_0,I_0} A_{I_1,I_1} \cdots A_{I_{k-1},I_{k-1}} C_k(B)_{\underline{I},\underline{J}}$ .
    - v. **Therefore verify that**  $(C_k(AB))_{\underline{I},*} = A_{I_1,I_1} A_{I_2,I_2} \cdots A_{I_k,I_k} * (C_k(B))_{\underline{I},*}$
    - vi. **Set**  $D_{\underline{I},\underline{I}}$  **to**  $A_{I_0,I_0} A_{I_1,I_1} \cdots A_{I_{k-1},I_{k-1}}$ .
  - (b) Otherwise if  $I_k \geq \min(m, n)$ , then do the following:

- i. Using the precondition, verify that  $A_{I_k,*} = 0_{1 \times n}$ .
- ii. Therefore verify that  $(AB)_{I_k,*} = 0_{1 \times n}$ .
- iii. Therefore verify that  $((AB)_{I,*})_{k,*} = 0_{1 \times n}$ .
- iv. Therefore for each column label  $J$ : verify that  $C_k(AB)_{\underline{I},\underline{J}} = \det((AB)_{I,J}) = 0$ .
- v. **Therefore verify that**  $(C_k(AB))_{\underline{I},*}$  **is zero.**
- (c) **Therefore verify that**  $C_k(AB)_{\underline{I},*} = D_{\underline{I},*} C_k(B)$ .

#### 4. Verify that $D$ is diagonal.

#### 5. Verify that $C_k(AB) = DC_k(B)$ .

#### 6. Yield $\langle D \rangle$ .

### Procedure V:34(4.22)

#### Objective

Choose an integer  $0 \leq k \leq \min(m, n)$ . Choose a  $m \times m$  tilt,  $A$ . Also choose a  $m \times n$  matrix,  $B$ . The objective of the following instructions is to show that  $C_k(AB) = C_k(A)C_k(B)$ .

#### Implementation

1. Execute **procedure V:32** on matrices  $A$  and  $1_m$  and let  $\langle D \rangle$  receive.
2. Using **procedure V:31**, verify that  $C_k(A) = C_k(A1_m) = DC_k(1_m) = D1_{\binom{m}{k}} = D$ .
3. Execute **procedure V:32** on  $\langle A, B \rangle$  and let  $\langle D' \rangle$  receive.
4. Verify that  $D' = D = C_k(A)$ .
5. **Therefore verify that**  $C_k(AB) = D' C_k(B) = C_k(A) C_k(B)$ .

### Procedure V:35(4.23)

#### Objective

Choose an integer  $0 \leq k \leq \min(m, n)$ . Choose an  $n \times n$  tilt,  $A$ . Also choose a  $m \times n$  matrix,  $B$ . The objective of the following instructions is to show that  $C_k(BA) = C_k(B)C_k(A)$ .

## Implementation

Instructions are analogous to those of [procedure V:34](#).

## Procedure V:36(4.24)

### Objective

Choose an integer  $0 \leq k \leq \min(m, n)$ . Choose an  $m \times n$  diagonal matrix,  $A$ . Also choose a  $n \times n$  matrix,  $B$ . The objective of the following instructions is to show that  $C_k(AB) = C_k(A)C_k(B)$ .

### Implementation

Instructions are analogous to those of [procedure V:34](#).

## Procedure V:37(4.25)

### Objective

Choose a  $m \times n$  matrix,  $A$ . Let  $D_{-1,-1} = 1$ . The objective of the following instructions is to construct a list of  $m \times m$  tilts,  $M$ , an  $m \times n$  diagonal matrix,  $D$ , a list of polynomials,  $v$ , and a list of  $n \times n$  tilts,  $N$ , such that  $M_*AN_* = D$ ,  $A = M^{-1}_*DN^{-1}_*$ , and  $D_{i,i} = v_iD_{i-1,i-1}$  for  $i$  in  $[0 : \min(m, n)]$ .

### Implementation

1. Let  $D$  be a copy of  $A$ .
2. Let  $\langle M, N \rangle$  receive the results of executing [procedure V:15](#) on the pair  $\langle m, n \rangle$  and the following procedure:
  - (a) Execute [procedure V:22](#) on the matrix  $D$  and let  $\langle v \rangle$  receive.
3. **Verify that**  $D_{i,i} = v_iD_{i-1,i-1}$  **for**  $i$  **in**  $[0 : \min(m, n)]$ .
4. **Verify that**  $M_*AN_* = D$ .
5. Hence verify that  $A = 1_m A 1_n = M^{-1}_*M_*AN_*N^{-1}_* = M^{-1}_*DN^{-1}_*$ .
6. **Yield the tuple**  $\langle M, D, v, N \rangle$ .

## Procedure V:38(4.26)

### Objective

Choose integers  $0 \leq k \leq \min(m, n, p)$ . Choose a  $m \times n$  matrix,  $A$ . Also choose a  $n \times p$  matrix,  $B$ . The objective of the following instructions is to show that  $C_k(AB) = C_k(A)C_k(B)$ .

### Implementation

1. Execute [procedure V:37](#) on  $A$  and let  $\langle M, D, , N \rangle$  receive.
2. Using repeated applications of [procedure V:36](#), verify that  $C_k(AB)$ 
  - (a)  $= C_k(M^{-1}_0 \cdots M^{-1}_{|M|-1} D N^{-1}_0 \cdots N^{-1}_{|N|-1} B)$
  - (b)  $= C_k(M^{-1}_0) \cdots C_k(M^{-1}_{|M|-1}) * C_k(D) * C_k(N^{-1}_0) \cdots C_k(N^{-1}_{|N|-1}) C_k(B)$
  - (c)  $= C_k(M^{-1}_0 \cdots M^{-1}_{|M|-1} D N^{-1}_0 \cdots N^{-1}_{|N|-1}) C_k(B)$
  - (d)  $= C_k(A) C_k(B)$ .

## Procedure V:39(4.27)

### Objective

Choose a  $m \times m$  matrix,  $A$ . Let  $D$  be a copy of  $A$ . Execute [procedure V:22](#) on  $D$ . The objective of the following instructions is to show that  $\det(A)$  is the product of the diagonal entries of  $D$ .

### Implementation

1. Execute [procedure V:37](#) on  $A$  and let  $\langle M, D, , N \rangle$  receive.
2. Using [procedure V:38](#), verify that  $\det(A)$ 
  - (a)  $= C_m(A)$
  - (b)  $= C_m(M^{-1}_0 \cdots M^{-1}_{|M|-1} D N^{-1}_0 \cdots N^{-1}_{|N|-1})$
  - (c)  $= C_m(M^{-1}_0) \cdots C_m(M^{-1}_{|M|-1}) C_m(D) C_m(N^{-1}_0) \cdots C_m(N^{-1}_{|N|-1})$
  - (d)  $= 1 \cdots 1 C_m(D) 1 \cdots 1 = C_m(D)$
  - (e)  $= \det(D)$
  - (f)  $= \prod_r^{[0:m]} D_{r,r}$ .

### Declaration V:20

The notation  $A^T$ , where  $A$  is a  $m \times n$  matrix, will be used to refer to the  $n \times m$  matrix such that  $A^T_{i,j} = A_{j,i}$  for  $i$  in  $[0 : n]$ , for  $j$  in  $[0 : m]$ .

### Procedure V:40(4.28)

#### Objective

Choose a  $m \times n$  matrix,  $A$ , and a  $n \times k$  matrix,  $B$ . The objective of the following instructions is to show that  $B^T A^T = (AB)^T$ .

#### Implementation

1. Verify that  $B^T A^T$  and  $(AB)^T$  have dimensions  $k \times m$ .
2. For  $i$  in  $[0 : k]$ : For  $j$  in  $[0 : m]$ :
  - (a) Verify that  $(B^T A^T)_{i,j} = \sum_l^{[0:n]} B_{l,i} A_{j,l} = \sum_l^{[0:n]} A_{j,l} B_{l,i} = (AB)_{j,i} = ((AB)^T)_{i,j}$ .
3. Therefore verify that  $B^T A^T = (AB)^T$ .

### Procedure V:41(4.29)

#### Objective

Choose a  $m \times m$  matrix,  $A$ . The objective of the following instructions is to show that  $\det(A^T) = \det(A)$ .

#### Implementation

1. Execute **procedure V:37** on  $A$  and let  $\langle M, D, , N \rangle$  receive.
2. Therefore using procedures **procedure V:39** and **procedure V:40**, verify that  $\det(A^T)$ 
  - (a)  $= \det((M^{-1}_0 \cdots M^{-1}_{|M|-1} D N^{-1}_0 \cdots N^{-1}_{|N|-1})^T)$
  - (b)  $= \det((N^{-1}_{|N|-1})^T \cdots (N^{-1}_0)^T D^T (M^{-1}_{|M|-1})^T \cdots (M^{-1}_0)^T)$
  - (c)  $= \det(D^T)$
  - (d)  $= \det(D)$
  - (e)  $= \det(M^{-1}_0 \cdots M^{-1}_{|M|-1} D N^{-1}_0 \cdots N^{-1}_{|N|-1})$

$$(f) = \det(A).$$

### Procedure V:42(4.30)

#### Objective

Choose a  $m \times n$  matrix,  $A$ , and an integer  $0 \leq k \leq \min(m, n)$ . The objective of the following instructions is to show that  $C_k(A)^T = C_k(A^T)$ .

#### Implementation

1. For each row label  $I$  of  $C_k(A^T)$ , do the following:
  - (a) For each column label  $J$  of  $C_k(A^T)$ , do the following:
    - i. Using **procedure V:41**, verify that  $(C_k(A^T))_{\underline{I}, \underline{J}} = \det((A^T)_{I, J}) = \det(A_{J, I}) = (C_k(A))_{\underline{J}, \underline{I}}$ .
2. Therefore verify that  $(C_k(A))^T = (C_k(A^T))$ .

# Chapter 17

## Polynomials and Normal Forms

### Procedure V:43(4.31)

#### Objective

Choose a  $m \times n$  rational matrix,  $A$ , and a  $m \times p$  rational matrix,  $B$ . Execute **procedure V:37** on  $A$  and let  $\langle M, D, , N \rangle$  receive the result. If the indices of the rows of  $D$  that are entirely zero are also the indices of the rows of  $M_*B$  that are entirely zero, then the objective of the following instructions is to construct a  $n \times p$  rational matrix  $E$  such that  $AE = B$ .

#### Implementation

1. Verify that  $A = M^{-1}_*DN^{-1}_*$ .
2. Verify that  $M^{-1}_*$ ,  $D$ , and  $N^{-1}_*$  are rational matrices.
3. Let  $C$  be an  $n \times p$  matrix with its  $i^{th}$  row given as follows:
  - (a) If  $D_{i,i} \neq 0$ , then do the following:
    - i. Let row  $i$  be row  $i$  of  $M_*B$  divided by  $D_{i,i}$ .
  - (b) Otherwise, do the following:
    - i. **Choose  $p$  rational numbers to fill up the row.**
4. Verify that  $DC = M_*B$ .
5. Let  $E$  be  $N_*C$ .
6. **Therefore using **procedure V:17**, verify that  $AE = M^{-1}_*DN^{-1}_*E =$**

$$\begin{aligned} M^{-1}_*DN^{-1}_*N_*C &= M^{-1}_*D1_nC = \\ M^{-1}_*DC &= M^{-1}_*M_*B = 1_mB = B. \end{aligned}$$

7. **Yield the tuple  $\langle E \rangle$ .**

#### Declaration V:21

The notation  $A \setminus B$  will be used to refer to the result yielded by executing **procedure V:43** on  $\langle A, B \rangle$ .

### Procedure V:44(4.32)

#### Objective

Choose a  $m \times n$  rational matrix,  $A$ , and a  $p \times n$  rational matrix,  $B$ . Execute **procedure V:37** on  $A$  and let  $\langle M, D, , N \rangle$  receive the result. If the indices of the columns of  $D$  that are entirely zero are also the indices of the columns of  $BN_*$  that are entirely zero, then the objective of the following instructions is to construct a  $p \times m$  rational matrix  $E$  such that  $EA = B$ .

#### Implementation

Instructions are analogous to those of **procedure V:43**.

#### Declaration V:22

The notation  $A/B$  will be used to refer to the result yielded by executing **procedure V:44** on  $\langle A, B \rangle$ .

### Procedure V:45(4.33)

#### Objective

Choose a  $m \times n$  rational matrix,  $A$ , a  $n \times p$  rational matrix,  $E$ , and a  $m \times p$  rational matrix,  $B$  such that  $AE = B$ . Execute [procedure V:37](#) on  $A$  and let  $\langle M, D, , N \rangle$  receive the result. If the indices of the rows of  $D$  that are entirely zero are not also the indices of the rows of  $M_*B$  that are entirely zero, then the objective of the following instructions is to show that  $0 \neq 0$ .

#### Implementation

1. Verify that  $M^{-1}_*DN^{-1}_*E = AE = B$ .
2. Therefore verify that  $DN^{-1}_*E = M_*B$ .
3. Let  $i$  be an integer such that  $D_{i,*}$  is zero and yet  $(M_*B)_{i,*}$  is not zero.
4. Verify that  $D_{i,*} = D_{i,*}N^{-1}_*E = (DN^{-1}_*E)_{i,*} = (M_*B)_{i,*}$ .
5. Let  $j$  be an integer such that  $(M_*B)_{i,j} \neq 0$ .
6. Now verify that  $0 = D_{i,j} = (M_*B)_{i,j} \neq 0$ .

### Procedure V:46(4.34)

#### Objective

Choose a  $p \times m$  rational matrix,  $E$ , a  $m \times n$  rational matrix,  $A$ , and a  $p \times n$  rational matrix,  $B$  such that  $EA = B$ . Execute [procedure V:37](#) on  $A$  and let  $\langle M, D, , N \rangle$  receive the result. If the indices of the columns of  $D$  that are entirely zero are not also the indices of the columns of  $BN_*$  that are entirely zero, then the objective of the following instructions is to show that  $0 \neq 0$ .

#### Implementation

Instructions are analogous to those of [procedure V:45](#).

### Procedure V:47(4.35)

#### Objective

Choose two  $m \times m$  rational matrices,  $A$  and  $B$ , such that  $AB = 1_m$ . The objective of the following instructions is to show that either  $0 = 1$  or  $BA = 1_m$ .

#### Implementation

1. Execute [procedure V:37](#) on  $B$  and let  $\langle M, D, , N \rangle$  receive the result.
2. Verify that  $B = M^{-1}_*DN^{-1}_*$ .
3. If  $D$  has a zero on its diagonal, then do the following:
  - (a) Using [procedure V:39](#), verify that  $\det(1_m) = \det(AB) = \det(A)\det(B) = \det(A)\det(D) = \det(A) * 0 = 0$ .
  - (b) Also verify that  $\det(1_m) = 1^m = 1$ .
  - (c) Therefore verify that  $0 = 1$ .
  - (d) **Abort procedure.**
4. Otherwise do the following:
  - (a) Verify that  $D$  does not have a zero on its diagonal.
  - (b) Verify that  $B \setminus 1_m = 1_m(B \setminus 1_m) = AB(B \setminus 1_m) = A(B(B \setminus 1_m)) = A1_m = A$ .
  - (c) **Therefore verify that  $BA = B(B \setminus 1_m) = 1_m$ .**

### Procedure V:48(4.36)

#### Objective

Choose an  $m \times m$  matrix,  $M$ , and an  $m \times m$  rational matrix,  $B$ . The objective of the following instructions is to construct a  $m \times m$  matrix,  $Q$ , and a  $m \times m$  rational matrix,  $R$ , such that  $M = (\lambda 1_m - B)Q + R$ .

#### Implementation

1. Let  $M_0\lambda^b + M_1\lambda^{b-1} + \dots + M_b\lambda^0 = M$ , where the  $M_i$  are  $m \times m$  rational matrices.
2. Now let  $R = B^bM_0 + B^{b-1}M_1 + \dots + B^0M_b$ .

3. Let  $Q = \sum_k^{[1:b]} (\lambda^{k-1} 1_m B^0 + \lambda^{k-2} 1_m B^1 + \dots + \lambda^0 1_m B^{k-1}) M_k$ .
4. Verify that  $M - R = (\lambda 1_m - B) \sum_k^{[1:b]} (\lambda^{k-1} 1_m B^0 + \lambda^{k-2} 1_m B^1 + \dots + \lambda^0 1_m B^{k-1}) M_k = (\lambda 1_m - B) Q$ .
5. **Verify that**  $M = (\lambda 1_m - B) Q + R$ .
6. **Yield the tuple**  $\langle Q, R \rangle$ .

### Procedure V:49(4.37)

#### Objective

Choose an  $m \times m$  matrix,  $M$ , and an  $m \times m$  rational matrix,  $B$ . The objective of the following instructions is to construct a  $m \times m$  matrix,  $Q$ , and a  $m \times m$  rational matrix,  $R$ , such that  $M = Q(\lambda 1_m - B) + R$ .

#### Implementation

The instructions are analogous to those of [procedure V:48](#).

### Procedure V:50(4.38)

#### Objective

Choose two  $m \times m$  rational matrices,  $B, A$ , and two lists of  $m \times m$  tilts such that  $\lambda 1_m - B = M(\lambda 1_m - A)N$ . The objective of the following instructions is to either show that  $0 = 1$  or to construct  $m \times m$  rational matrices  $R_1$  and  $R_3$  such that  $1_m = R_1 R_3$  and  $B = R_1 A R_3$ .

#### Implementation

1. Verify that  $(\lambda 1_m - B)N^{-1} = M(\lambda 1_m - A)NN^{-1} = M(\lambda 1_m - A)1_m = M(\lambda 1_m - A)$ .
2. Execute [procedure V:49](#) on  $\langle M, B \rangle$  and let  $\langle Q_1, R_1 \rangle$  receive.
3. Verify that  $M = (\lambda 1_m - B)Q_1 + R_1$ .
4. Execute [procedure V:49](#) on  $\langle N^{-1}, A \rangle$  and let  $\langle Q_2, R_2 \rangle$  receive.
5. Verify that  $N^{-1} = Q_2(\lambda 1_m - A) + R_2$ .

6. By substituting  $M$  and  $N^{-1}$  into (2), verify that  $(\lambda 1_m - B)(Q_2(\lambda 1_m - A) + R_2) = ((\lambda 1_m - B)Q_1 + R_1)(\lambda 1_m - A)$ .
7. By rearranging both sides, verify that  $(\lambda 1_m - B)(Q_2 - Q_1)(\lambda 1_m - A) = R_1(\lambda 1_m - A) - (\lambda 1_m - B)R_2$ .
8. By equating the coefficients of different powers of  $\lambda$  both sides, verify that  $Q_2 - Q_1 = 0_{m \times m}$ .
9. Verify that  $R_1(\lambda 1_m - A) - (\lambda 1_m - B)R_2 = (\lambda 1_m - B)(Q_2 - Q_1)(\lambda 1_m - A) = (\lambda 1_m - B)0_{m \times m}(\lambda 1_m - A) = 0_{m \times m}$ .
10. Therefore by adding  $(\lambda 1_m - B)R_2$  to both sides, verify that  $\lambda R_1 - R_1 A = R_1(\lambda 1_m - A) = (\lambda 1_m - B)R_2 = \lambda R_2 - B R_2$ .
11. By equating the coefficients of  $\lambda$  on both sides, verify that  $R_1 = R_2$ .
12. Therefore verify that  $R_1 A = B R_1$ .
13. Execute [procedure V:49](#) on  $\langle M^{-1}, A \rangle$  and let  $\langle Q_3, R_3 \rangle$  receive.
14. Verify that  $M^{-1} = (\lambda 1_m - A)Q_3 + R_3$ .
15. Verify that  $1_m = M M^{-1} = ((\lambda 1_m - B)Q_1 + R_1)M^{-1} = (\lambda 1_m - B)Q_1 M^{-1} + R_1 M^{-1} = (\lambda 1_m - B)Q_1 M^{-1} + R_1(\lambda I - A)Q_3 + R_1 R_3 = (\lambda 1_m - B)Q_1 M^{-1} + (\lambda I - B)R_1 Q_3 + R_1 R_3 = (\lambda 1_m - B)(Q_1 M^{-1} + R_1 Q_3) + R_1 R_3$ .
16. By equating the powers of  $\lambda$  on both sides, verify that  $Q_1 M^{-1} + R_1 Q_3 = 0$ .
17. By substituting zero for  $Q_1 M^{-1} + R_1 Q_3$ , **verify that**  $1_m = (\lambda 1_m - B)0_{m \times m} + R_1 R_3 = R_1 R_3$ .
18. **Therefore using [procedure V:47](#), verify that**  $R_3 R_1 = 1_m$ .
19. **Also, verify that**  $B = B 1_m = B R_1 R_3 = R_1 A R_3$ .
20. **Yield the pair**  $(R_1, R_3)$ .

### Procedure V:51(4.39)

#### Objective

Choose a  $m \times n$  matrix,  $A$ . Choose two integers  $0 \leq i, j < m$  such that  $i \neq j$ . The objective of the following instructions is to negate row  $i$  and swap it with row  $j$  using only elementary row operations.

### Implementation

1. Let  $A$  be our working matrix.
2. Subtract row  $j$  from row  $i$ .
3. Add row  $i$  to row  $j$ .
4. Subtract row  $j$  from row  $i$ .
5. **Verify that the  $i^{th}$  row has been negated and swapped with the  $j^{th}$  row.**

### Procedure V:52(4.40)

#### Objective

Choose a  $m \times n$  matrix,  $A$ . Choose two integers  $0 \leq i, j < n$  such that  $i \neq j$ . The objective of the following instructions is to negate column  $i$  and swap it with row  $j$  using only elementary column operations.

#### Implementation

The instructions are analogous to those of [procedure V:51](#).

### Procedure V:53(4.41)

#### Objective

Choose an  $m \times n$  diagonal matrix,  $A$ . Choose two integers  $0 \leq i, j < \min(m, n)$  such that  $i \neq j$ . The objective of the following instructions is to swap  $B_{i,i}$  and  $B_{j,j}$  using only elementary row and column operations.

#### Implementation

1. Let  $A$  be our working matrix.
2. Use [procedure V:52](#) to negate the  $i^{th}$  row and swap it with the  $j^{th}$  row.
3. Use [procedure V:52](#) to negate the  $i^{th}$  column and swap it with the  $j^{th}$  column.
4. **Therefore, overall verify that  $B_{i,i}$  and  $B_{j,j}$  have been swapped.**

### Procedure V:54(4.42)

#### Objective

Choose an  $m \times n$  diagonal matrix,  $A$ . Choose two integers  $0 \leq i, j < \min(m, n)$  such that  $i \neq j$ . Choose a rational  $k \neq 0$ . The objective of the following instructions is to multiply  $B_{i,i}$  by  $k$  and  $B_{j,j}$  by  $\frac{1}{k}$  using only elementary row and column operations.

#### Implementation

1. Let  $A$  be our working matrix.
2. Add  $k$  times row  $i$  to row  $j$ .
3. Subtract  $\frac{1}{k}$  times row  $j$  from row  $i$ .
4. Add  $k$  times row  $i$  to row  $j$ .
5. Verify that the  $i^{th}$  row has been scaled by  $k$ , the  $j^{th}$  row by  $-\frac{1}{k}$ , and that both these rows are swapped.
6. Use [procedure V:52](#) to negate the  $i^{th}$  row and swap it with the  $j^{th}$  row.
7. **Therefore, overall verify that  $B_{i,i}$  has been multiplied by  $k$ , and  $B_{j,j}$  by  $\frac{1}{k}$ .**

### Procedure V:55(4.43)

#### Objective

Choose a  $m \times m$  rational matrix,  $A$ . Execute [procedure V:22](#) on the polynomial matrix  $\lambda I - A$  and let  $\langle B \rangle$  be the result. The objective of the following instructions is to show that either none of the diagonal entries of  $B$  are equal to zero, or  $1 = 0$ .

#### Implementation

1. Verify that  $\det(\lambda I - A)$  is a monic polynomial of degree  $m$ .
2. Therefore using [procedure V:39](#), verify that  $\det(B) = \det(\lambda I - A)$ .
3. Therefore verify that  $\det(B)$  is a monic polynomial of degree  $m$ .
4. If any of the diagonal entries of  $B$  equal zero, then do the following:



- (a) Verify that  $\det(B) = B_{0,0}B_{1,1} \cdots B_{m-1,m-1} = 0$ .
  - (b) Therefore using (3) and (4a), verify that  $1 = 0$ .
  - (c) **Abort procedure.**
5. Otherwise do the following:
- (a) **Verify that none of the diagonal entries of  $B$  equal zero.**
  - ii. Verify that  $B_{i,i} \neq 0$ .
  - iii. Therefore verify that  $u_i \neq 0$ .
  - iv. **Therefore verify that**  $\deg(B_{i,i}) = \deg(u_i B_{i-1,i-1}) \geq \deg(B_{i-1,i-1}) > 0$ .
  - (e) **Yield the tuple  $\langle a \rangle$ .**

## Procedure V:56(4.44)

### Objective

Choose a positive integer  $m$  and an  $m \times m$  rational matrix,  $A$ . Execute **procedure V:37** on the polynomial matrix  $\lambda 1_m - A$  and let  $\langle B, v, \rangle$  be the result. The objective of the following instructions is to either show that  $0 < 0$  or to construct an integer  $a$  such that  $\sum_i^{[a:m]} \deg(B_{i,i}) = m$ ,  $\deg(B_{i,i}) > 0$  for  $i$  in  $[a : m]$ , and  $\deg(B_{i,i}) = 0$  for  $i$  in  $[0 : a]$ .

### Implementation

1. Execute **procedure V:55** on  $A$ .
2. If  $\deg(B_{i,i}) = 0$  for  $i$  in  $[0 : m]$ , then do the following:
  - (a) Verify that  $\det(\lambda 1_m - A) = \det(B) = B_{0,0}B_{1,1} \cdots B_{m-1,m-1}$ .
  - (b) **Therefore verify that**  $0 < m = \deg(\det(\lambda 1_m - A)) = \deg(B_{0,0}B_{1,1} \cdots B_{m-1,m-1}) = 0 + 0 + \cdots + 0 = 0$ .
  - (c) **Abort procedure.**
3. Otherwise do the following:
  - (a) Let  $0 \leq a < m$  be the least integer such that  $\deg(B_{a,a}) > 0$ .
  - (b) **Verify that**  $\deg(B_{i,i}) = 0$  for  $i$  in  $[0 : a]$ .
  - (c) **Verify that**  $\sum_i^{[a:m]} \deg(B_{i,i}) = \sum_i^{[0:m]} \deg(B_{i,i}) = \deg(B_{0,0}B_{1,1} \cdots B_{m-1,m-1}) = \deg(\det(B)) = \deg(\lambda 1_m - A) = m$ .
  - (d) For  $i$  in  $[a + 1 : m]$ , do the following:
    - i. Verify that  $B_{i,i} = u_i B_{i-1,i-1}$ .

### Declaration V:23

The notation  $(e_i)_{k \times 1}$  will be used to refer to the  $k \times 1$  rational matrix such that its  $i^{th}$  entry, 1, is the only non-zero entry.

### Declaration V:24

The notation  $\text{mat}_t(p)$  will be used as a shorthand for  $\sum_j^{[0:t]} p_j e_j$ .

### Declaration V:25

The notation  $\text{comp}(p)$ , where  $p \neq 0$  is a monic polynomial such that  $\deg(p) > 0$ , will be used as a shorthand for the  $\deg(p) \times \deg(p)$  rational matrix of the following constitution:

1. Its first  $\deg(p) - 1$  columns equal the last  $\deg(p) - 1$  columns of  $1_k$ .
2. Its last column is  $-\text{mat}_{\deg(p)}(p)$ .

## Procedure V:57(4.45)

### Objective

Choose a monic polynomial,  $p$  such that  $\deg(p) > 0$ . Let  $k = \deg(p)$ . Choose a  $k \times k$  matrix,  $D$ , such that  $D = \lambda 1_k - \text{comp}(p)$ . The objective of the following instructions is to transform  $D$  into  $\text{diag}(1, \dots, 1, p)$  by a sequence of elementary operations.

### Implementation

1. Let the matrix  $D$  be our working matrix.
2. For  $i$  in  $[k : 1]$ , add  $\lambda$  times row  $i$  to row  $i - 1$ .
3. Verify that  $D$ 's first  $k - 1$  columns are now the last  $k - 1$  columns of  $-1_k$ .

4. Verify that  $D$ 's last column is  $p$  followed by some other polynomials.
5. For  $i$  in  $[1 : k]$ , subtract  $D_{i,k-1}$  times column  $i - 1$  from column  $k - 1$ .
6. Verify that  $D$ 's last column is now  $p$  followed by zeros.
7. For  $i$  in  $[1 : k]$ , negate row  $i - 1$  and exchange it with row  $i$  using [procedure V:52](#).
8. **Therefore verify that**  $D = \text{diag}(1, \dots, 1, p)$ .

### Procedure V:58(4.46)

#### Objective

Choose a positive integer  $m$  and an  $m \times m$  rational matrix,  $A$ . Execute [procedure V:15](#) on the polynomial matrix  $\lambda 1_m - A$  and let  $\langle B, , \rangle$  receive the result. Execute [procedure V:56](#) on  $A$  and let  $\langle a \rangle$  receive the result. Let  $E_i = \text{comp}(\text{mon}(B_{a+i,a+i}))$  for  $i$  in  $[0 : m - a]$ . The objective of the following instructions is to first show that  $\text{cols}(\text{diag}(E)) = m$ , and second to apply a sequence of elementary operations on  $\lambda 1_m - \text{diag}(E)$  to obtain the matrix  $B$ .

#### Implementation

1. Verify that the diagonal of  $B$  comprises  $a$  rationals followed by  $B_{a,a}, B_{a+1,a+1}, \dots, B_{m-1,m-1}$ .
2. **Using [procedure V:57](#), verify that**  

$$\begin{aligned} \text{cols}(\text{diag}(E)) &= \sum_i^{[0:|E|]} \text{cols}(E_i) = \\ &= \sum_i^{[0:|E|]} \text{cols}(\text{comp}(\text{mon}(B_{a+i,a+i}))) = \\ &= \sum_i^{[0:|E|]} \text{deg}(\text{mon}(B_{a+i,a+i})) = \sum_i^{[0:m-a]} \text{deg}(B_{a+i,a+i}) = \\ &= \sum_i^{[a:m]} \text{deg}(B_{i,i}) = m. \end{aligned}$$
3. Let  $F = \lambda 1_m - \text{diag}(E)$ .
4. Now for  $i$  in  $[0 : |E|]$ :
  - (a) Let  $j = \sum_r^{[0:i]} \text{cols}(E_r)$ .
  - (b) Let  $k = j + \text{cols}(E_i)$ .
  - (c) Apply [procedure V:57](#) on the tuple  $\langle \text{mon}(B_{a+i,a+i}), F_{[j:k],[j:k]} \rangle$ .
5. Now verify that  $F$  is an  $m \times m$  diagonal rational matrix.

6. Also verify that the diagonal of  $F$  comprises  $\text{mon}(B_{a,a}), \text{mon}(B_{a+1,a+1}), \dots, \text{mon}(B_{m-1,m-1})$  and  $a$  1s.
7. Rearrange the diagonal of  $F$  so that  $\text{mon}(B_{i,i})$  is at the  $i^{\text{th}}$  position on the diagonal for  $i$  in  $[a : m]$  by doing pairwise swaps. In general, swap the  $i^{\text{th}}$  and  $j^{\text{th}}$  diagonal entries using [procedure V:53](#).
8. For  $i$  in  $[0 : m - 1]$ , do the following:
  - (a) Let  $k = \frac{(B_{i,i})_{\text{deg}(B_{i,i})}}{(F_{i,i})_{\text{deg}(F_{i,i})}}$ .
  - (b) Scale  $B_{i,i}$  by  $k$  and  $B_{i+1,i+1}$  by  $\frac{1}{k}$  using [procedure V:54](#).
  - (c) Now verify that  $F_{i,i} = B_{i,i}$ .
9. Now verify that  $\det(F)_m = \det(\lambda 1_m - \text{diag}(E))_m = 1 = \det(\lambda 1_m - A)_m = \det(B)_m$ .
10. Therefore verify that  $(F_{m,m})_{\text{deg}(F_{m,m})}$ 
  - (a)  $= \frac{\det(F)_m}{(\det(F_{[1:m],[1:m]}))_{m-\text{deg}(F_{m,m})}}$
  - (b)  $= \frac{\det(B)_m}{(\det(B_{[1:m],[1:m]}))_{m-\text{deg}(B_{m,m})}}$
  - (c)  $= (B_{m,m})_{\text{deg}(B_{m,m})}$ .
11. Therefore verify that  $F_{m,m} = B_{m,m}$ .
12. **Therefore verify that**  $F = B$ .

### Procedure V:59(4.47)

#### Objective

Choose a  $m \times m$  rational matrix,  $A$ . Execute [procedure V:56](#) on  $A$  and let  $\langle a \rangle$  receive the result. Let  $E_i = \text{comp}(\text{mon}(B_{a+i,a+i}))$  for  $i$  in  $[0 : m - a]$ . The objective of the following instructions is to either show that  $0 = 1$  or to construct  $m \times m$  rational matrices  $R, T$  such that  $A = R \text{diag}(E) T$ ,  $RT = 1_m$ , and  $TR = 1_m$ .

#### Implementation

1. Execute [procedure V:37](#) on the polynomial matrix  $\lambda 1_m - A$  and let  $\langle P, B, , Q \rangle$  be the result.
2. Verify that  $P_*(\lambda 1_m - A)Q_* = B$ .
3. Verify that  $\lambda 1_m - A = P^{-1} {}_* B Q^{-1} {}_*$ .

4. Let  $Z$  be a variant of **procedure V:37** where every occurrence of **procedure V:22** in its instructions is replaced with **procedure V:58**, and where every mention of  $v$  is ignored.
5. Execute procedure  $Z$  on the matrix  $\lambda 1_m - \text{diag}(E)$  and let  $\langle M, \cdot, N \rangle$  receive the result.
6. Verify that  $M_*(\lambda 1_m - \text{diag}(E))N_* = B$ .
7. Verify that  $\lambda 1_m - A = P^{-1}_* B Q^{-1}_* = P^{-1}_* M(\lambda 1_m - \text{diag}(E)) N Q^{-1}_*$ .
8. Execute **procedure V:50** on the matrices  $\langle A, P^{-1}M, \text{diag}(E), NQ^{-1} \rangle$ . Let the tuple  $\langle R, T \rangle$  be the result.
9. **Verify that**  $A = R \text{diag}(E) T$ .
10. **Verify that**  $RT = 1_m$ .
11. **Verify that**  $TR = 1_m$ .
12. **Yield the tuple**  $\langle R, E, T \rangle$ .

#### Procedure V:60(4.86)

##### Objective

Choose two polynomials  $a, b$  and an  $m \times m$  matrix  $C$  such that  $a = b$ . The objective of the following instructions is to show that  $\Lambda(a, C) = \Lambda(b, C)$ .

##### Implementation

Implementation is analogous to that of **procedure II:34**.

#### Procedure V:61(4.87)

##### Objective

Choose two polynomials  $a, b$  and an  $m \times n$  matrix  $C$ . The objective of the following instructions is to show that  $\Lambda(a + b, C) = \Lambda(a, C) + \Lambda(b, C)$ .

##### Implementation

Implementation is analogous to that of **procedure II:39**.

#### Procedure V:62(4.88)

##### Objective

Choose a polynomial  $a$  and an  $m \times m$  matrix  $B$ . The objective of the following instructions is to show that  $\Lambda(-a, B) = -\Lambda(a, B)$ .

##### Implementation

Implementation is analogous to that of **procedure II:45**.

#### Procedure V:63(4.89)

##### Objective

Choose two polynomials  $a, b$  and an  $m \times m$  matrix  $C$ . The objective of the following instructions is to show that  $\Lambda(ab, C) = \Lambda(a, C)\Lambda(b, C)$ .

##### Implementation

Implementation is analogous to that of **procedure II:48**.

#### Procedure V:64(4.48)

##### Objective

Choose a polynomial,  $r$ , and  $m \times m$  rational matrices,  $R, A, S$  such that  $SR = 1_m$ . The objective of the following instructions is to show that  $\Lambda(r, RAS) = R\Lambda(r, A)S$ .

##### Implementation

1. Verify that  $\Lambda(r, RAS)$ 
  - (a)  $= \sum_j^{[0:|r|]} r_j (RAS)^j$
  - (b)  $= \sum_j^{[0:|r|]} r_j R A^j S$
  - (c)  $= R (\sum_j^{[0:|r|]} r_j A^j) S$
  - (d)  $= R \Lambda(r, A) S$ .

## Procedure V:65(4.49)

### Objective

Choose a list of  $m \times m$  rational matrices,  $A$ , and a polynomial,  $r$ . The objective of the following instructions is to show that  $\Lambda(r, \text{diag}(A)) = \text{diag}(\Lambda(r, A))$ .

### Implementation

1. For  $i = 0$  up to  $i = t$ , by repeated applications of **procedure V:21**, verify that  $\text{diag}(A)^i$  evaluates to  $\text{diag}(A^i)$ .
2. Therefore verify that  $\Lambda(r, \text{diag}(A))$ 
  - (a)  $= \sum_j^{[0:|r|]} r_j \text{diag}(A)^j$
  - (b)  $= \sum_j^{[0:|r|]} r_j \text{diag}(A^j)$
  - (c)  $= \sum_j^{[0:|r|]} \text{diag}(r_j A^j)$
  - (d)  $= \text{diag}(\sum_j^{[0:|r|]} r_j A^j)$
  - (e)  $= \text{diag}(\Lambda(r, A))$ .

## Procedure V:66(4.50)

### Objective

Choose a  $m \times m$  rational matrix,  $A$ , and a polynomial,  $r$ . Execute **procedure V:59** on the matrix  $A$  and let the tuple  $\langle R_1, E, R_3 \rangle$  receive the result. The objective of the following instructions is to show that  $\Lambda(r, A) = R_1 \text{diag}(\Lambda(r, E))R_3$ .

### Implementation

1. Verify that  $R_3 R_1 = 1_m$ .
2. Using **procedure V:64**, verify that  $\Lambda(r, A) = \Lambda(r, R_1 \text{diag}(E)R_3) = R_1 \Lambda(r, \text{diag}(E))R_3$ .
3. Using **procedure V:65**, verify that  $\Lambda(r, \text{diag}(E)) = \text{diag}(\Lambda(r, E))$ .
4. **Therefore verify that**  $\Lambda(r, A) = R_1 \text{diag}(\Lambda(r, E))R_3$ .

## Procedure V:67(4.51)

### Objective

Choose a monic polynomial  $p \neq 0$  such that  $\deg(p) > 0$ . The objective of the following instructions is to show that  $\Lambda(p, \text{comp}(p)) = 0_{\deg(p) \times \deg(p)}$ .

### Implementation

1. Let  $G = \text{comp}(p)$ .
2. For  $i$  in  $[0 : \deg(p)]$ , verify that  $G^i e_0 = G^{i-1} e_1 = \dots = G^0 e_i = e_i$ .
3. Therefore, for  $i \in [0 : \deg(p)]$ , do the following:
  - (a) Using (1), verify that  $\Lambda(p, G)e_i$ 
    - i.  $= (\sum_j^{[0:|p|]} p_j G^j) e_i$
    - ii.  $= (\sum_j^{[0:|p|]} p_j G^j) G^i e_0$
    - iii.  $= G^i (G G^{\deg(p)-1} + \sum_j^{[0:\deg(p)]} p_j G^j) e_0$
    - iv.  $= G^i (G e_{\deg(p)-1} + \sum_j^{[0:\deg(p)]} p_j e_j)$
    - v.  $= G^i 0_{\deg(p) \times 1}$
    - vi.  $= 0_{\deg(p) \times 1}$ .
4. **Therefore verify that**  $\Lambda(p, \text{comp}(p)) = \Lambda(p, G) = 0_{\deg(p) \times \deg(p)}$ .

### Declaration V:26

The notation **last<sub>A</sub>**, where  $A$  is an  $m \times m$  rational matrix, will be used as a shorthand for the polynomial yielded by executing the following instructions:

1. Execute **procedure V:37** on the polynomial matrix  $\lambda 1_m - A$  and let the tuple  $\langle B, , \rangle$  receive the result.
2. Yield  $\langle B_{m-1, m-1} \rangle$ .

## Procedure V:68(4.52)

### Objective

Choose a  $m \times m$  rational matrix,  $A$ . The objective of the following instructions is to show that either  $1 = 0$  or **last<sub>A</sub>**  $\neq 0$ .

## Implementation

1. Execute **procedure V:55** on  $A$ .
2. **Therefore verify that**  $\text{last}_A \neq 0$ .

## Procedure V:69(4.53)

### Objective

Choose a  $m \times m$  rational matrix,  $A$ . The objective of the following instructions is to either show that  $0 < 0$  or to show that  $\Lambda(\text{last}_A, A) = 0_{m \times m}$ .

## Implementation

1. Execute **procedure V:37** on the matrix  $A$  and let the tuple  $\langle M, B, v, N \rangle$  receive the result.
2. Execute **procedure V:56** on  $A$  and let  $\langle a \rangle$  receive.
3. Execute **procedure V:59** on  $A$  and let  $\langle R, E, T \rangle$  receive.
4. For  $j$  in  $[0 : |E|]$ :
  - (a) Verify that  $E_j = \text{comp}(\text{mon}(B_{a+j, a+j}))$ .
  - (b) Verify that  $\text{last}_A = B_{m-1, m-1} = B_{a+j, a+j} \prod_r^{[a+j+1:m]} v_r$ .
  - (c) Let  $k = \deg(\text{mon}(B_{a+j, a+j}))$ .
  - (d) Therefore using **procedure V:67** verify that  $\Lambda(\text{last}_A, E_j) = \Lambda(B_{m-1, m-1}, E_j) = \Lambda(B_{a+j, a+j}, \text{comp}(\text{mon}(B_{a+j, a+j}))) \prod_r^{[a+j+1:m]} \Lambda(v_r, E_j) = 0_{k \times k} \prod_r^{[a+j+1:m]} \Lambda(v_r, E_j) = 0_{k \times k}$ .
5. **Therefore using procedure V:66 verify that**  $\Lambda(\text{last}_A, A) = R \text{diag}(\Lambda(\text{last}_A, E))T = R \text{diag}(\Lambda(B_{m-1, m-1}, E))T = R 0_{m \times m} T = 0_{m \times m}$ .

## Procedure V:70(4.54)

### Objective

Choose a monic polynomial  $p$  such that  $\deg(p) > 0$ . Choose a polynomial  $g \neq 0$  such that  $\deg(g) < \deg(p)$ . The objective of the following instructions is to show that  $\Lambda(g, \text{comp}(p)) \neq 0_{\deg(p) \times \deg(p)}$ .

## Implementation

1. Let  $G = \text{comp}(p)$ .
2. Therefore using **declaration V:25**, verify that  $\Lambda(g, G)e_0 = (\sum_j^{[0:\deg(g)+1]} g_j G^j)e_0 = \sum_j^{[0:\deg(g)+1]} g_j e_j \neq 0_{\deg(p) \times 1}$ .
3. **Therefore verify that**  $\Lambda(g, G) \neq 0_{\deg(p) \times \deg(p)}$ .

## Procedure V:71(4.55)

### Objective

Choose a polynomial  $g$  and a monic polynomial  $p$  such that  $\deg(p) = \deg(g) > 0$  and  $\Lambda(g, \text{comp}(p)) = 0_{\deg(g) \times \deg(g)}$ . The objective of the following instructions is to show that  $g = g_{\deg(g)}p$ .

## Implementation

1. Let  $G = \text{comp}(p)$ .
2. Using **declaration V:25**, verify that  $0_{\deg(g) \times 1} = \Lambda(g, G)e_0 = (\sum_j^{[0:|g|]} g_j G^j)e_0 = g_{\deg(g)} G e_{\deg(g)-1} + \sum_j^{[0:\deg(g)]} g_j e_j$ .
3. Therefore for  $i$  in  $[0 : \deg(g)]$ , do the following:
  - (a) Verify that  $0 = (g_{\deg(g)} G e_{\deg(g)-1} + \sum_j^{[0:\deg(g)]} g_j e_j)_{i,0}$ .
  - (b) Therefore using **declaration V:25**, verify that  $-g_{\deg(g)} p_i + g_i = 0$ .
  - (c) Therefore verify that  $g_i = g_{\deg(g)} p_i$ .
4. **Therefore verify that**  $g = g_{\deg(g)}p$ .

## Procedure V:72(4.56)

### Objective

Choose a  $m \times m$  rational matrix,  $A$ . Choose a polynomial  $p \neq 0$ , such that  $\Lambda(p, A) = 0_{m \times m}$ . The objective of the following instructions is to either show that  $0 \neq 0$  or to construct a polynomial  $f$  such that  $p = f \text{last}_A$ .

## Implementation

1. Let  $F$  be the  $1 \times 2$  matrix  $\langle \langle p, \text{last}_A \rangle \rangle$ .
2. Execute **procedure V:37** on  $F$  and let  $\langle M, D, , N \rangle$  receive the result.
3. Verify that  $D_{0,0} \neq 0$ .
4. Let  $g = D_{0,0}$ .
5. Verify that  $F = M^{-1} * D N^{-1} * = D N^{-1} *$ .
6. Verify that  $\text{last}_A = F_{0,1} = D_{0,0} N^{-1} *_{0,1} + D_{0,1} N^{-1} *_{1,1} = D_{0,0} N^{-1} *_{0,1} = g N^{-1} *_{0,1}$ .
7. Therefore verify that  $N^{-1} *_{0,1} \neq 0$ .
8. Let  $u = \deg(\text{last}_A)$ .
9. Now verify that  $u = \deg(\text{last}_A) = \deg(D_{0,0} N^{-1} *_{0,1}) \geq \deg(D_{0,0}) = \deg(g)$ .
10. Verify that  $D = M * F N * = F N *$ .
11. Therefore verify that  $g = D_{0,0} = N *_{0,0} p + N *_{1,0} \text{last}_A$ .
12. Therefore using **procedure V:67**, verify that  $\Lambda(g, A) = \Lambda(N *_{0,0}, A) \Lambda(p, A) + \Lambda(N *_{1,0}, A) \Lambda(\text{last}_A, A) = \Lambda(N *_{0,0}, A) 0_{m \times m} + \Lambda(N *_{1,0}, A) 0_{m \times m} = 0_{m \times m}$ .
13. Execute **procedure V:59** on the matrix  $A$  and let the tuple  $\langle R_1, E, R_3 \rangle$  receive the result.
14. Using **procedure V:66**, and **procedure V:59**, verify that  $\text{diag}(\Lambda(g, E)) = 1_m \text{diag}(\Lambda(g, E)) 1_m = R_3 R_1 \text{diag}(\Lambda(g, E)) R_3 R_1 = R_3 \Lambda(g, A) R_1 = R_3 0_{m \times m} R_1 = 0_{m \times m}$ .
15. Let  $G = \text{comp}(\text{mon}(\text{last}_A))$ .
16. Verify that  $\Lambda(g, G) = \Lambda(g, E_{|E|-1}) = \text{diag}(\Lambda(g, E))_{[m-u:m], [m-u:m]} = 0_{u \times u}$ .
17. If  $\deg(g) < u$ , then:
  - (a) Using **procedure V:70**, verify that  $\Lambda(g, G) \neq 0_{u \times u}$ .
  - (b) **Therefore using (16), verify that**  $0_{u \times u} = \Lambda(g, G) \neq 0_{u \times u}$ .
  - (c) **Abort procedure.**
18. Otherwise, do the following:
  - (a) Verify that  $\deg(g) = u$ .
  - (b) Using **procedure V:71**, verify that  $g = g_{\deg(g)} \text{last}_A$ .

$$(c) \text{ Therefore verify that } p = F_{0,0} = D_{0,0} N^{-1} *_{0,0} + D_{0,1} N^{-1} *_{1,0} = N^{-1} *_{0,0} g + N^{-1} *_{1,0} * 0 = N^{-1} *_{0,0} g = N^{-1} *_{0,0} g_{\deg(g)} \text{last}_A.$$

$$(d) \text{ Yield the tuple } \langle N^{-1} *_{0,0} g_{\deg(g)} \rangle.$$

## Procedure V:73(4.57)

### Objective

Choose an  $m \times n$  rational matrix,  $A$ , and an  $n \times m$  rational matrix,  $B$ , such that  $AB = 1_m$ . The objective of the following instructions is to show that either  $0 = 1$  or every column of  $B$  is non-zero.

### Implementation

1. If any column  $i$  of  $B$ ,  $Be_i$ , is equal to zero, then:
  - (a) Verify that  $0_{n \times 1} = A 0_{n \times 1} = A(Be_i) = (AB)e_i = 1_m e_i = e_i$ .
  - (b) **Therefore verify that  $0=1$ .**
  - (c) **Abort procedure.**

## Procedure V:74(4.58)

### Objective

Choose a  $m \times m$  rational matrix,  $A$ . Choose a polynomial  $p$  such that  $p \neq 0$ ,  $\Lambda(p, A) = 0$ , and  $\deg(p) < \deg(\text{last}_A)$ . The objective of the following instructions is to show that  $0 < 0$ .

### Implementation

1. Execute **procedure V:72** on  $A$  and  $p$  and let  $f$  receive.
2. Now verify that  $p = f \text{last}_A$ .
3. Now using the precondition and (2), verify that  $f \neq 0$  and  $\text{last}_A \neq 0$ .
4. **Therefore using the precondition, (2), and (3), verify that**  $\deg(\text{last}_A) > \deg(p) = \deg(f \text{last}_A) \geq \deg(\text{last}_A)$ .
5. **Abort procedure.**

### Declaration V:27

The notation  $\text{pows}(A)$ , where  $A$  is a  $m \times m$  rational matrix, will be used as a shorthand for the result yielded by executing the following instructions:

1. Let  $t = \deg(\text{last}_A)$ .
2. Make an  $m^2 \times t$  matrix,  $B$ , whose  $i^{\text{th}}$  column is the sequential concatenation of the columns of  $A^i$ .
3. **Yield**  $\langle B \rangle$ .

### Procedure V:75(4.59)

#### Objective

Choose a  $m \times m$  rational matrix,  $A$ . Execute **procedure V:37** on  $\text{pows}(A)$  and let the tuple  $\langle M, D, , N \rangle$  receive the result. Let  $t = \text{cols}(\text{pows}(A))$ . The objective of the following instructions is to show that either  $0 < 0$  or to show that  $C_t(D) = C_t(D)_{0,0}e_0 \neq 0$ .

#### Implementation

1. Execute **procedure V:37** on  $\text{pows}(A)$  and let the tuple  $\langle M, D, , N \rangle$  receive the result.
2. Verify that  $M_* \text{pows}(A)N_* = D$ .
3. Using **procedure V:17**, verify that  $M^{-1}_* M_* \text{pows}(A)N_* = 1_{m^2} \text{pows}(A)N_* = \text{pows}(A)N_* = M^{-1}_* D$ .
4. If  $C_t(D)_{0,0} = 0$ , then:
  - (a) Verify that for some  $0 \leq i < t$ ,  $D_{i,i} = 0$ .
  - (b) Therefore verify that  $De_i = 0_{m^2 \times 1}$ .
  - (c) Therefore verify that  $\text{pows}(A)(Ne_i) = (\text{pows}(A)N)e_i = (M^{-1}D)e_i = M^{-1}(De_i) = 0_{m^2 \times 1}$ .
  - (d) Let  $p = N_{0,i}\lambda^0 + N_{1,i}\lambda^1 + \dots + N_{t-1,i}\lambda^{t-1}$ .
  - (e) Therefore verify that  $\Lambda(p, A) = 0_{m \times m}$ .
  - (f) Execute **procedure V:73** on  $N^{-1}_*$  and  $N_*$ .
  - (g) Therefore verify that  $p \neq 0$ .
  - (h) Execute **procedure V:74** on  $A$  and  $p$ .
  - (i) **Abort procedure.**

5. Otherwise, do the following:

- (a) Execute **procedure V:33** on  $\langle D, 1_t, t \rangle$  and let  $E$  receive.
- (b) Verify that  $C_t(D) = C_t(D1_t) = EC_t(1_t) = E * 1 = E$ .
- (c) Verify that  $E$  is a  $\binom{m^2}{t} \times \binom{t}{t}$  diagonal matrix.
- (d) Therefore verify that  $C_t(D)$  is a  $\binom{m^2}{t} \times 1$  diagonal matrix.
- (e) **Therefore verify that**  $C_t(D) = C_t(D)_{0,0}e_0 \neq 0$ .

### Procedure V:76(4.60)

#### Objective

Choose a  $m \times m$  rational matrix,  $A$ . Let  $t = \text{cols}(\text{pows}(A))$ . The objective of the following instructions is to show that either  $0 < 0$  or to show that  $C_t(\text{pows}(A)) \neq 0$ .

#### Implementation

1. Execute **procedure V:37** on  $\text{pows}(A)$  and let the tuple  $\langle M, D, , N \rangle$  receive the result.
2. Verify that  $\text{pows}(A) = M^{-1}_* DN^{-1}_*$ .
3. Execute **procedure V:73** on  $C_t(M_*)$ ,  $C_t(M^{-1}_*)$ .
4. Hence verify that all columns of  $C_t(M^{-1}_*)$  are non-zero.
5. Execute **procedure V:75** on  $A$ .
6. Verify that  $C_t(D) = C_t(D)_{0,0}e_0 \neq 0$ .
7. Therefore verify that  $C_t(D)_{0,0} \neq 0$ .
8. Execute **procedure V:73** on  $C_t(N_*)$ ,  $C_t(N^{-1}_*)$ .
9. Hence verify that  $C_t(N^{-1}_*) \neq 0$ .
10. **Verify that**  $C_t(\text{pows}(A)) = C_t(M^{-1}_* DN^{-1}_*) = C_t(M^{-1}_*)C_t(D)C_t(N^{-1}_*) = C_t(M^{-1}_*)C_t(D)_{0,0}e_0C_t(N^{-1}_*) = C_t(D)_{0,0}C_t(N^{-1}_*)C_t(M^{-1}_*)e_0 \neq 0_{\binom{m^2}{t} \times 1}$ .

### Declaration V:28

The notation  $\text{tr}(A)$ , where  $A$  is a square matrix, will be used as a shorthand for the sum of its diagonal entries.

### Procedure V:77(4.68)

#### Objective

Choose two  $m \times m$  matrices  $A, B$ . The objective of the following instructions is to show that  $\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B)$ .

#### Implementation

1. Verify that  $\text{tr}(A + B)$ 
  - (a)  $= \sum_r^{[0:m]} (A + B)_{r,r}$
  - (b)  $= \sum_r^{[0:m]} (A_r + B_r)_{r,r}$
  - (c)  $= \sum_r^{[0:m]} A_{r,r} + \sum_r^{[0:m]} B_{r,r}$
  - (d)  $= \text{tr}(A) + \text{tr}(B)$ .

### Procedure V:78(4.69)

#### Objective

Choose a polynomial  $b$  and an  $m \times m$  matrix  $A$ . The objective of the following instructions is to show that  $\text{tr}(bA) = b \text{tr}(A)$ .

#### Implementation

1. Verify that  $\text{tr}(bA)$ 
  - (a)  $= \text{tr}(b_{m \times m} A)$
  - (b)  $= \sum_r^{[0:m]} (b_{m \times m} A)_{r,r}$
  - (c)  $= \sum_r^{[0:m]} \sum_t^{[0:m]} (b_{m \times m})_{r,t} A_{t,r}$
  - (d)  $= \sum_r^{[0:m]} (b_{m \times m})_{r,r} A_{r,r}$
  - (e)  $= \sum_r^{[0:m]} b A_{r,r}$
  - (f)  $= b \sum_r^{[0:m]} A_{r,r}$
  - (g)  $= b \text{tr}(A)$ .

### Procedure V:79(4.70)

#### Objective

Choose an  $m \times n$  matrix  $A$  and an  $n \times m$  matrix  $B$ . The objective of the following instructions is to show that  $\text{tr}(AB) = \text{tr}(BA)$ .

#### Implementation

1. Verify that  $\text{tr}(AB)$ 
  - (a)  $= \sum_r^{[0:m]} (AB)_{r,r}$
  - (b)  $= \sum_r^{[0:m]} \sum_t^{[0:n]} A_{r,t} B_{t,r}$
  - (c)  $= \sum_t^{[0:n]} \sum_r^{[0:m]} B_{t,r} A_{r,t}$
  - (d)  $= \sum_t^{[0:n]} (BA)_{t,t}$
  - (e)  $= \text{tr}(BA)$ .

### Procedure V:80(4.71)

#### Objective

Choose an  $m \times n$  matrix  $A$  such that  $A \neq 0$ . The objective of the following instructions is to show that  $\text{tr}(A^T A) > 0$ .

#### Implementation

1. Verify that  $\text{tr}(A^T A)$ 
  - (a)  $= \sum_r^{[0:n]} (A^T A)_{r,r}$
  - (b)  $= \sum_r^{[0:n]} \sum_t^{[0:m]} (A^T)_{r,t} A_{t,r}$
  - (c)  $= \sum_r^{[0:n]} \sum_t^{[0:m]} A_{t,r} A_{t,r}$
  - (d)  $= \sum_r^{[0:n]} \sum_t^{[0:m]} (A_{t,r})^2$
  - (e)  $> 0$ .

### Declaration V:29

The phrase "symmetric matrix" will be used to refer to matrices  $A$  such that " $A^T = A$ ".



## Procedure V:81(4.61)

### Objective

Choose a symmetric  $m \times m$  rational matrix,  $A$ . Let  $t = \deg(\text{last}_A)$ . Choose two polynomials  $u, w$  such that  $\deg(u) < t$  and  $\deg(w) < t$ . The objective of the following instructions is to show that  $\text{tr}(\Lambda(uw, A)) = \text{mat}(u)^T \text{pows}(A)^T \text{pows}(A) \text{mat}_t(w)$ .

### Implementation

1. Verify that  $\text{tr}(\Lambda(uw, A))$ 
  - (a)  $= \text{tr}(\Lambda(u, A)\Lambda(w, A))$
  - (b)  $= \text{tr}((\sum_p^{[0:t]} u_p A^p)(\sum_q^{[0:t]} w_q A^q))$
  - (c)  $= \text{tr}(\sum_p^{[0:t]} \sum_q^{[0:t]} u_p w_q A^p A^q)$
  - (d)  $= \sum_p^{[0:t]} \sum_q^{[0:t]} u_p w_q \text{tr}(A^p A^q)$
  - (e)  $= \sum_p^{[0:t]} \sum_q^{[0:t]} u_p w_q \sum_e^{[0:m]} \sum_f^{[0:m]} A^p_{e,f} \cdot A^q_{f,e}$
  - (f)  $= \sum_p^{[0:t]} \sum_q^{[0:t]} u_p w_q \sum_e^{[0:m]} \sum_f^{[0:m]} A^p_{f,e} \cdot A^q_{f,e}$
  - (g)  $= \sum_p^{[0:t]} \sum_q^{[0:t]} u_p w_q \sum_g^{[0:m^2]} \text{pows}(A)_{g,p} \text{pows}(A)_{g,q}$
  - (h)  $= \sum_p^{[0:t]} \sum_q^{[0:t]} u_p w_q (\text{pows}(A)^T \text{pows}(A))_{p,q}$
  - (i)  $= \sum_p^{[0:t]} u_p (\text{pows}(A)^T \text{pows}(A) \text{mat}_t(w))_p$
  - (j)  $= \text{mat}_t(u)^T \text{pows}(A)^T \text{pows}(A) \text{mat}_t(w)$

### Declaration V:30

The notation  $\text{sel}_A$ , where  $A$  is an  $m \times m$  rational matrix, will be used as a shorthand for the result yielded by executing the following instructions:

1. Using **procedure V:42**, **procedure V:76**, and **procedure V:80**, verify that  $C_t(\text{pows}(A)^T \text{pows}(A)) = C_t(\text{pows}(A)^T) C_t(\text{pows}(A)) = C_t(\text{pows}(A))^T C_t(\text{pows}(A)) = \text{tr}(C_t(\text{pows}(A))^T C_t(\text{pows}(A))) > 0$ .
2. Let  $t = \deg(\text{last}_A)$ .
3. Let  $H = (\text{pows}(A)^T \text{pows}(A)) \setminus e_{t-1}$ .
4. **Yield**  $\langle \frac{\sum_j^{[0:t]} H_{j,0} \lambda^j}{(\text{last}_A)_t} \rangle$ .

## Procedure V:82(4.62)

### Objective

Choose a symmetric  $m \times m$  rational matrix,  $A$ . Let  $t = \deg(\text{last}_A)$ . Choose a polynomial  $u$  such that  $\deg(u) < t$ . The objective of the following instructions is to show that  $\text{tr}(\Lambda(u \text{sel}_A, A)) = \frac{u_t - 1}{(\text{last}_A)_t}$ .

### Implementation

1. Using **procedure V:81**, verify that  $\text{tr}(\Lambda(u \text{sel}_A, A))$ 
  - (a)  $= \text{mat}(u)^T \text{pows}(A)^T \text{pows}(A) \text{mat}_t(\text{sel}_A)$
  - (b)  $= \frac{\text{mat}(u)^T \text{pows}(A)^T \text{pows}(A) ((\text{pows}(A)^T \text{pows}(A)) \setminus e_{t-1})}{(\text{last}_A)_t}$
  - (c)  $= \frac{\text{mat}(u)^T e_{t-1}}{(\text{last}_A)_t}$
  - (d)  $= \frac{\text{mat}(u)_{t-1,0}}{(\text{last}_A)_t}$
  - (e)  $= \frac{u_t - 1}{(\text{last}_A)_t}$ .

## Procedure V:83(4.63)

### Objective

Choose a symmetric  $m \times m$  rational matrix,  $A$ . The objective of the following instructions is to either show that  $0 \neq 0$  or construct polynomials  $u, v$  such that  $u \text{last}_A + v \text{sel}_A = 1$ .

### Implementation

1. Let  $t = \deg(\text{last}_A)$ .
2. Let  $G$  be the  $1 \times 2$  matrix  $\langle \langle \text{last}_A, \text{sel}_A \rangle \rangle$ .
3. Execute **procedure V:37** on  $G$  and let the tuple  $\langle M, D, , N \rangle$  receive.
4. Verify that  $G = M^{-1} * D N^{-1} *$ .
5. Verify that  $\text{last}_A \neq 0$ .
6. Therefore verify that  $D_{0,0} \neq 0$ .
7. If  $\deg(D_{0,0}) > 0$ , then do the following:
  - (a) Let  $b = N^{-1} *_{0,0}$ .
  - (b) Verify that  $\text{last}_A = b D_{0,0}$ .
  - (c) Therefore verify that  $b \neq 0$ .

- (d) Let  $z = \deg(b)$ .
  - (e) Verify that  $t = \deg(\text{last}_A) = \deg(bD_{0,0}) = \deg(b) + \deg(D_{0,0}) > \deg(b) = z$ .
  - (f) Let  $c = N^{-1}_{*0,1}$ .
  - (g) Verify that  $\text{sel}_A = cD_{0,0}$ .
  - (h) Let  $u = \lambda^{t-z-1}b$ .
  - (i) Execute **procedure V:82** on  $A$  and  $u$ .
  - (j) Hence verify that  $(\text{last}_A)_t \text{tr}(\Lambda(u \text{sel}_A, A)) = u_{t-1} = b_z \neq 0$ .
  - (k) Also verify that  $\text{tr}(\Lambda(u \text{sel}_A, A))$ 
    - i.  $= \text{tr}(\Lambda(\lambda^{t-z-1}bcD_{0,0}, A))$
    - ii.  $= \text{tr}(\Lambda(\lambda^{t-z-1}c \text{last}_A, A))$
    - iii.  $= \text{tr}(\Lambda(\lambda^{t-z-1}c, A)\Lambda(\text{last}_A, A))$
    - iv.  $= \text{tr}(\Lambda(\lambda^{t-z-1}c, A)0_{m \times m})$
    - v.  $= \text{tr}(0_{m \times m})$
    - vi.  $= 0$ .
  - (l) **Therefore verify that  $0 \neq 0$ .**
  - (m) **Abort procedure.**
8. Otherwise, do the following:
- (a) Verify that  $\deg(D_{0,0}) = 0$ .
  - (b) Let  $u = \frac{N_{0,0}}{D_{0,0}}$ .
  - (c) Let  $v = \frac{N_{1,0}}{D_{0,0}}$ .
  - (d) **Verify that  $u \text{last}_A + v \text{sel}_A = 1$ .**
  - (e) **Yield the tuple  $\langle u, v \rangle$ .**

## Procedure V:84(4.64)

### Objective

Choose a symmetric  $m \times m$  rational matrix  $A$ , where  $m > 0$ . Let  $t = \deg(\text{last}_A)$ . The objective of the following instructions is to either show that  $0 \neq 0$  or to construct lists of polynomials  $s, q$  such that

- 1. For  $i = 0$  to  $i = t$ ,  $\deg(s_i) = i$ .
- 2. For  $i = 0$  to  $i = t$ ,  $\text{sgn}((s_i)_i) = \text{sgn}((s_t)_t)$ .
- 3. For  $i = 1$  to  $i = t - 1$ ,  $s_{i-1} + s_{i+1} = q_i s_i$ .
- 4.  $s_t = \text{last}_A$ .

### Implementation

- 1. Let  $s_t = \text{last}_A$ .
- 2. Execute **procedure V:83** on  $A$  and let  $\langle u, s_{t+1} \rangle$  receive the result.
- 3. Hence verify that  $us_t + s_{t+1} \text{sel}_A = 1$ .
- 4. Let  $q_t = s_{t+1} \text{div } s_t$ .
- 5. Let  $s_{t-1} = s_{t+1} \bmod s_t$ .
- 6. Verify that  $s_{t+1} = q_t s_t + s_{t-1}$ , where  $\deg(s_{t-1}) < \deg(s_t) = t$ .
- 7. Therefore verify that  $us_t + (q_t s_t + s_{t-1}) \text{sel}_A = 1$ .
- 8. Therefore verify that  $\Lambda(s_{t-1} \text{sel}_A, A) = \Lambda(us_t + (q_t s_t + s_{t-1}) \text{sel}_A, A) = \Lambda(1, A) = 1_m$ .
- 9. Therefore using **procedure V:82**, verify that  $\frac{(s_{t-1})_{t-1}}{(s_t)_t} = \text{tr}(\Lambda(s_{t-1} \text{sel}_A, A)) = \text{tr}(1_m) = m > 0$ .
- 10. For  $i \in [t : 1]$ , do the following:
  - (a) Let  $q_i = (-s_{i+1}) \text{div}(-s_i)$ .
  - (b) Let  $s_{i-1} = (-s_{i+1}) \bmod (-s_i)$ .
  - (c) Verify that  $\deg(q_i) = 1$ .
  - (d) Verify that  $(q_i)_1 = \frac{(s_{i+1})_{i+1}}{(s_i)_i}$ .
  - (e) Also verify that  $-s_{i+1} = -q_i s_i + s_{i-1}$ .
  - (f) Therefore verify that  $q_i s_i = s_{i+1} + s_{i-1}$ .
  - (g) Therefore verify that  $q_i s_i - s_{i+1} = s_{i-1}$ .
  - (h) Execute **procedure II:76** on the tuple  $\langle s, q, i - 1 \rangle$  and let  $\langle p, j \rangle$  receive.
  - (i) Verify that  $s_{i-1} = ps_{t-1} + js_t$ .
  - (j) Verify that  $\deg(p) = t - 1 - (i - 1) = t - i$ .
  - (k) Verify that  $\deg(j) = t - 2 - (i - 1) = t - 1 - i$ .
  - (l) Therefore verify that  $\Lambda(s_{i-1}, A) = \Lambda(ps_{t-1} + js_t, A) = \Lambda(ps_{t-1}, A) + \Lambda(j, A)\Lambda(s_t, A) = \Lambda(ps_{t-1}, A) + \Lambda(j, A)0_{m \times m} = \Lambda(ps_{t-1}, A)$ .
- (m) If  $\Lambda(p, A) = 0$ , then do the following:
  - i. Execute **procedure V:74** on  $A$  and  $p$ .
  - ii. **Abort procedure.**

(n) Otherwise, if  $\Lambda(s_{i-1}, A) = 0_{m \times m}$ , then do the following:

- i. Verify that  $\Lambda(ps_{t-1} \text{sel}_A, A) = \Lambda(ps_{t-1}, A)\Lambda(\text{sel}_A, A) = \Lambda(s_{i-1}, A)\Lambda(\text{sel}_A, A) = 0_{m \times m}\Lambda(\text{sel}_A, A) = 0_{m \times m}$ .
- ii. Verify that  $\Lambda(ps_{t-1} \text{sel}_A, A) = \Lambda(p, A)\Lambda(s_{t-1} \text{sel}_A, A) = \Lambda(p, A)1_m = \Lambda(p, A) \neq 0_{m \times m}$ .

iii. Therefore verify that  $0 \neq 0$ .

iv. **Abort procedure.**

(o) Otherwise if  $\Lambda(s_{i-1} \text{sel}_A, A) = 0_{m \times m}$ , then do the following:

- i. Verify that  $\Lambda(s_{i-1} \text{sel}_A s_{t-1}, A) = \Lambda(s_{i-1} \text{sel}_A, A)\Lambda(s_{t-1}, A) = 0_{m \times m}\Lambda(s_{t-1}, A) = 0_{m \times m}$ .
- ii. Verify that  $\Lambda(s_{i-1} \text{sel}_A s_{t-1}, A) = \Lambda(s_{i-1}, A)\Lambda(\text{sel}_A s_{t-1}, A) = \Lambda(s_{i-1}, A)1_m = \Lambda(s_{i-1}, A) \neq 0_{m \times m}$ .

iii. Therefore verify that  $0_{m \times m} \neq 0_{m \times m}$ .

iv. **Abort procedure.**

(p) Otherwise, do the following:

- i. Verify that  $\deg(s_{i-1}) < i$ .
- ii. Verify that  $\Lambda(s_{i-1} \text{sel}_A, A) \neq 0_{m \times m}$ .
- iii. Execute the **subprocedure V:85:0** on the tuple  $(i-1, s_{i-1})$ .
- iv. Hence using **procedure V:80**, verify that  $\frac{(s_{i-1})_{i-1}}{(s_i)_i} = \text{tr}(\Lambda(s_{i-1}^2 \text{sel}_A^2, A)) = \text{tr}((\Lambda(s_{i-1} \text{sel}_A, A))^2) = \text{tr}((\Lambda(s_{i-1} \text{sel}_A, A))^T (\Lambda(s_{i-1} \text{sel}_A, A))) > 0$ .

v. **Therefore verify that**  $\text{sgn}((s_{i-1})_{i-1}) = \text{sgn}((s_i)_i)$ .

11. Yield the tuple  $\langle s_{[0:t+1]}, q_{[0:t]} \rangle$ .

### Subprocedure V:85:0

**Objective** Choose an integer  $0 \leq k \leq t$  such that polynomial  $s_k$  is defined. Choose a polynomial  $g$  such that  $\deg(g) \leq \min(k, t-1)$ . The objective of the following instructions is to show that  $\text{tr}(\Lambda(gs_k \text{sel}_A^2, A)) = \frac{g_k}{(s_{k+1})_{k+1}}$ .

### Implementation

1. If  $k = t$ , then verify that  $\text{tr}(\Lambda(gs_k \text{sel}_A^2, A))$

- (a)  $= \text{tr}(\Lambda(gs_t \text{sel}_A^2, A))$
- (b)  $= \text{tr}(\Lambda(g \text{sel}_A^2, A)\Lambda(s_t, A))$
- (c)  $= \text{tr}(\Lambda(g \text{sel}_A^2, A)0_{m \times m})$
- (d)  $= 0$
- (e)  $= \frac{g_k}{(s_{k+1})_{k+1}}$ .

2. Otherwise if  $k = t-1$ , then verify that  $\text{tr}(\Lambda(gs_k \text{sel}_A^2, A))$

- (a)  $= \text{tr}(\Lambda(gs_{t-1} \text{sel}_A^2, A))$
- (b)  $= \text{tr}(\Lambda(g \text{sel}_A, A)\Lambda(s_{t-1} \text{sel}_A, A))$
- (c)  $= \text{tr}(\Lambda(g \text{sel}_A, A)1_m)$
- (d)  $= \text{tr}(\Lambda(g \text{sel}_A, A))$
- (e)  $= \frac{g_k}{(s_{k+1})_{k+1}}$ .

3. Otherwise if  $k < t-1$ , then do the following:

- (a) Verify that  $\deg(gq_{k+1}) = k+1 \leq t-1$ .
- (b) Execute the **subprocedure V:85:0** on the tuple  $\langle k+1, gq_{k+1} \rangle$ .
- (c) Now verify that  $\text{tr}(\Lambda((gq_{k+1})s_{k+1} \text{sel}_A^2, A)) = \frac{(s_{k+2})_{k+2}}{(s_{k+1})_{k+1}} \frac{g_k}{(s_{k+2})_{k+2}} = \frac{g_k}{(s_{k+1})_{k+1}}$ .
- (d) Verify that  $\deg(g) \leq k \leq t-2$ .
- (e) Execute the **subprocedure V:85:0** on the tuple  $\langle k+2, g \rangle$ .
- (f) Now verify that  $\text{tr}(\Lambda(gs_{k+2} \text{sel}_A^2, A)) = \frac{g_{k+2}}{(s_{k+3})_{k+3}} = \frac{0}{(s_{k+3})_{k+3}} = 0$ .
- (g) Therefore verify that  $\text{tr}(\Lambda(gs_k \text{sel}_A^2, A))$ 
  - i.  $= \text{tr}(\Lambda(g(q_{k+1}s_{k+1} + s_{k+2}) \text{sel}_A^2, A))$
  - ii.  $= \text{tr}(\Lambda(gq_{k+1}s_{k+1} \text{sel}_A^2 + gs_{k+2} \text{sel}_A^2, A))$
  - iii.  $= \text{tr}(\Lambda(gq_{k+1}s_{k+1} \text{sel}_A^2, A) + \Lambda(gs_{k+2} \text{sel}_A^2, A))$
  - iv.  $= \text{tr}(\Lambda(gq_{k+1}s_{k+1} \text{sel}_A^2, A)) + \text{tr}(\Lambda(gs_{k+2} \text{sel}_A^2, A))$
  - v.  $= \frac{g_k}{(s_{k+1})_{k+1}} + 0$
  - vi.  $= \frac{g_k}{(s_{k+1})_{k+1}}$ .

## Procedure V:85(4.65)

### Objective

Choose a symmetric  $m \times m$  rational matrix,  $A$ . Let  $t = \deg(\text{last}_A)$ . The objective of the following instructions is to either show that  $0 < 0$  or to construct two lists of rational numbers  $c, d$  such that  $c_0 < d_0 \leq c_1 < d_1 \leq \dots \leq c_{t-1} < d_{t-1}$  and  $0 \neq \text{sgn}(\Lambda(\text{last}_A, c_i)) = -\text{sgn}(\Lambda(\text{last}_A, d_i))$  for  $i$  in  $[0 : t]$ .

### Implementation

1. Execute **procedure V:84** on the matrix  $A$  and let the tuple  $\langle s, q \rangle$  receive the result.
2. Execute **procedure II:75** supplying the tuple  $\langle s, q \rangle$ . Let the tuple  $\langle c, d \rangle$  receive the result.
3. **Verify that**  $c_0 < d_0 \leq c_1 < d_1 \leq \dots \leq c_{t-1} < d_{t-1}$ .
4. **Verify that**  $\text{sgn}(\Lambda(\text{last}_A, c_i)) = -\text{sgn}(\Lambda(\text{last}_A, d_i))$  **for**  $i$  **in**  $[0 : t]$ .
5. **Yield**  $\langle c, d \rangle$ .

## Procedure V:86(4.66)

### Objective

Choose a symmetric  $m \times m$  rational matrix,  $A$ . Let  $t = \deg(\text{last}_A)$ . Execute **procedure V:85** on  $A$  and let the tuple  $\langle c, d \rangle$  receive the result. Execute **procedure V:37** on  $A$  and let the tuple  $\langle, u, \rangle$  receive the result. The objective of the following instructions is to either show that  $1 = -1$  or to construct a list of non-negative integers  $k$  such that  $0 \neq \text{sgn}(\Lambda(u_{k_i}, c_i)) = -\text{sgn}(\Lambda(u_{k_i}, d_i))$  for  $i$  in  $[0 : t]$ .

### Implementation

1. Verify that  $\text{last}_A = u_0 u_1 \dots u_{m-1}$ .
2. For  $i$  in  $[0 : t]$ , do the following:
  - (a) Using the precondition, verify that  $0 \neq \text{sgn}(\Lambda(\text{last}_A, c_i)) = -\text{sgn}(\Lambda(\text{last}_A, d_i))$ .
  - (b) If  $0 \in \text{sgn}(\Lambda(u, c_i))$ , then do the following:
    - i. Verify that 0

- A.  $= \text{sgn}(\Lambda(u_0, c_i)) \text{sgn}(\Lambda(u_1, c_i)) \dots \text{sgn}(\Lambda(u_{m-1}, c_i))$
- B.  $= \text{sgn}(\Lambda(u_0, c_i) \Lambda(u_1, c_i) \dots \Lambda(u_{m-1}, c_i))$
- C.  $= \text{sgn}(\Lambda(u_0 u_1 \dots u_{m-1}, c_i))$
- D.  $= \text{sgn}(\Lambda(\text{last}_A, c_i))$
- E.  $\neq 0$ .

(c) If  $0 \in \text{sgn}(\Lambda(u, d_i))$ , then do the following:

i. Verify that 0

- A.  $= \text{sgn}(\Lambda(u_0, d_i)) \text{sgn}(\Lambda(u_1, d_i)) \dots \text{sgn}(\Lambda(u_{m-1}, d_i))$
- B.  $= \text{sgn}(\Lambda(u_0, d_i) \Lambda(u_1, d_i) \dots \Lambda(u_{m-1}, d_i))$
- C.  $= \text{sgn}(\Lambda(u_0 u_1 \dots u_{m-1}, d_i))$
- D.  $= \text{sgn}(\Lambda(\text{last}_A, d_i))$
- E.  $\neq 0$ .

(d) If  $\text{sgn}(\Lambda(u_j, c_i)) = \text{sgn}(\Lambda(u_j, d_i))$  for  $j \in [0 : m]$ , then do the following:

i. Verify that  $\text{sgn}(\Lambda(\text{last}_A, c_i))$

- A.  $= \text{sgn}(\Lambda(u_0 u_1 \dots u_{m-1}, c_i))$
- B.  $= \text{sgn}(\Lambda(u_0, c_i)) \text{sgn}(\Lambda(u_1, c_i)) \dots \text{sgn}(\Lambda(u_{m-1}, c_i))$
- C.  $= \text{sgn}(\Lambda(u_0, d_i)) \text{sgn}(\Lambda(u_1, d_i)) \dots \text{sgn}(\Lambda(u_{m-1}, d_i))$
- D.  $= \text{sgn}(\Lambda(u_0 u_1 \dots u_{m-1}, d_i))$
- E.  $= \text{sgn}(\Lambda(\text{last}_A, d_i))$ .

ii. **Therefore verify that**  $1 = -1$ .

iii. **Abort procedure.**

(e) Otherwise do the following:

- i. **Let**  $k_i$  **be the least integer such that**  $0 \neq \text{sgn}(\Lambda(u_{k_i}, c_i)) = -\text{sgn}(\Lambda(u_{k_i}, d_i))$ .

3. **Yield**  $\langle k \rangle$ .

## Procedure V:87(4.67)

### Objective

Choose a symmetric  $m \times m$  rational matrix,  $A$ . Execute [procedure V:37](#) on  $A$  and let the tuple  $\langle, u, \rangle$  receive the result. Execute [procedure II:69](#) on  $A$  and let  $k$  receive. Let  $t = \deg(\text{last}_A)$ . Let  $n_j = \sum_i^{[0:t]} [k_i = j]$  for  $j$  in  $[0 : m]$ . The objective of the following instructions is to either show that  $0 < 0$ , or to show that  $n_i = \deg(u_i)$  for  $i$  in  $[0 : m]$ .

### Implementation

1. Verify that  $\sum_j^{[0:m]} n_j = \sum_j^{[0:m]} \sum_i^{[0:t]} [k_i = j] = \sum_i^{[0:t]} 1 = t$ .
2. If for any  $i$  in  $[0 : m]$ ,  $n_i > \deg(u_i)$ , then do the following:
  - (a) Execute [procedure II:69](#) on the polynomial  $u_i$  along with  $\deg(u_i) + 1$  of the distinct pairs  $\langle c_l, d_l \rangle$  such that  $k_l = i$ .
  - (b) **Abort procedure.**
3. Otherwise if for any  $i$  in  $[0 : m]$ ,  $n_i < \deg(u_i)$ , then do the following:
  - (a) Verify that  $\sum_i^{[0:m]} n_j < \sum_i^{[0:m]} \deg(u_j) = t$ .
  - (b) Therefore using (1) and (a), verify that  $\sum_i^{[0:m]} n_j < \sum_i^{[0:m]} n_j$ .
  - (c) **Abort procedure.**
4. Otherwise, do the following:
  - (a) **For all  $i$  in  $[0 : m]$ , verify that  $n_i = \deg(u_i)$ .**

## Procedure V:88(4.72)

### Objective

Choose a symmetric  $m \times m$  rational matrix,  $A$ . Let  $t = \deg(\text{last}_A)$ . Execute [procedure V:86](#) on the matrix  $A$  and let the tuple  $\langle k \rangle$  receive the result. The objective of the following instructions is to either show that  $0 < 0$  or to show that  $\sum_i^{[0:t]} (m - k_i) = m$ .

### Implementation

1. Execute [procedure V:37](#) on the matrix  $A$  and let the tuple  $\langle, D, u, \rangle$ .
2. Using [procedure V:87](#), verify that  $\sum_i^{[0:t]} (m - k_i)$ 
  - (a)  $= \sum_i^{[0:t]} \sum_j^{[0:m]} [k_i \leq j]$
  - (b)  $= \sum_j^{[0:m]} \sum_i^{[0:t]} [k_i \leq j]$
  - (c)  $= \sum_j^{[0:m]} \sum_i^{[0:t]} [k_i \leq j] \sum_l^{[0:m]} [k_i = l]$
  - (d)  $= \sum_j^{[0:m]} \sum_l^{[0:m]} \sum_i^{[0:t]} [k_i \leq j] [k_i = l]$
  - (e)  $= \sum_j^{[0:m]} \sum_l^{[0:m]} \sum_i^{[0:t]} [l \leq j] [k_i = l]$
  - (f)  $= \sum_j^{[0:m]} \sum_l^{[0:m]} [l \leq j] \sum_i^{[0:t]} [k_i = l]$
  - (g)  $= \sum_j^{[0:m]} \sum_l^{[0:m]} [l \leq j] \deg u_l$
  - (h)  $= \sum_j^{[0:m]} \sum_l^{[0:j+1]} \deg u_l$
  - (i)  $= \sum_j^{[0:m]} \deg D_{j,j}$
  - (j)  $= m$

# Bibliography

- [1] Harold Edwards. *Linear Algebra*. Springer Science+Business Media, 1995.
- [2] Hugh L. Montgomery, Ivan Niven, Herbert S. Zuckerman. *An Introduction to the Theory of Numbers*. John Wiley & Sons, 1991.
- [3] Ludwig Wittgenstein. *Philosophical Grammar*. Edited by Rush Rhees. Translated by Anthony Kenny. Basil Blackwell, Oxford, 1974.