

REMARKS ON POINCARÉ DUALITY OF MATROIDS

RYOSUKE MUROOKA

ABSTRACT. Adiprasito, Huh, and Katz show the Kähler package for matroids using matroidal flips, which are purely combinatorial notions. Following their work, Amini and Piquerez give a new proof of the Kähler package, more generally for quasi-projective unimodular Bergman fans using Keel's lemma and the weak factorization theorem for smooth toric varieties. Later, in the matroid case, Hsiao shows that the weak factorization step can be replaced by an explicit sequence of toric blow-ups (i.e., star subdivisions of unimodular fans). This is an expository paper reviewing a simple proof of Poincaré duality for matroids using this sequence of star subdivisions and Keel's lemma from algebraic geometry.

1. INTRODUCTION

Let n, r be non-negative integers. In this paper, a *matroid* M refers to a loopless matroid of rank $r + 1$ on an $(n + 1)$ -set E . Write $\mathcal{L}(M)$ for the lattice of flats of M . Let $\bigoplus_{i \in E} \mathbf{Z}e_i$ be the free abelian group with E as its basis. Put $e_S := \sum_{i \in S} e_i$ for a subset $S \subset E$, $N_E := (\bigoplus_{i \in E} \mathbf{Z}e_i)/\mathbf{Z}e_E$, and $N_{E, \mathbf{R}} := N_E \otimes_{\mathbf{Z}} \mathbf{R}$. The *Bergman fan* Σ_M of a matroid M is the pure r -dimensional unimodular fan

$$\{ \text{Cone}(e_F : F \in \mathcal{F}) : \mathcal{F} \subset \mathcal{P}(M) \text{ is a flag} \}$$

in $N_{E, \mathbf{R}}$. Here, $\mathcal{P}(M) := \mathcal{L}(M) \setminus \{\emptyset, E\}$, and $\mathcal{F} \subset \mathcal{P}(M)$ is called a *flag* if it is linearly ordered with respect to the inclusion relation. Associated to Σ_M , we can consider the n -dimensional smooth quasi-projective complex toric variety X_{Σ_M} , which is possibly non-compact. Then, the *Chow ring* $A^*(M)$ of M is defined to be the Chow ring $A^*(X_{\Sigma_M})$ of X_{Σ_M} . Adiprasito, Huh, and Katz show Poincaré duality of M , which is part of the Kähler package, in the following sense.

Theorem 1 ([1, Theorem 6.19]). *For a matroid M of rank $r + 1$, $A^*(M)$ satisfies Poincaré duality of dimension r , i.e., it satisfies*

- 1) *there exists a group isomorphism $\deg : A^r(M) \xrightarrow{\sim} \mathbf{Z}$, called a degree map,*
- 2) *for every integer $q > r$, we have $A^q(M) = 0$, and*
- 3) *for every integer $q \leq r$, the map*

$$A^{r-q}(M) \rightarrow \text{Hom}(A^q(M), \mathbf{Z}); \quad x \mapsto (y \mapsto \deg(xy))$$

is a group isomorphism, i.e., the multiplication $A^q(M) \times A^{r-q}(M) \rightarrow A^r(M) \xrightarrow{\sim} \mathbf{Z}$ is a perfect pairing.

The key concepts of their proof are matroidal flips, which are purely combinatorial technical notions. Amini and Piquerez [2, Section 3] develop a new approach to the Kähler package, more generally for quasi-projective unimodular Bergman fans, using tropical modifications, Keel's lemma, and the weak factorization theorem for smooth toric varieties. See also [3, Section 1.9] for a later account of this argument. In the matroid case, the weak factorization step can be replaced by an explicit sequence of toric blow-ups. Motivated by [4], Hsiao, Karu, and Yang [6] calculate mixed volumes of matroids using the deletion of matroids. For their computation, they utilize the following important fact. For a non-coloop $a \in E$ of M and the deletion $M \setminus a$, the projection $\pi : N_E \rightarrow N_{E \setminus \{a\}}$ naturally induces a morphism of fans $\pi : \Sigma_M \rightarrow \Sigma_{M \setminus a}$, and this morphism factors as

$$(\clubsuit) \quad \Sigma_M = \Sigma_m \xrightarrow{\pi_m} \Sigma_{m-1} \xrightarrow{\pi_{m-1}} \cdots \xrightarrow{\pi_2} \Sigma_1 \xrightarrow{\pi_1} \Sigma_0 \xrightarrow{\pi_0} \Sigma_{M \setminus a},$$

where the morphism $\pi_0 : \Sigma_0 \rightarrow \Sigma_{M \setminus a}$ is induced by the projection $\pi : N_E \rightarrow N_{E \setminus \{a\}}$, $\pi_i : \Sigma_i \rightarrow \Sigma_{i-1}$ is induced by the identity map, and Σ_i is a star subdivision of Σ_{i-1} at a 2-dimensional unimodular cone for each $i = 1, \dots, m$. Using the factorization $(\clubsuit)$, Hsiao [5] gives a simple combinatorial proof that $A^*(M) \otimes_{\mathbf{Z}} \mathbf{R}$ satisfies the hard Lefschetz property, the Hodge-Riemann relations, and therefore Poincaré duality. In §3, we will give a proof of Theorem 1 in the language of algebraic geometry using $(\clubsuit)$ and Keel's lemma.

2. KEEL'S LEMMA

In this section, we recall an algebro-geometric lemma, which is known as Keel's lemma.

Theorem 2 (Keel's lemma, cf. [7, Appendix, Theorem 1]). *Let X be a smooth variety, $i : Z \hookrightarrow X$ a smooth closed subvariety of codimension z , $\pi : \tilde{X} \rightarrow X$ the blow-up of X along Z , and E the exceptional divisor, that is, we consider the pullback diagram*

$$\begin{array}{ccc} E & \hookrightarrow & \tilde{X} \\ \downarrow & \lrcorner & \downarrow \pi \\ Z & \xhookrightarrow{i} & X. \end{array}$$

If the pullback map $i^ : A^*(X) \rightarrow A^*(Z)$ is surjective, then the map $\pi^* : A^*(X)[T] \rightarrow A^*(\tilde{X})$; $T \mapsto -[E]$ induces an isomorphism*

$$\frac{A^*(X)[T]}{(P(T), T \cdot \text{Ker } i^*)} \xrightarrow{\sim} A^*(\tilde{X})$$

where $P(T) \in A^*(X)[T]$ is a monic polynomial of degree z . In particular, we have a decomposition

$$(\diamond) \quad A^q(\tilde{X}) = A^q(X) \oplus [E] \cdot A^{q-1}(Z) \oplus \dots \oplus [E]^{z-1} \cdot A^{q-z+1}(Z)$$

for all q , where we embed $\pi^* : A^*(X) \hookrightarrow A^*(\tilde{X})$, and for all positive integers p, q ,

$$A^{q-p}(Z) \xrightarrow[(i^*)^{-1}]{\sim} A^{q-p}(X)/\text{Ker } i^* \xhookrightarrow{[E]^p \cdot (-)} A^q(\tilde{X}).$$

Now we go back to our toric case. Let Σ be a unimodular fan in $N_{E, \mathbf{R}}$. For a z -dimensional cone $\sigma \in \Sigma$, let Σ_σ denote the star fan of Σ at σ , and let $\Sigma^*(\sigma)$ denote the star subdivision of Σ at σ . The orbit closure $V(\sigma)$ associated to σ is the smooth toric subvariety X_{Σ_σ} of codimension z , and $\pi : X_{\Sigma^*(\sigma)} \rightarrow X_\Sigma$ is the blow-up of X_Σ along $V(\sigma)$. Since the ring $A^*(X_\Sigma)$ is generated by the classes of torus-invariant divisors, the pullback homomorphism $A^*(X_\Sigma) \rightarrow A^*(X_{\Sigma_\sigma})$ is always surjective, so the assumption of Theorem 2 is satisfied if we substitute $X = X_\Sigma$, $Z = V(\sigma) = X_{\Sigma_\sigma}$, $\tilde{X} = X_{\Sigma^*(\sigma)}$. We also note that the q -th degree part $A^q(X_\Sigma)$ is a finitely generated abelian group.

Corollary 3. *We further assume that $A^*(X_\Sigma)$ and $A^*(X_{\Sigma_\sigma})$ satisfy Poincaré duality of dimension r and $r - z$ respectively. Then $A^*(X_{\Sigma^*(\sigma)})$ satisfies Poincaré duality of dimension r .*

Proof. The decomposition $(\diamond)$ tells us that $A^r(X_{\Sigma^*(\sigma)}) \cong \mathbf{Z}$ and $A^q(X_{\Sigma^*(\sigma)}) = 0$ for each $q > r$. By Poincaré duality of $A^*(X)$ and $A^*(Z)$, each summand of $(\diamond)$ is a finitely generated free abelian group. With an appropriately ordered basis chosen from each summand, the representation matrix of the bilinear pairing $A^q(X_{\Sigma^*(\sigma)}) \times A^{r-q}(X_{\Sigma^*(\sigma)}) \rightarrow \mathbf{Z}$ is a block lower triangular matrix

$$\begin{bmatrix} P_0 & O & \cdots & \cdots & O \\ O & P_1 & \ddots & \ddots & \vdots \\ \vdots & * & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & O \\ O & * & * & * & P_{z-1} \end{bmatrix}.$$

Poincaré duality of $A^*(X_\Sigma)$ and $A^*(X_{\Sigma_\sigma})$ imply that $\det P_q = \pm 1$ for all $q = 0, 1, \dots, z-1$, and therefore the above representation matrix has determinant ± 1 . ■

3. A PROOF OF POINCARÉ DUALITY BY STAR SUBDIVISIONS

In this section, we prove Theorem 1 by induction on n . Before starting the proof, we recall the definition of a coloop.

Definition 4. An element $a \in E$ is called a *coloop* of M if the subset $E \setminus \{a\}$ is a flat of M , or equivalently, $\dim \Sigma_{M \setminus a} < \dim \Sigma_M$.

If all elements of E are coloops of M , then M must be the Boolean matroid on E . The toric variety X_{Σ_M} is a smooth projective complex variety of dimension $r = n$. By the Jurkiewicz-Danilov theorem, the cycle map $A^*(X_{\Sigma_M}) \rightarrow H^{2*}(X_{\Sigma_M}, \mathbf{Z})$ is an isomorphism, and thus Poincaré duality of $A^*(M)$ follows in the usual sense. In particular, this establishes the statement for $n = 0$.

We assume that $n > 0$ and there exists a non-coloop $a \in E$ of M . If the singleton $\{a\}$ is not a flat of M , then the deletion map $d : \mathcal{L}(M) \rightarrow \mathcal{L}(M \setminus a); F \mapsto F \setminus \{a\}$ is bijective, and $\Sigma_M \subset (e_a - e_b)^\perp$ for some element b parallel to a . This implies that $X_{\Sigma_M} \cong X_{\Sigma_{M \setminus a}} \times \mathbf{G}_m$, and hence $A^*(M) \cong A^*(M \setminus a)$, where the right-hand side satisfies Poincaré duality of dimension r by the inductive hypothesis. Hence we may assume that $\{a\}$ is a flat of M . Now we construct the factorization ($\clubsuit$). Logically, the set $\mathcal{P}(M)$ can be partitioned into four disjoint subsets

$$\begin{aligned} \mathcal{P}(M) &= \{F : a \notin F, F \cup \{a\} \in \mathcal{P}(M)\} \sqcup \{F : a \notin F, F \cup \{a\} \notin \mathcal{P}(M)\} \\ &\quad \sqcup \{F : a \in F, F \setminus \{a\} \in \mathcal{P}(M)\} \sqcup \{F : a \in F, F \setminus \{a\} \notin \mathcal{P}(M)\} \\ &=: \mathcal{P}_1 \sqcup \mathcal{P}_2 \sqcup \mathcal{P}_3 \sqcup \mathcal{P}_4. \end{aligned}$$

Under the deletion map $d : \mathcal{P}(M) \rightarrow \mathcal{P}(M \setminus a)$, we have

$$\mathcal{P}(M \setminus a) = \mathcal{P}_1 \sqcup \mathcal{P}_2 \sqcup d(\mathcal{P}_4), \quad d(\mathcal{P}_3) \subset \mathcal{P}_1, \quad \mathcal{P}(M/a) = d(\mathcal{P}_3) \sqcup d(\mathcal{P}_4).$$

Consider the inverse map $d^{-1} : \mathcal{P}(M \setminus a) \rightarrow \mathcal{P}(M) \setminus \mathcal{P}_3$. We construct a fan Σ_0 in $N_{E, \mathbf{R}}$ by

$$\begin{aligned} \Sigma_0 &:= \{ \text{Cone}(e_{d^{-1}(F)} : F \in \mathcal{F}) : \mathcal{F} \subset \mathcal{P}(M \setminus a) \}_{\text{flag}} \\ &\quad \cup \{ \text{Cone}(e_{d^{-1}(F)} : F \in \mathcal{F}) + \text{Cone}(e_a) : \mathcal{F} \subset \mathcal{P}(M/a) \}_{\text{flag}} \end{aligned}$$

(the latter cones are intended to recover the rays associated to $\mathcal{P}_3$). Observing that Σ_0 has exactly one more ray than $\Sigma_{M \setminus a}$ (the ray is $\text{Cone}(e_a)$), one can show that the projection $\pi_0 : \Sigma_0 \rightarrow \Sigma_{M \setminus a}$ induces an isomorphism $\pi_0^* : A^*(M \setminus a) \xrightarrow{\sim} A^*(X_{\Sigma_0})$. We order the flats $d(\mathcal{P}_3) = \{F_1, F_2, \dots, F_m\}$ so that $F_i \subset F_j$ implies that $i \leq j$. Recall that, since $d(\mathcal{P}_3) \subset \mathcal{P}_1$, we have $d^{-1}(F_i) = F_i$ for all i , and hence $\text{Cone}(e_a, e_{F_i}) \in \Sigma_0$ for all i . Let Σ_1 be the star subdivision of Σ_0 at $\text{Cone}(e_{F_m}, e_a)$. Σ_1 has an extra ray $\text{Cone}(e_{F_m \cup \{a\}})$, and all new cones arising from this subdivision are of the form

$$\begin{aligned} &\text{Cone}(e_{d^{-1}(F)} : F \in \mathcal{F}) + \text{Cone}(e_{F_m \cup \{a\}}) \quad \text{where } \mathcal{F} \text{ is a flag of } \mathcal{P}(M/a), \text{ or} \\ &\text{Cone}(e_{d^{-1}(F)} : F \in \mathcal{F}) + \text{Cone}(e_a) + \text{Cone}(e_{F_m \cup \{a\}}) \quad \text{where } \mathcal{F} \text{ is a flag of } \mathcal{P}(M/a) \setminus \{F_m\}. \end{aligned}$$

The cones of the former form are already cones of Σ_M . By recursively constructing the star subdivision Σ_{i+1} of Σ_i at $\text{Cone}(e_{F_{m-i}}, e_a)$ for each $i = 1, \dots, m-1$, we obtain $\Sigma_m = \Sigma_M$ and the desired factorization

$$(\clubsuit) \quad \Sigma_M = \Sigma_m \xrightarrow{\pi_m} \Sigma_{m-1} \xrightarrow{\pi_{m-1}} \dots \xrightarrow{\pi_2} \Sigma_1 \xrightarrow{\pi_1} \Sigma_0 \xrightarrow{\pi_0} \Sigma_{M \setminus a}.$$

Finally, the star fan of Σ_i at $\text{Cone}(e_{F_{m-i}}, e_a)$ is isomorphic to the product $\Sigma_{[\emptyset, F_{m-i}]} \times \Sigma_{[F_{m-i} \cup \{a\}, E]}$, where $[\emptyset, F_{m-i}]$ denotes the restriction of M to F_{m-i} and $[F_{m-i} \cup \{a\}, E]$ denotes the contraction of M by $F_{m-i} \cup \{a\}$. This implies that its Chow ring is isomorphic to $A^*([\emptyset, F_{m-i}]) \otimes A^*([F_{m-i} \cup \{a\}, E])$, which satisfies Poincaré duality of dimension $(\text{rk}_M(F_{m-i}) - 1) + (\text{crk}_M(F_{m-i} \cup \{a\}) - 1) = r - 2$ by the inductive hypothesis. Applying Corollary 3 with $z = 2$, we conclude that $A^*(M)$ satisfies Poincaré duality of dimension r . $\blacksquare$

We give some remarks of the proof.

Remark 5. If we subdivide Σ_0 at $\text{Cone}(e_{F_1}, e_a)$, the resultant fan differs from Σ_M . For instance, when $F_1 \subset F_2$, $\text{Cone}(e_{F_1}, e_{F_2}, e_a)$ is divided into $\text{Cone}(e_{F_1 \cup \{a\}}, e_{F_2}, e_a)$ and $\text{Cone}(e_{F_1 \cup \{a\}}, e_{F_1}, e_{F_2})$. The former has a face $\text{Cone}(e_{F_1 \cup \{a\}}, e_{F_2})$, which remains in the final fan and is not a cone of Bergman fan.

Remark 6. In the language of tropical geometry, the fan Σ_0 is known as the tropical modification of the tropical fan $\Sigma_{M/a}$ along the tropical divisor $\Sigma_{M/a}$ with respect to a suitable piecewise linear function on $\Sigma_{M/a}$. Then the isomorphism $\pi^* : A^*(M/a) \rightarrow A^*(X_{\Sigma_0})$ follows from [3, Theorem 6.4].

ACKNOWLEDGEMENTS

The author is grateful to his advisor, Yusuke Nakamura, for steady guidance and numerous insightful comments. The author would like to express his sincere gratitude to Masahiko Yoshinaga for giving him helpful comments to improve the proof of Poincaré duality of matroids. The author would also like to thank Kosuke Mizuno and Kentaro Tanaka for informative comments.

REFERENCES

- [1] Karim Adiprasito, June Huh, and Eric Katz, *Hodge theory for combinatorial geometries*, Ann. Math. (2) **188** (2018), no. 2, 381–452. DOI 10.4007/annals.2018.188.2.1.
- [2] Omid Amini and Matthieu Piquerez, *Hodge theory for tropical varieties*, 2020. arXiv preprint, arXiv:2007.07826v1.
- [3] ———, *Hodge theory for tropical fans*, 2023. arXiv preprint, arXiv:2310.15367v1.
- [4] Tom Braden, June Huh, Jacob P. Matherne, Nicholas Proudfoot, and Botong Wang, *A semi-small decomposition of the Chow ring of a matroid*, Adv. Math. **409** (2022), Article 108646. DOI 10.1016/j.aim.2022.108646.
- [5] Andy Hsiao, *Hodge theory of matroids by star subdivisions* (2025), preprint, available at arXiv:2509.10909v1.
- [6] Andy Hsiao, Kalle Karu, and Jonathan Yang, *Mixed Volumes of Matroids*, Discrete Comput. Geom. (2026). DOI 10.1007/s00454-026-00832-y.
- [7] Sean Keel, *Intersection theory of moduli space of stable n -pointed curves of genus zero*, Trans. Amer. Math. Soc. **330** (1992), 545–574. DOI 10.1090/S0002-9947-1992-1034665-0.

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, NAGOYA, JAPAN

Email address: ryosuke.murooka.c1@math.nagoya-u.ac.jp