

Pytesimal: A Python package for modelling small planetary bodies

29 April 2021

Summary

Planetesimals were small bodies that existed in the protoplanetary disk, some of which coalesced to form the planets, while others are sampled by the meteorite record. Thermal processing recorded in meteorites can be linked to the thermal evolution of their parent planetesimals. This can inform us of the size and geometries of these parent bodies and how deep within them the meteorite material resided. Modelling the temperatures and cooling rates within planetesimals that existed 4.5 billion years ago is one of the key ways to understand the geological context of meteorites. The **Pytesimal** package focuses on the conductive cooling stage of planetesimal evolution, and provides a toolkit for modelling the temperature and cooling rate distribution inside meteorite parent bodies in 1D, with and without temperature-dependent material properties.

Statement of need

Pytesimal is a Python package for modelling the thermal evolution of planetesimals and other small planetary bodies. Meteorite parent body modelling is an active field in small-body planetary science. There are two broad categories of models: those focusing on the accretion and differentiation of planetesimals, and those investigating the later conductive cooling of parent bodies (Bryson et al. 2015; Elkins-Tanton, Weiss, and Zuber 2011; Haack, Rasmussen, and Warren 1990; Murphy Quinlan et al. 2021; Nichols et al. 2016; Sahijpal 2021). **Pytesimal** models the later conductive-cooling stage.

Pytesimal will enable groups to develop models of planetesimals and investigate the thermal history of meteorite parent bodies without having to rebuild the same basic architecture each time. **Pytesimal** provides a framework for modelling the conductive cooling of planetesimals, and is designed to be modular to allow future contributions and developments to be included. **Pytesimal** also includes plotting functionality to visualise the results of model runs, and a number of specialised tools designed to investigate pallasite meteorites specifically.

Method

The **Pytesimal** package focuses on the conductive cooling of differentiated planetesimals, with the ability to alter the model set-up to also investigate primitive bodies that have not segregated a core. The basic 1D set-up includes a conductively cooling discretised region which can include a low-diffusivity megaregolith layer, and an isothermal convecting core. The core can be removed to closer approximate primitive meteorite parent bodies, with a zero flux boundary condition applied across the centre to ensure symmetry (Figure 1).





Figure 1: TEMP PLACEHOLDER - TO BE PROPERLY DRAWN UP. Cartoon sketch of model set-up.

The 1D conductive cooling of the discretised region is controlled by the heat equation:

$$\frac{\partial T}{\partial t} \rho C = \frac{1}{r^2} \frac{\partial}{\partial r} \left(kr^2 \frac{\partial T}{\partial r} \right) = \overbrace{\frac{dk}{dT} \left(\frac{\partial T}{\partial r} \right)^2}^{\text{non-linear term}} + \underbrace{\frac{2k}{r} \frac{\partial T}{\partial r}}_{\text{geometric term}} + \overbrace{k \frac{\partial^2 T}{\partial r^2}}^{\text{linear term}}, \quad (1)$$

where T is the temperature, r is the radial value, t is time, and k , ρ and C are the conductivity, density and heat capacity respectively. **Pytesimal** provides the capability to use temperature-dependent conductivity, heat capacity and density, with functions suitable for an olivine mantle included. These T -dependent

material properties of olivine (??) are based on experimental results and mineral physics theory from Fei (2013), Robie, Hemingway, and Takei (1982), Su et al. (2018), Suzuki (1975), Xu et al. (2004), with more information in Murphy Quinlan et al. (2021).



The numerical_methods

module uses the explicit Forward-Time Central-Space (FTCS) scheme which is conditionally stable and must satisfy Von Neumann stability criteria in 1D: $\frac{\kappa \delta t}{\delta r^2} \leq \frac{1}{2}$, where κ is the thermal diffusivity of the material, δt is the timestep of the numerical scheme, and δr is the radial step (Crank and Nicolson 1947). `Pytesimal.numerical_methods` includes functions to calculate the diffusivity from k , ρ and C , and to check whether the chosen timestep will result in instabilities.

Boundary conditions for the top and bottom of the discretised region are passed into `numerical_methods.discretisation` as callable objects to allow for user-defined functions to be incorporated. Two different boundary conditions are currently provided, illustrated in ??: a fixed temperature condition which can be applied to either the top or bottom boundary of the discretised region, and a zero flux boundary condition that can be applied at the bottom boundary when the core is removed.

The core interacts with the mantle through heat extracted across the core-mantle boundary over one timestep in the form of power (P , in Watts). The heat extracted in one timestep (P_{CMB}) is calculated:

$$P_{\text{CMB}} = -A_c k_m \left. \frac{\partial T}{\partial r} \right|_{r=r_c} \quad (2)$$

where A_c is the core surface area, r_c is the core radius, and k_m is the thermal conductivity at the base of the mantle or discretised region. From this, the core boundary temperature is then updated by ΔT :

$$\Delta T = -\frac{P_{\text{CMB}}}{\rho_c C_c V_c} \delta t \quad (3)$$

where ρ_c and C_c are the density and heat capacity of the core, and V_c is the volume of the core. Once the core reaches its freezing temperature, the temperature is held constant. Latent heat is extracted until the total latent heat associated with core crystallisation has been removed. This simple eutectic core model ignores inner core formation and treats the liquid and solid fraction as identical, but is implemented in a way that allows the `IsothermalEutecticCore` object to be replaced with a more complex core model where applicable.

`Pytesimal` also contains the functionality to quickly plot results, which allows for both on-the-go data visualisation and for saved results to be loaded and plotted at a later time (Figure 2).

Examples of applications

An earlier version of `Pytesimal` has been used in a scientific publication to demonstrate that the inclusion of temperature dependent thermal properties in place of constant values can result in different interpretations of the meteorite record, with pallasite meteorites used as an example (Murphy Quinlan et al. 2021). Figure 2 reproduces the results of Murphy Quinlan et al. (2021).

Benefits of this package

1. `Pytesimal` only requires the commonly available Python packages `numpy` and `matplotlib`, with `Jupyter` useful for running the provided examples, but not essential.
2. Simple models can be set up and run in a single function call with an input parameter file, while more bespoke set ups only require a few extra lines of code.
3. `Pytesimal` is designed to be modular and extensible so that it can be applied to a wide range of modelling requirements, to speed up development of meteorite parent body models.
4. Quick and simple visualisation of the results can be achieved with a single function call, and can be modified easily to produce publication-quality figures.

Acknowledgements

MMQ was supported by the Leeds-York Natural Environment Research Council Doctoral Training Partnership (NE/L002574/1). The authors would like to thank Christopher J Davies for his comments on the paper.




Figure 2: Temperatures and cooling rates in a 250 km radius planetesimal, using temperature dependent material properties. Annotations and lines to show the mantle, core and core crystallisation period are added later, outside of the `pytesimal.load_plot_save` functions.

References

- Bryson, James F. J., Claire I. O. Nichols, Julia Herrero-Albillos, Florian Kronast, Takeshi Kasama, Hossein Alimadadi, Gerrit van der Laan, Francis Nimmo, and Richard J. Harrison. 2015. “Long-lived magnetism from solidification-driven convection on the pallasite parent body.” *Nature* 517 (7535): 472–75.
- Crank, J., and P. Nicolson. 1947. “A Practical Method for Numerical Evaluation of Solutions of Partial Differential Equations of the Heat-Conduction Type.” *Mathematical Proceedings of the Cambridge Philosophical Society* 43 (1): 50–67.
- Elkins-Tanton, Linda T., Benjamin P. Weiss, and Maria T. Zuber. 2011. “Chondrites as samples of differentiated planetesimals.” *Earth and Planetary Science Letters* 305 (1-2): 1–10.
- Fei, Yingwei. 2013. “Thermal Expansion.” In *Mineral Physics & Crystallography*, 29–44. American Geophysical Union (AGU).
- Haack, Henning, Kaare L. Rasmussen, and Paul H. Warren. 1990. “Effects of regolith/megaregolith insulation on the cooling histories of differentiated asteroids.” *Journal of Geophysical Research* 95 (B4): 5111–24.
- Murphy Quinlan, M., A. M. Walker, C. J. Davies, J. E. Mound, T. Müller, and J. Harvey. 2021. “The Conductive Cooling of Planetesimals with Temperature-Dependent Properties.” *Journal of Geophysical Research: Planets* 126 (4). <https://doi.org/https://doi.org/10.1029/2020JE006726>.
- Nichols, Claire I. O., James F. J. Bryson, Julia Herrero-Albillos, Florian Kronast, Francis Nimmo, and Richard J. Harrison. 2016. “Pallasite paleomagnetism: Quiescence of a core dynamo.” *Earth and Planetary Science Letters* 441: 103–12.
- Robie, R. A., B. S. Hemingway, and H. Takei. 1982. “Heat Capacities and Entropies of Mg_2SiO_4 , Mn_2SiO_4 , and Co_2SiO_4 Between 5 and 380 K.” *American Mineralogist* 67 (5-6): 470–82.
- Sahijpal, Sandeep. 2021. “Thermal Evolution of Non-Spherical Asteroids in the Early Solar System.” *Icarus* 362: 114439. <https://doi.org/https://doi.org/10.1016/j.icarus.2021.114439>.
- Su, Chang, Yonggang Liu, Wei Song, Dawei Fan, Zhigang Wang, and Hongfeng Tang. 2018. “Thermodynamic properties of San Carlos olivine at high temperature and high pressure.” *Acta Geochimica* 37 (2): 171–79.
- Suzuki, Isao. 1975. “Thermal Expansion of Periclase and Olivine, and their Anharmonic Properties.” *Journal of Physics of the Earth* 23 (2): 145–59.
- Xu, Yousheng, Thomas J Shankland, Sven Linhardt, David C Rubie, Falko Langenhorst, and Kurt Klasinski. 2004. “Thermal diffusivity and conductivity

of olivine, wadsleyite and ringwoodite to 20 GPa and 1373 K.” *Physics of the Earth and Planetary Interiors* 143-144: 321–36.