

DIAGNOSTIC IMAGING REPORT - PLAIN RADIOGRAPH

Riverside Orthopedic Imaging Center | ACR Accredited | STAT/Routine

Patient: Sam Rodriguez | DOB: 09/04/1972 | Sex: M | MRN: EHR-SR-001

Insurance: United Healthcare Choice Plus | Member ID: UHC9SR00287

Ordering Physician: Dr. James Patel, MD - Orthopedic Surgery | NPI: 0987654321

Radiologist: Dr. Marcus Webb, MD - Diagnostic Radiology | NPI: 5544332211

Exam: Right Knee - AP, Lateral, and Sunrise (Merchant) Views | 3 views

Date of Exam: January 14, 2026 | Reason: Right knee pain, suspected internal derangement

Clinical Indication: 4-month history of right knee pain following twisting injury. Rule out fracture, assess joint space.

TECHNIQUE

Three-view plain radiograph of the right knee performed: AP weight-bearing, lateral, and sunrise (Merchant) views. Radiation dose within standard diagnostic range. No contrast administered.

FINDINGS

Alignment: Normal tibiofemoral and patellofemoral alignment. No fracture, dislocation, or subluxation identified.

Bone density: Normal bone mineral density for patient age. No lytic or blastic lesions.

Joint spaces: Medial tibiofemoral joint space: mildly decreased (approximately 3.0 mm; reference >4 mm). Lateral tibiofemoral joint space: preserved. Patellofemoral joint space: preserved.

Articular surfaces: Mild irregularity of the medial tibial plateau. Subchondral sclerosis, medial compartment.

Osteophytes: Small marginal osteophytes at the medial femoral condyle and medial tibial plateau, consistent with early degenerative changes.

Soft tissues: Mild soft tissue swelling medial compartment. No calcification within joint space or periarticular soft tissues.

Patella: Patella: normal position and morphology. No bipartite patella.

Comparison: No prior imaging available for comparison.

IMPRESSION

1. No acute fracture or dislocation of the right knee.
2. Mild medial compartment joint space narrowing with early degenerative changes (small osteophytes, subchondral sclerosis medial tibial plateau). Kellgren-Lawrence Grade I-II.
3. Mild soft tissue swelling, medial compartment -- correlate with clinical examination.
4. Plain radiographs are limited for evaluation of soft tissue structures (menisci, ligaments, articular cartilage). If clinical suspicion for internal derangement remains, MRI is recommended for definitive soft tissue assessment.

RECOMMENDATION: Given the clinical history of twisting injury with positive meniscal provocation tests on examination and medial joint line tenderness, MRI of the right knee is recommended to evaluate for suspected meniscal pathology and associated chondral or ligamentous injury. Plain films alone are insufficient to exclude internal derangement.

Dr. Marcus Webb, MD