

ASSIGNMENT-13

M. polireddy
19, 32, 50, 56

- 1) Given an array of $[4, -2, 5, 3, 10, -5, 2, 8, -3, 6, 7, -4, 1, 9, -10, -6, -8, 11, -9]$ integers find the max and min product that can be obtained by multiplying two integers from the array.

Solution:-

Array is $[4, -2, 5, 3, 10, -5, 2, 8, -3, 6, 7, -4, 1, 9, -10, -6, -8, 11, -9]$

We need to consider the longest and smallest product that can be formed by selecting two numbers from the array.

1) Sort the array

sorted array

$[-9, -8, -6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]$

2) identify possible candidates for maximum product

3) identify possible candidates for minimum product

calculating maximum product:-

* The two longest positive numbers are 10 and 11

$$10 \times 11 = 110$$

* The two smallest negative numbers are -9 and -8

$$-9 \times -8 = 72$$

The maximum product is 110

calculating minimum product

The longest positive and negative numbers is 11 and -9

$$11 \times -9 = -99$$

The smaller positive negative numbers are

$$-9 \times -8 =$$

$$-9 \times -8 = 72$$

-99 is smaller than 72, 50

maximum product = 110 and minimum product = -99.



Shot on OnePlus

poli | 2024.06.25 07:00

- ⑧ Demonstrate the primary search method for the key = 23 from the array = {2, 5, 8, 2, 16, 23, 38, 56, 72, 91}.

Solution

Given key = 23 and array = {2, 5, 8, 2, 16, 23, 38, 56, 72, 91}

1. initialise pointers

low = 0 and high = 9

Calculate

$$\text{mid} = \left\lceil \frac{\text{low} + \text{high}}{2} \right\rceil = \left\lceil \frac{0 + 9}{2} \right\rceil = 4$$

Compare arr[mid] with key:

$$\text{arr}[4] = 16$$

Since $16 < 23$ update $\text{low} = \text{mid} + 1 = 5$

$$\text{Calculate mid} = \left\lceil \frac{\text{low} + \text{high}}{2} \right\rceil = \left\lceil \frac{5 + 9}{2} \right\rceil = 7$$

Compare arr[mid] with key:

$$\text{arr}[7] = 56$$

Since $56 > 23$ update $\text{high} = \text{mid} - 1 = 6$

$$\text{mid} = \left\lceil \frac{5 + 6}{2} \right\rceil = 5 \quad \text{arr}[\text{mid}] = \text{arr}[5] = 23$$


$23 = 23$ (The key is found at index 5.)

\therefore The key = 23 is found at index 5.

- 3) Apply merge sort and other list of 8 elements, Data = {5, 6, 7, 12, 15, 22, 30, 50, 20} set up a recurrence relation for the no. of key comparisons made by merge sort.

Q1

Mergesort



Find the no. of times to perform swapping for selection sort also estimate the time.



Shot on OnePlus

poli 1 2024.06.25 07:01

Recursive Relation for comparisons.

$$T(n) = 2T(n/2) + O(n)$$

If $n=1$, $T(1) = 0$,

→ At each level of recursion we make at most $n-1$ comparisons to merge two halves of size $n/2$ so it becomes,

$$T(n) = 2T(n/2) + (n-1)$$

Solve recurrence relation we get

$$T(n) = n \log_2(n) - n + 1$$

$$\therefore T(n) = O(n \log n)$$

→ The recurrence Relation is $T(n) = 2T(n/2) + O(n)$

Or more precisely

$$T(n) = n \log_2(n) - n + 1$$

5) find the no of times of perform swapping for selection sort also estimate the p.c for the order of notation set $S(12, 7, 5, -2, 18, 6, 13, 4)$

The selection sort algorithm always moves exactly $n-1$ swaps in the worst case, where n is the no of elements in the list

given $S = \{12, 7, 5, -2, 18, 6, 13, 4\}$

No of elements $n = 8$

$$\text{No of swaps} = n-1 = 8-1 = 7$$

Time complexity:- $O(n^2)$

So the no of swaps is 7, and the p.c is $O(n^2)$.



Shot on OnePlus

poll | 2024.06.25 07:02

Find the index the target value 10 using binary search from the following list of elements [2, 4, 6, 8, 10, 12, 14, 16, 18, 20]

Given list = {2, 4, 6, 8, 10, 12, 14, 16, 18, 20} and val = 10

low = 0 and high = 9

$$\text{mid} = \frac{\text{low} + \text{high}}{2} = \frac{0 + 9}{2} = 4$$

mid = 10 mid == value

Since 10 == 10 the target is found at index 4

∴ The target value = 10 is found at index 4.

