

> West Afr J Med. Sep-Oct 2011;30(5):325-30.

Dysglycaemia and its risk factors in an urban Fulani population of northern Nigeria

A A Sabir¹, S A Isezuo, A E Ohwovoriole

Affiliations + expand

PMID: 22752819

Abstract

Background: The Fulani are a normally nomadic people known for covering great distances on foot with a resulting lean physique and presumably low incidence of diabetes mellitus (DM). The occurrence of glucose intolerance among the urban Nigerian Fulani has not been reported.

Objective: To determine the prevalence of dysglycaemia and its correlates among urban Fulani.

Methods: Three hundred and eighty-nine Fulani were recruited from urban areas of Sokoto state using a multi-stage sampling technique. Using a modification of the WHO STEPS instrument, information on socio-demographic and anthropometric data was gathered. Either casual or fasting plasma glucose was obtained in all subjects while oral glucose tolerance test (OGTT) was performed in a randomly selected subset of 48 subjects. Glucose intolerance or dysglycaemia was defined using WHO criteria as the presence of diabetes mellitus, impaired fasting glycaemia (IFG) or impaired glucose tolerance (IGT). Classifications of adiposity as body mass index (BMI) and waist circumference (WC) and blood pressure were made using standard criteria.

Results: Of the 389 subjects studied, 190 (48.8%) were females while 199 (51.2%) subjects were males (c2=5.43, p> 0.05). The overall mean (SD) age of the sample population was 39.3 (14.2) years with the males [42.0 (13.8) years] being significantly older than the females [36.4 (14.1) years], p< 0.05. Eighteen (4.6%) subjects had previously undiagnosed type 2 diabetes mellitus while 37(16.9%) of the subjects had IFG and seven (14.6%) subjects had impaired glucose tolerance. The mean FPG was higher in the males [5.49 (1.7) mmol/l] than in the females [5.25 (1.9) mmol/l],p=0.33. The overall mean two-hour plasma post glucose-load was 6.5 (1.6) mmol/l; higher in the males [6.9 (2)] mmol/l than in the females [6.2 (1) mmol/l] , p=0.55.The mean age, BMI, waist circumference and blood pressure of subjects with dysglycaemia were significantly higher than those in subjects with normoglycaemia (p<0.05).

Conclusion: Prevalence of dysglycaemia in the urban Fulani population of Northern Nigeria is high and much higher in the male than female Fulani. Occurrence of obesity and hypertension appears lower than in other Nigerian reports. Occurrence of dysglycaemia in the urban Fulani increases with age, weight and waist circumference. There is need to conduct a similar study among rural Fulani to better appreciate the environmental influence on these variables.

Similar articles

Type 2 diabetes mellitus and its risk factors among the rural Fulanis of Northern Nigeria.

Sabir A, Ohwovoriole A, Isezuo S, Fasanmade O, Abubakar S, Iwuala S.

Ann Afr Med. 2013 Oct-Dec;12(4):217-22. doi: 10.4103/1596-3519.122689.

PMID: 24309410

A large proportion of prediabetes and diabetes goes undiagnosed when only fasting plasma glucose and/or HbA1c are measured in overweight or obese patients.

Cosson E, Hamo-Tchatchouang E, Banu I, Nguyen MT, Chiheb S, Ba H, Valensi P.

Diabetes Metab. 2010 Sep;36(4):312-8. doi: 10.1016/j.diabet.2010.02.004.

PMID: 20627649

Glucose intolerance (diabetes and IGT) in a selected South Indian population with special reference to family history, obesity and lifestyle factors--the Chennai Urban Population Study (CUPS 14).

Mohan V, Shanthirani CS, Deepa R.

J Assoc Physicians India. 2003 Aug;51:771-7.

PMID: 14651136

THE DIABETES EPIDEMIC IN CHINA: AN INTEGRATED REVIEW OF NATIONAL SURVEYS.

Shen X, Vaidya A, Wu S, Gao X.

Endocr Pract. 2016 Sep;22(9):1119-29. doi: 10.4158/EP161199.RA. Epub 2016 Jun 13.

PMID: 27295015 [Free PMC article.](#) [Review.](#)

Estimating the prevalence, hospitalisation and mortality from type 2 diabetes mellitus in Nigeria: a systematic review and meta-analysis.

Adeloye D, Ige JO, Aderemi AV, Adeleye N, Amoo EO, Auta A, Oni G.

BMJ Open. 2017 May 11;7(5):e015424. doi: 10.1136/bmjopen-2016-015424.

PMID: 28495817 [Free PMC article.](#) [Review.](#)

See all similar articles

Cited by 7 articles

Mobile phone ownership and willingness to receive mHealth services among patients with diabetes mellitus in South-West, Nigeria.

Olamoyegun MA, Raimi TH, Ala OA, Fadare JO.

Pan Afr Med J. 2020 Sep 8;37:29. doi: 10.11604/pamj.2020.37.29.25174. eCollection 2020.

PMID: 33062130 [Free PMC article.](#)

The Burden of Undiagnosed Diabetes Mellitus in Adult African Population: A Systematic Review and Meta-Analysis.

Asmelash D, Asmelash Y.

J Diabetes Res. 2019 Apr 28;2019:4134937. doi: 10.1155/2019/4134937. eCollection 2019.

PMID: 31183383 [Free PMC article.](#)

Prevalence and correlates of traditional risk factors for cardiovascular disease in a Nigerian ART-naïve HIV population: a cross-sectional study.

Ekrikpo UE, Akpan EE, Ekott JU, Bello AK, Okpechi IG, Kengne AP.

BMJ Open. 2018 Jul 19;8(7):e019664. doi: 10.1136/bmjopen-2017-019664.

PMID: 30030310 [Free PMC article.](#)

Prevalence and Risk Factors for Diabetes Mellitus in Nigeria: A Systematic Review and Meta-Analysis.

Uloko AE, Musa BM, Ramalan MA, Gezawa ID, Puepet FH, Uloko AT, Borodo MM, Sada KB.

Diabetes Ther. 2018 Jun;9(3):1307-1316. doi: 10.1007/s13300-018-0441-1. Epub 2018 May 14.

PMID: 29761289 [Free PMC article.](#)

Effect of short and long moderate-intensity exercises in modifying cardiometabolic markers in sedentary Kenyans aged 50 years and above.

Magutah K, Meiring R, Patel NB, Thairu K.

BMJ Open Sport Exerc Med. 2018 Apr 25;4(1):e000316. doi: 10.1136/bmjsem-2017-000316. eCollection 2018.

PMID: 29719726 [Free PMC article.](#)

See all "Cited by" articles

MeSH terms

- > Adolescent

> Adult

> Age Distribution

> Aged

> Blood Glucose / metabolism*

> Body Mass Index

> Female

> Glucose Intolerance / diagnosis

> Glucose Intolerance / epidemiology*

> Humans

> Incidence

> Male

> Middle Aged

> Nigeria / epidemiology

> Prevalence

> Risk Factors

> Sex Distribution

> Urban Population*

> Young Adult

Substances

> Blood Glucose

Related information

MedGen

PubChem Compound

PubChem Substance

LinkOut – more resources

Medical

MedlinePlus Health Information

Miscellaneous

NCI CPTAC Assay Portal

ACTIONS

“ Cite

☆ Favorites

SHARE



PAGE NAVIGATION

Title & authors

< Abstract

Similar articles

Cited by

MeSH terms

Substances

Related information

LinkOut - more resources

