Frequency of Abstract Words for Years 2007 to 2013


model
data
method
propose
study
analysis
estimate
effect
approach
base
test
time
sample
simulation
likelihood
regression
estimator
bayes
parameter
measure
process
multiple
patient
association
longitudinal
survival
clinical
factor
assumption
biomarker
covariate
treatment
variable
distribution
statistic
inference
performance
interaction
procedure
gene
function
disease
trial
outcome
illustrate
design
cluster
power
prior
challenge
comparison
dependence
change
value
different
present
identify
information
problem
paper
algorithm
standard
functional
perform
equation
markov
obtained
known
expected
involving
distributed
technology
role
much
dynamic
novel
mcmc
quantitative
scientific
mechanism
brain
nonparametric
assess
measurement
randomize
variation
marginal
simulate
detection
advantage
recently
limited
end
carry
community
learning
behavior
microarray
traditional
assessment
national
influence
accuracy
validation
effectiveness
popular
even
matrix
image
space
low
involve
define
blood
source
amount
furthermore
experimental
medical
monitor
quality
family
posterior
basis
good
scenario
exact
arise
growth
incidence
semiparametric
investigated
evaluation
extend
temporal
independence
particularly
indicate
main
powerful
phenotype
death
theory
especially
kernel
regarding
quantify
evaluating
mapping
understanding
competing
researcher
software
called
determined
accommodate
computation
highly
made
right
subgroup
pm
generally
calculate
critical
target
approximate
limitations
substantial
generation
analytic
interpretation
future
established
nonlinear
environment
attention
extended
infection
simultaneous
strong
genotype
birth
assignment
pollution
threshold
throughput
hospital
achieve
reported
implementation
induce
observational
monte
best
shown
assessment
observe
high
level
estimation
size
population
potential
exposure
toxicity
participant
penalty
assume
consider
log
focus
better
proportional
degree
capture
subject
demonstrate
cancer
feature
system
environmental
categorical
evaluate
hierarchical
characterize
region
status
detecting
surrogate
investigate
review
roc
class
setting
consistent
accurate
operating
provide
trait
positive
coverage
across
variant
unit
finding
activity
asymptotic
random
spline
explore
needed
event
expression
location
project
logistic
obtain
assessing
easily
compare
analyze
motivate
endpoint
reduction
prevalence
mean
normal
square
frequent
contrast
rule
health
cross
constraint
applying
specific
reduce
gwas
new
predict
joint
inverse
short
one
finite
challenging
predictor
case
exist
probability
empirical
criterion
yield
exponential
weight
genomic
repeat
odd
natural
curve
usually
count
employ
censor
introduce
criteria
play
addition
differential
mass
increase
latent
efficacy
become
optimal
certain
support
within
technique
detect
instead
first
dependent
carlo
genome
cohort
able
require
typically
finally
development
sources
semi
give
property
common
rather
mortality
relate
pathway
hazard
without
cell
available
area
practice
vector
early
distance
discuss
strategy
collect
loss
range
characteristic
relative
pair
interim
frailty
progression
key
pregnancy
polymorphism
intensity
program
adjustment
robustness
separate
complexity
transformation
valid
stochastic
designed
specifically
help
next
copy
nature
specificity
assuming
currently
lack
counts
significance
agreement
cumulative
occur
unbiased
initial
sequential
additive
principal
computationally
adverse
handle
match
express
uncertainty
poisson
assumed
recurrent
produce
intervention
another
likely
courses
smoking
respect
allowing
controls
index
theoretical
average
unknown
specified
univariate
graphical
residual
chain
step
significant
imputation
classification
informative
ability
usual
examined
higher
practical
impact
situation
commonly
fmri
diagnostic
tree
sparse
corresponding
predicting
classical
suggest
incorporate
difference
condition
make
implement
long
bivariate
period
jointly
safety
least
use
follow
sensitive
dataset
combination
gaussian
diagnosis
thousand
cause
primary
robust
decision
chip
coefficient
improve
overall
complicate
evidence
literature
pattern
maximum
relationship
parametric
independent
variability
partial
stratification
combining
need
appropriate
form
since
account
several
research
methodology
depend
smoothing
represent
length
studied
benefit
smaller
multivariate
local
framework
structure
network
sensitivity
particular
tumor
shape
frequency
aspect
pre
survey
alternative
component
real
genetic
individual
due
biological
must
utility
combined
hypothesis
allow
dimension
group
correlation
given
conditional
simultaneouslygee
working
significantly
flexible
either
cox
presencediscrete
purpose
composite
structural
great
limit
adaptive
false
covariance
null
along
making
normality
causal
generalize
control
small
tool
potential
rank
article
whose
difficult
although
biomedical
effective
efficient
meta
fix
exposure
log
focus
applicable
true
vary
ratio
scale
stage
include
risk
distribution
show
reference
propensity
post
address
shrinkage
density
left
interest
single
identified
binary
develop
result
two
error
fit
restricted
graph
correct
examine
directly
context
work
type
care
useful
select
lung
penalized
little
large
direct
previously
breast
question
year
adjust
derive
efficiency
application
regulatory
major
continuous
ordinal
therapy
possible
lead
conventional
fully
take
issue
validity
objective
experiment
confounding
field
signal
various
heart
center
proportion
baseline
four
second
failure
additional
extensive
three
lower
found
describe
identification
precision
relevant
less
current
affect
generate
phase
mixture
complete
determine
general
potentially
discovery
screening
conduct
conclusion
free
similar
correlate
connectivity
link
underlying
summary
part
quantile
em
recent
choice
relatively
heterogeneity
extreme
presentation
frequentist
bootstrap
protein
previous
following
numerical
dependence
women
haplotype
distributed
involving
expected
amount
furthermore
required
monitor
good
scenario
exact
arise
growth
incidence
investigated
evaluation
extend
independence
particularly
indicate
mapping
accommodate
computation
highly
substantial
generation
analytic
interpretation
future
established
nonlinear
environment
attention
extended
infection
simultaneous
strong
genotype
birth
assignment
pollution
threshold
throughput
hospital
achieve
reported
implementation
induce
limitations
approximate
critical
target
determined
calculate
generally
subgroup
pm
right
made
understanding
competing
researcher
software
called
evaluating
death
phenotype
powerful
main
indicate
particularly
independence
temporal
extend
evaluation
investigated
exact
arise
growth
incidence
semiparametric
detection
advantage
recently
limited
end
carry
community
learning
behavior
microarray
traditional
assessment
national
influence
accuracy
validation
effectiveness
popular
even
matrix
image
space
low
involve
define
blood
source
amount
furthermore
experimental
medical
monitor
quality
family
posterior
basis
good
scenario
observational
monte
best
shown
assess
measurement
randomize
variation
marginal
simulate
detection
advantage
recently
limited
end
carry
community
learning
behavior
microarray
traditional
assessment
national
influence
accuracy
validation
effectiveness
popular
even
matrix
image
space
low
involve
define
blood
source
amount
furthermore
experimental
medical
monitor
quality
family
posterior
basis
good
scenario
exact
arise
growth
incidence
semiparametric
investigated
evaluation
extend
temporal

Frequency of Abstract Words for Years 2007 to 2013




Frequency of Abstract Words for Years 2007 to 2013
model
data
method
study
propose
test
approach
analysis
estimate
effect
base
trial
sample
two
bayes
multiple
likelihood
function
treatment
regression
simulation
variable
statistic
estimator
distribution
parameter
estimation
covariate
longitudinal
bias
time
risk
probability
genetic
cancer
disease
design
outcome
model data
study estimate
analysis variable
statistic estimator
distribution parameter
estimation covariate
longitudinal bias
time risk
probability genetic
cancer disease
design outcome