

# From preferences to rewards: A brief introduction

Fernando E. Rosas

Iliad Intensive

August 11, 2026

# The Question

- Agents are powerful — but increasing capabilities lead to increasing risks.

# The Question

- Agents are powerful — but increasing capabilities lead to increasing risks.
- To understand agentic risk, we need to understand how agents work.

# The Question

- Agents are powerful — but increasing capabilities lead to increasing risks.
- To understand agentic risk, we need to understand how agents work.
- Core of agency: *doing things (planning) for reasons (beliefs)*.


# The Question

- Agents are powerful — but increasing capabilities lead to increasing risks.
- To understand agentic risk, we need to understand how agents work.
- Core of agency: *doing things (planning) for reasons (beliefs)*.
- What are goals? Hard question.

# The Question


- Agents are powerful — but increasing capabilities lead to increasing risks.
- To understand agentic risk, we need to understand how agents work.
- Core of agency: *doing things (planning) for reasons (beliefs)*.
- What are goals? Hard question.
- **Goal of today:** understand what are preferences.

# The Route



- Each arrow adds structure.


# The Route



- Each arrow adds structure.
- Each structure buys a more specific representation.




# The Route



- Each arrow adds structure.
- Each structure buys a more specific representation.
- *Outcome 1*: Understand the assumptions of each step.

# The Route



- Each arrow adds structure.
- Each structure buys a more specific representation.
- *Outcome 1*: Understand the assumptions of each step.
- *Outcome 2*: Recognise the difference between preference, utility, and reward.

# Basic setting

- Let  $\mathcal{O}$  be a finite observation set, and  $\mathcal{A}$  a finite action set.

# Basic setting

- Let  $\mathcal{O}$  be a finite observation set, and  $\mathcal{A}$  a finite action set.
- A one-step interaction is

$$t = (o, a) \in \mathcal{O} \times \mathcal{A}.$$

# Basic setting

- Let  $\mathcal{O}$  be a finite observation set, and  $\mathcal{A}$  a finite action set.
- A one-step interaction is

$$t = (o, a) \in \mathcal{O} \times \mathcal{A}.$$

- A finite trajectory of length  $n$  is

$$h = (o_1, a_1, o_2, a_2, \dots, o_n, a_n) \in (\mathcal{O} \times \mathcal{A})^n.$$

# Basic setting

- Let  $\mathcal{O}$  be a finite observation set, and  $\mathcal{A}$  a finite action set.
- A one-step interaction is

$$t = (o, a) \in \mathcal{O} \times \mathcal{A}.$$

- A finite trajectory of length  $n$  is

$$h = (o_1, a_1, o_2, a_2, \dots, o_n, a_n) \in (\mathcal{O} \times \mathcal{A})^n.$$

- The full finite trajectory space is

$$\mathcal{H}^* = \bigcup_{n=0}^{\infty} (\mathcal{O} \times \mathcal{A})^n.$$

# Preferences

- A weak preference relation  $\succsim$  compares complete trajectories.

$$h \succsim h'$$

means “ $h$  is at least as good as  $h'$ .”

- A weak preference relation  $\succsim$  compares complete trajectories.

$$h \succsim h'$$

means “ $h$  is at least as good as  $h'$ .”

- From this we define:

$$h \sim h' \iff h \succsim h' \text{ and } h' \succsim h,$$

$$h \succ h' \iff h \succsim h' \text{ and not } h' \succsim h.$$



- A weak preference relation  $\succsim$  compares complete trajectories.

$$h \succsim h'$$

means “ $h$  is at least as good as  $h'$ .”

- From this we define:

$$h \sim h' \iff h \succsim h' \text{ and } h' \succsim h,$$

$$h \succ h' \iff h \succsim h' \text{ and not } h' \succsim h.$$

- At this point,  $\succsim$  need not be complete, transitive, numerical, or ‘reward-like’.

# Three Kinds of Trouble

By choosing the “wrong type of preference”, an agent can run into three kinds of trouble (from less to more harmful):

- **Representability failure:** preference has no convenient formal representation.

# Three Kinds of Trouble

By choosing the “wrong type of preference”, an agent can run into three kinds of trouble (from less to more harmful):

- **Representability failure:** preference has no convenient formal representation.
- **Self-defeat:** agents take suboptimal choices or plans — there are better alternatives that are not being taken.

# Three Kinds of Trouble

By choosing the “wrong type of preference”, an agent can run into three kinds of trouble (from less to more harmful):


- **Representability failure:** preference has no convenient formal representation.
- **Self-defeat:** agents take suboptimal choices or plans — there are better alternatives that are not being taken.
  - ▷ E.g. **dominance:** another policy yields preferred trajectories at least as preferred in every state of the world, and strictly so in at least one.

# Three Kinds of Trouble

By choosing the “wrong type of preference”, an agent can run into three kinds of trouble (from less to more harmful):


- **Representability failure:** preference has no convenient formal representation.
- **Self-defeat:** agents take suboptimal choices or plans — there are better alternatives that are not being taken.
  - ▷ E.g. **dominance:** another policy yields preferred trajectories at least as preferred in every state of the world, and strictly so in at least one.
- **Vulnerability:** individually acceptable trades can combine into guaranteed loss.
  - ▷ E.g. **Money pumps:** circular preferences ( $h_1 \succ h_2 \succ h_3 \succ h_1$ ) lead to indefinite loss.

# Money Pump Picture




- If the agent pays for every strict improvement, a cycle can be exploited.

# Money Pump Picture



- If the agent pays for every strict improvement, a cycle can be exploited.
- It returns to the same trajectory with less money.

# Money Pump Picture



- If the agent pays for every strict improvement, a cycle can be exploited.
- It returns to the same trajectory with less money.
- Parallel with thermodynamics: the second law of thermodynamics can be read as stating that *'nature cannot be money-pumped'*.



# Three Kinds of Results

There are three types of results regarding preferences:

- **Representation results:** axioms imply that preferences can be written in a particular mathematical form.
  - ▷ E.g. some preferences can be represented as if maximising a utility function.

# Three Kinds of Results

There are three types of results regarding preferences:

- **Representation results:** axioms imply that preferences can be written in a particular mathematical form.
  - ▷ E.g. some preferences can be represented as if maximising a utility function.
- **Coherence results:** violations imply some penalty for the agent.
  - ▷ E.g. money pumps or Dutch books.

# Three Kinds of Results

There are three types of results regarding preferences:

- **Representation results:** axioms imply that preferences can be written in a particular mathematical form.
  - ▷ E.g. some preferences can be represented as if maximising a utility function.
- **Coherence results:** violations imply some penalty for the agent.
  - ▷ E.g. money pumps or Dutch books.
- **Selection results:** some optimization pressure tends to remove agents with a given failure mode.
  - ▷ E.g. market competition or natural evolution.

# Three Kinds of Results

There are three types of results regarding preferences:

- **Representation results:** axioms imply that preferences can be written in a particular mathematical form.
  - ▷ E.g. some preferences can be represented as if maximising a utility function.
- **Coherence results:** violations imply some penalty for the agent.
  - ▷ E.g. money pumps or Dutch books.
- **Selection results:** some optimization pressure tends to remove agents with a given failure mode.
  - ▷ E.g. market competition or natural evolution.

These results are related (one can lead to the other) but not identical: form, vulnerability, and survival are different claims.

# Axiom 1: Completeness

Completeness says every pair can be compared:

$$\forall h, h' \in \mathcal{H}^*, \quad h \succcurlyeq h' \quad \text{or} \quad h' \succcurlyeq h.$$

# Axiom 1: Completeness

Completeness says every pair can be compared:

$$\forall h, h' \in \mathcal{H}^*, \quad h \succcurlyeq h' \quad \text{or} \quad h' \succcurlyeq h.$$

- Incomparability is not the same as indifference.

# Axiom 1: Completeness

Completeness says every pair can be compared:

$$\forall h, h' \in \mathcal{H}^*, \quad h \succcurlyeq h' \quad \text{or} \quad h' \succcurlyeq h.$$

- Incomparability is not the same as indifference.
- If it fails, some trajectories are genuinely incomparable.

## Axiom 2:Transitivity

Transitivity says comparisons fit together:

$$h \succ h' \text{ and } h' \succ h'' \implies h \succ h''.$$



## Axiom 2:Transitivity

Transitivity says comparisons fit together:

$$h \succ h' \text{ and } h' \succ h'' \implies h \succ h''.$$

- If it fails, local pairwise judgments need not form a global ranking.

## Axiom 2: Transitivity

Transitivity says comparisons fit together:

$$h \succ h' \text{ and } h' \succ h'' \implies h \succ h''.$$

- If it fails, local pairwise judgments need not form a global ranking.
- The classic failure is a cycle: money pumps.

$$h_1 \succ h_2, \quad h_2 \succ h_3, \quad h_3 \succ h_1.$$

- Similar to Helmholtz-Hodge decomposition: transitivity is equivalent to requiring no rotational component.

# Axioms 1 + 2 $\rightarrow$ Ordinal Utility

- Completeness + transitivity make  $\succsim$  a weak order (total order with equivalent objects).

# Axioms 1 + 2 $\rightarrow$ Ordinal Utility

- Completeness + transitivity make  $\succsim$  a weak order (total order with equivalent objects).
- On a countable space  $\mathcal{H}^*$ , weak orders admit an ordinal utility: there exists a function  $u : \mathcal{H}^* \rightarrow \mathbb{R}$  such that

$$h \succsim h' \iff u(h) \geq u(h').$$

# Axioms 1 + 2 $\rightarrow$ Ordinal Utility

- Completeness + transitivity make  $\succsim$  a weak order (total order with equivalent objects).
- On a countable space  $\mathcal{H}^*$ , weak orders admit an ordinal utility: there exists a function  $u : \mathcal{H}^* \rightarrow \mathbb{R}$  such that

$$h \succsim h' \iff u(h) \geq u(h').$$

- Only the ranking matters:

$$u, \quad 2u + 17, \quad \exp(u)$$

all represent the same deterministic preferences.

# Beyond Ordinal Utility

Ordinal utility ranks outcomes, but give a structure to mixtures of them.

- Example:

$$x \succ y \succ z.$$

How should we rank  $y$  with respect to  $\frac{1}{2}x + \frac{1}{2}z$ ? Should they be equivalent?

# Beyond Ordinal Utility

Ordinal utility ranks outcomes, but give a structure to mixtures of them.

- Example:

$$x \succ y \succ z.$$

How should we rank  $y$  with respect to  $\frac{1}{2}x + \frac{1}{2}z$ ? Should they be equivalent?

Ordinal rank doesn't say how to rank mixtures, which can arise naturally out of uncertainty.

# Lotteries Over Trajectories

- Let  $\Delta(X)$  be the set of probability distributions over  $X \subseteq \mathcal{H}^*$ .



# Lotteries Over Trajectories

- Let  $\Delta(X)$  be the set of probability distributions over  $X \subseteq \mathcal{H}^*$ .
- A lottery is written

$$\mu = \sum_i p_i \delta_{x_i}.$$

# Lotteries Over Trajectories

- Let  $\Delta(X)$  be the set of probability distributions over  $X \subseteq \mathcal{H}^*$ .
- A lottery is written

$$\mu = \sum_i p_i \delta_{x_i}.$$

- A sure trajectory  $x$  is the degenerate lottery  $\delta_x$ .

# Lotteries Over Trajectories


- Let  $\Delta(X)$  be the set of probability distributions over  $X \subseteq \mathcal{H}^*$ .
- A lottery is written

$$\mu = \sum_i p_i \delta_{x_i}.$$

- A sure trajectory  $x$  is the degenerate lottery  $\delta_x$ .
- We can mix lotteries:

$$\lambda\mu + (1 - \lambda)\nu.$$

# Lottery Picture



- vNM asks when preferences over these objects are expectations.

# vNM representaiton: The big picture

- 1 **Axiom 1:Completeness:** any two lotteries can be compared.

# vNM representaiton: The big picture

- ① **Axiom 1:Completeness:** any two lotteries can be compared.
- ② **Axiom 2:Transitivity:** comparisons over lotteries fit together.

# vNM representaiton: The big picture

- ① **Axiom 1:Completeness:** any two lotteries can be compared.
- ② **Axiom 2:Transitivity:** comparisons over lotteries fit together.
- ③ **Axiom 3:Continuity:** intermediate prospects can be matched by a mixture of better and worse prospects.

# vNM representaiton: The big picture

- ① **Axiom 1:Completeness:** any two lotteries can be compared.
- ② **Axiom 2:Transitivity:** comparisons over lotteries fit together.
- ③ **Axiom 3:Continuity:** intermediate prospects can be matched by a mixture of better and worse prospects.
- ④ **Axiom 4:Independence:** common lottery components cancel.



# vNM representaiton: The big picture

- ① **Axiom 1:Completeness:** any two lotteries can be compared.
- ② **Axiom 2:Transitivity:** comparisons over lotteries fit together.
- ③ **Axiom 3:Continuity:** intermediate prospects can be matched by a mixture of better and worse prospects.
- ④ **Axiom 4:Independence:** common lottery components cancel.

**Result:** The first two axioms give ranking structure based on a cardinal utility; the last two axioms make the utility to be an expected value.

# Axiom 3: Continuity

- If  $\mu \succcurlyeq \nu \succcurlyeq \xi$ , continuity says there always exists some value  $\lambda \in [0, 1]$  such that

$$\nu \sim \lambda\mu + (1 - \lambda)\xi.$$

# Axiom 3: Continuity

- If  $\mu \succcurlyeq \nu \succcurlyeq \xi$ , continuity says there always exists some value  $\lambda \in [0, 1]$  such that

$$\nu \sim \lambda\mu + (1 - \lambda)\xi.$$

- Intuition: no outcome is infinitely better or worse than the others.

# Axiom 3: Continuity

- If  $\mu \succcurlyeq \nu \succcurlyeq \xi$ , continuity says there always exists some value  $\lambda \in [0, 1]$  such that

$$\nu \sim \lambda\mu + (1 - \lambda)\xi.$$

- Intuition: no outcome is infinitely better or worse than the others.

Classic counter-examples are lexicographic preferences, in which events can be infinitely different from each other.

*Example:*  $\mu$  is eating chocolate,  $\nu$  is eating potatoes, and  $\xi$  is accepting genocide.

## Axiom 4: Independence

- If  $\mu \succcurlyeq \nu$ , then mixing both with the same background lottery  $\xi$  should not reverse the comparison:

$$\mu \succcurlyeq \nu \iff \lambda\mu + (1 - \lambda)\xi \succcurlyeq \lambda\nu + (1 - \lambda)\xi.$$

# Axiom 4: Independence

- If  $\mu \succcurlyeq \nu$ , then mixing both with the same background lottery  $\xi$  should not reverse the comparison:

$$\mu \succcurlyeq \nu \iff \lambda\mu + (1 - \lambda)\xi \succcurlyeq \lambda\nu + (1 - \lambda)\xi.$$

- Independence is what gives linear averaging.

# Axiom 4: Independence

- If  $\mu \succcurlyeq \nu$ , then mixing both with the same background lottery  $\xi$  should not reverse the comparison:

$$\mu \succcurlyeq \nu \iff \lambda\mu + (1 - \lambda)\xi \succcurlyeq \lambda\nu + (1 - \lambda)\xi.$$

- Independence is what gives linear averaging.

When it fails, common consequences do not psychologically or behaviorally cancel.

*Example (Allais paradox):* Most people would say

- Getting \$100 for sure *is better than* a coin flip for \$250.
- A 1-in-100 shot at \$100 *is worst than* a 1-in-200 shot at \$250.

These are the same choice, just scaled down.

# Theorem 1: The von Neumann-Morgenstern representation

Assumptions: a finite set of options  $X \subseteq \mathcal{H}^*$ , the four axioms hold.

- **Result 1 (existence):** there exists  $u : X \rightarrow \mathbb{R}$  such that

$$\mu \succsim \nu \iff \sum_{x \in X} \mu(x) u(x) \geq \sum_{x \in X} \nu(x) u(x).$$



# Theorem 1: The von Neumann-Morgenstern representation

Assumptions: a finite set of options  $X \subseteq \mathcal{H}^*$ , the four axioms hold.

- **Result 1 (existence):** there exists  $u : X \rightarrow \mathbb{R}$  such that

$$\mu \succsim \nu \iff \sum_{x \in X} \mu(x) u(x) \geq \sum_{x \in X} \nu(x) u(x).$$

- **Result 2 (uniqueness):**  $u$  is unique up to positive affine transformations:

$$u'(x) = a + bu(x), \quad b > 0.$$

## Ordinal utility

- Any set of outcomes
- Preserves ranking
- Indifferent to non-decreasing transformation
- Cannot display 'meaningful gaps'

## vNM utility

- Lotteries (probabilistic mixtures)
- Preserves risky choices
- Only affine transforms are okay
- Inconmesurable gaps can exist

## Ordinal utility

- Any set of outcomes
- Preserves ranking
- Indifferent to non-decreasing transformation
- Cannot display 'meaningful gaps'

## vNM utility

- Lotteries (probabilistic mixtures)
- Preserves risky choices
- Only affine transforms are okay
- Inconmesurable gaps can exist

vNM utility further refines ordinal utility (it is not just a relabeling).

# Beyond expected utility: Temporal structure

- vNM utility assigns a single scalar to a whole trajectory.


# Beyond expected utility: Temporal structure

- vNM utility assigns a single scalar to a whole trajectory.
- But agents make decisions over time, so they need indications of utility that unfold step-by-step.

# Beyond expected utility: Temporal structure

- vNM utility assigns a single scalar to a whole trajectory.
- But agents make decisions over time, so they need indications of utility that unfold step-by-step.
- From a global utility  $u(h)$ , one can always define the increment


$$r(t; h) = u(t \cdot h) - u(h), \quad \text{satisfying} \quad u(t \cdot h) = r(t; h) + u(h).$$



# Beyond expected utility: Temporal structure

- vNM utility assigns a single scalar to a whole trajectory.
- But agents make decisions over time, so they need indications of utility that unfold step-by-step.
- From a global utility  $u(h)$ , one can always define the increment

$$r(t; h) = u(t \cdot h) - u(h), \quad \text{satisfying} \quad u(t \cdot h) = r(t; h) + u(h).$$



- But the increment  $r(t; h)$  may be non-local, and hence cannot be optimised for directly.

# Axiom 5: Temporal equanimity

Let  $T = \mathcal{O} \times \mathcal{A}$  be the transition set.

- The fifth axiom says there is a function  $\gamma : T \rightarrow [0, 1]$  such that prepending  $t$  rescales continuation differences, so that

$$u(t \cdot \mu) - u(t \cdot \nu) = \gamma(t)(u(\mu) - u(\nu)).$$

$\gamma(t)$  measures how much the future still matters after  $t$ .



# Axiom 5: Temporal equanimity

Let  $T = \mathcal{O} \times \mathcal{A}$  be the transition set.

- The fifth axiom says there is a function  $\gamma : T \rightarrow [0, 1]$  such that prepending  $t$  rescales continuation differences, so that

$$u(t \cdot \mu) - u(t \cdot \nu) = \gamma(t)(u(\mu) - u(\nu)).$$

$\gamma(t)$  measures how much the future still matters after  $t$ .

- $\gamma(t) = 1$ : future differences matter as much as present differences.

# Axiom 5: Temporal equanimity

Let  $T = \mathcal{O} \times \mathcal{A}$  be the transition set.

- The fifth axiom says there is a function  $\gamma : T \rightarrow [0, 1]$  such that prepending  $t$  rescales continuation differences, so that

$$u(t \cdot \mu) - u(t \cdot \nu) = \gamma(t)(u(\mu) - u(\nu)).$$

$\gamma(t)$  measures how much the future still matters after  $t$ .

- $\gamma(t) = 1$ : future differences matter as much as present differences.
- $0 < \gamma(t) < 1$ : future differences are damped.

# Axiom 5: Temporal equanimity

Let  $T = \mathcal{O} \times \mathcal{A}$  be the transition set.

- The fifth axiom says there is a function  $\gamma : T \rightarrow [0, 1]$  such that prepending  $t$  rescales continuation differences, so that

$$u(t \cdot \mu) - u(t \cdot \nu) = \gamma(t)(u(\mu) - u(\nu)).$$

$\gamma(t)$  measures how much the future still matters after  $t$ .

- $\gamma(t) = 1$ : future differences matter as much as present differences.
- $0 < \gamma(t) < 1$ : future differences are damped.
- $\gamma(t) = 0$ : after  $t$ , continuation does not matter.

This is transition-dependent discounting, not necessarily a single global constant.

# Theorem 2: Markov reward representation

Assumptions: Axioms 1-5 over lotteries over trajectories.

- Then, utility has a local recursive form:

$$u(t \cdot h) = r(t) + \gamma(t)u(h), \quad u(\varepsilon) = 0.$$

## Theorem 2: Markov reward representation

Assumptions: Axioms 1-5 over lotteries over trajectories.

- Then, utility has a local recursive form:

$$u(t \cdot h) = r(t) + \gamma(t)u(h), \quad u(\varepsilon) = 0.$$

- Preferences over lotteries are still represented by expected utility.

## Theorem 2: Markov reward representation

Assumptions: Axioms 1-5 over lotteries over trajectories.

- Then, utility has a local recursive form:

$$u(t \cdot h) = r(t) + \gamma(t)u(h), \quad u(\varepsilon) = 0.$$

- Preferences over lotteries are still represented by expected utility.

This establishes a bridge from expected utility maximisation over whole-trajectory utility to RL-style expected sum of future reward maximisation.

# Unrolling the recursion

- For  $h = (t_1, t_2, \dots, t_n)$ :

$$\begin{aligned} u(h) &= r(t_1) + \gamma(t_1)r(t_2) \\ &\quad + \gamma(t_1)\gamma(t_2)r(t_3) + \dots \\ &\quad + \left( \prod_{i=1}^{n-1} \gamma(t_i) \right) r(t_n). \end{aligned}$$

# Unrolling the recursion

- For  $h = (t_1, t_2, \dots, t_n)$ :


$$\begin{aligned} u(h) &= r(t_1) + \gamma(t_1)r(t_2) \\ &\quad + \gamma(t_1)\gamma(t_2)r(t_3) + \dots \\ &\quad + \left( \prod_{i=1}^{n-1} \gamma(t_i) \right) r(t_n). \end{aligned}$$

- If  $\gamma(t) \equiv \gamma$ , this becomes standard discounted return:

$$u(h) = \sum_{k=1}^n \gamma^{k-1} r(t_k).$$



# Summary: Preference, utility, and reward



# Open questions regarding AI alignment

- Will powerful AIs necessarily become expected utility maximisers?

# Open questions regarding AI alignment

- Will powerful AIs necessarily become expected utility maximisers?
- Instead of 'given a rewards, how to optimise it' we could ask 'what kind of properties we want our reward to satisfy'. For example, Are any of these properties (or their negation) useful for fostering:
  - Corregibility? (e.g. incompleteness)

# Open questions regarding AI alignment

- Will powerful AIs necessarily become expected utility maximisers?
- Instead of 'given a rewards, how to optimise it' we could ask 'what kind of properties we want our reward to satisfy'. For example, Are any of these properties (or their negation) useful for fostering:
  - Corregibility? (e.g. incompleteness)
  - Robust values? (e.g. continuity)

# Open questions regarding AI alignment

- Will powerful AIs necessarily become expected utility maximisers?
- Instead of 'given a rewards, how to optimise it' we could ask 'what kind of properties we want our reward to satisfy'. For example, Are any of these properties (or their negation) useful for fostering:
  - Corregibility? (e.g. incompleteness)
  - Robust values? (e.g. continuity)
  - Something else?
- Is RLHF providing an ordinal or cardinal utility signal? Are there implications of this?

# Open questions regarding AI alignment

- Will powerful AIs necessarily become expected utility maximisers?
- Instead of 'given a rewards, how to optimise it' we could ask 'what kind of properties we want our reward to satisfy'. For example, Are any of these properties (or their negation) useful for fostering:
  - Corregibility? (e.g. incompleteness)
  - Robust values? (e.g. continuity)
  - Something else?
- Is RLHF providing an ordinal or cardinal utility signal? Are there implications of this?